

## Comparison of the Microbial Quality of Traditional Ice Cream Produced by Small-Scale Manufacturers in *Kazeroon* with the Iranian National Standard

<sup>1</sup>M.S.A. Ghasemi, <sup>2</sup>P. Azadnia and <sup>3</sup>A. Maghsoodi

<sup>1</sup>School of Veterinary Medicine, Shiraz University, Iran

<sup>2</sup>Department of Food Hygiene and Public Health, School of Veterinary Medicine, Islamic Azad University, Kazeroon Branch, Iran

<sup>3</sup>School of Veterinary Medicine, Islamic Azad University, Kazeroon Branch, Iran

**Abstract:** Since, ice-cream harbors many potent pathogens, its microbial quality has always been crucially important in public health. In this study, 50 samples of ice cream produced by 30 different small-scale traditional ice cream manufacturers in *Kazeroon* were studied for Total Bacterial Count (TBC), Enterobacteriaceae count, coagulase positive Staphylococcal count, *E. coli* search and *Salmonella* sp. search, in order to determine if the samples meet the Ice cream standard set by the Iranian National Standard Center. Out of total ice cream of *Kazeroon* 50 samples (100%), 50 samples (100%) and 18 samples (36%) exceeded standard value of mesophilic aerobic count, Enterobacteriaceae count and coagulase positive Staphylococcal count, respectively. Also, according to the results obtained from *E. coli* search, it was determined that 26% of ice-cream samples examined were not fit bacteriologically to the Iranian Ice-Cream Standard. However, non of the samples were found to be contaminated with *Salmonella* sp. In the case of coagulase positive Staphylococcal count, the microbial quality of grade 2 ice cream samples was poorer in comparison to grade 1 samples but these two grade samples did not show significant differences in other cases. However, this study shows that the overall microbial quality of traditional ice cream samples being sold in *Kazeroon* is poor.

**Key words:** Traditional ice cream, standard, *Kazeroon*, total bacterial count, coliform, *Staphylococcus*, *E. coli*, *Salmonella*

---

### INTRODUCTION

Ice cream is a frozen dairy food made by freezing a pasteurized mix. The mix is composed of a combination of milk products (milk, condensed milk, milk powder, cream), sugar, emulsifiers, stabilizers, flavouring and coloring agents. Ice cream is a palatable, nutritive and healthful food. Therefore, ice cream production has increased rapidly in recent years in many countries of the world (Kocak *et al.*, 1998). In Iran, the production of ice cream is made in two different ways: traditional methods and the industrial production. The traditional production method refers to the manufacturing of open/artisanal ice cream. This kind of ice cream is generally manufactured in small-scale production units, which do not follow completely a standard procedure for production of ice cream. As, ice cream can become contaminated, while adding some ingredients after pasteurization or by means of improper sanitation of the equipment and environment, many psychrotroph and psychrotolerant microorganisms like *Listeria monocytogens*, *Staphylococcus aureus*,

*Bacillus*, *Salmonella*, *Shigella*, *Streptococcus*, *Pseudomonas*, *Campylobacter*, *Brucella* sp. and coliform bacteria are generally present in this kind of product (Jay, 1992). Therefore, evaluation of the presence of microorganisms in these products can give us the information about the raw ingredients quality and the sanitary nature of the processing and packaging stages. Therefore, there are lots of surveys on the bacteriological quality of ice cream and related products (Massa *et al.*, 1989). The present research was conducted to determine the sanitary quality of traditionally produced ice cream in *Kazeroon* (a city of Fars province) and comparing it to the Iranian Ice Cream National Standard.

### MATERIALS AND METHODS

In this study, fifty samples of traditional ice cream collected randomly from 30 different local markets of *Kazeroon*. The markets which had hygienic and acceptable appearance (according to the Iranian Institute of Food Control and Inspection) were nominated grade 1

and those, which had poor appearance were called grade 2. The samples were collected in an ice box and immediately taken to the laboratory where they were kept at -20°C, until they were examined for the bacteriological quality. Ice cream samples were prepared according to the Iranian Institute of standard and Industrial studies, 2001a. Bacteriological examination included the total bacteria, Enterobacteriaceae and coagulase positive *Staphylococcus aureus* counts, *E. coli* and *Salmonella* sp. search were also, done according to the Iranian Institute of Standard and Industrial Studies (1991, 1994, 1995a, b; 2001b), respectively. SPSS software (chi-square ( $\chi^2$ ) and Fishers exact test) were employed to statistically evaluate the data.

### RESULTS AND DISCUSSION

Out of total 50 samples, 50 ice cream samples (100%) exceeded standard value of total mesophilic aerobic count, which is  $5 \times 10^5$  cfu g<sup>-1</sup> (Iranian Institute of Standard and Industrial Studies, 2001a). In case of total Enterobacteriaceae count, 50 samples (100%) crossed the standard value which is 100 cfu g<sup>-1</sup>. Similarly, 18 samples (36%) and 13 samples (26%) were found crossing the standard value of Staphylococcal count (100 cfu g<sup>-1</sup>) and *E. coli* (0 cfu g<sup>-1</sup>), respectively. However, none of the samples were found to be contaminated with *Salmonella* sp. The average counts for total mesophilic aerobic bacteria, Enterobacteriaceae and coagulase positive *Staphylococcus aureus* were found to be  $4.9 \times 10^6$ ,  $2.9 \times 10^4$  and  $1.55 \times 10^2$  cfu g<sup>-1</sup>, respectively. In the case of coagulase positive Staphylococcal count, the microbial quality of grade 2 ice cream samples was poorer in comparison to grade 1 samples ( $p < 0.05$ ) but these two grade samples did not show significant differences in other cases ( $p > 0.05$ ). However, this study shows that the overall microbial quality of traditional ice cream samples being sold in *Kazeroon* is poor.

The bacteriological status of the samples are presented in Table 1 and the comparison of the bacteriological status of the samples (cfu g<sup>-1</sup>) with the standard value is shown in Table 2.

Ice cream is a fairly complex food containing sugar, emulsifiers and fats. This investigation presents the current status of microbial quality of ice cream being sold in *Kazeroon*. From the total analyzed ice cream samples (n = 50), 100% exceeded standard value of mesophilic aerobic count, 100% crossed the standard value of total Enterobacteriaceae count, 36% were beyond the standard value of staphylococcal count and 26% were found to carry *E. coli*, but none of the samples were found to be contaminated with *Salmonella* sp. These results are generally similar to the findings of other researchers and can be supported by the research of Massa *et al.* (1989) and Tamminga *et al.* (1980). Microbial quality of ice cream examined in Phillipines by Orallo *et al.* (1999) and Bangladesh by M-E Elahi *et al.* (2002) also showed the more or less comparable results.

*Staphylococcus aureus* is found to survive longer in ice cream (Gogov *et al.*, 1984). Staphylococci might enter the milk products from food handlers which either suffering from acute pyogenic infections or being healthy carriers harboring the organisms in nose or throat. The presence of coagulase positive *Staphylococcus*, which is mainly *S. aureus* can lead to the staphylococcal intoxication as a result of the growth and toxin production of this organism. The presence of starch and protein can also encourage the toxin production of these organisms (Jay, 1992).

The survival rate of the organisms depended both on the specific peculiarities of the individual strains and on the composition of the ice cream mixture. Exceeding the standard value in terms of mesophilic aerobic count may be because of poor handling of retailers. However, in terms of total coliform its cutting of standard value indicates the presence of fecal contamination of ice cream.

Table 1: The bacteriological status of the traditional ice cream samples (cfu g<sup>-1</sup>)

Grade of samples	No. of samples	Total bacterial count		Enterobacteriaceae count		Staphylococcal count	
		SV*	Average	SV*	Average	SV*	Average
1	25	$5.0 \times 10^5$	$4.8 \times 10^6$	$1.0 \times 10^2$	$2.8 \times 10^4$	$1.0 \times 10^2$	$1.0 \times 10^2$
2	25	-	$5.0 \times 10^6$	-	$3.0 \times 10^4$	-	$2.1 \times 10^2$

\*Standard value

Table 2: Comparison of the bacteriological status of the traditional ice cream samples (cfu g<sup>-1</sup>) with the standard value

Grade of samples	No. of samples	TBC*	Enterobacteriaceae	Staphylococcus	<i>E. coli</i>	<i>Salmonella</i>
1	25 no. of samples exceeded standard	25	25	4	7	0
	Samples exceeded standard (%)	100	100	16	28	0
2	25 no. of samples exceeded standard	25	25	14	6	0
	Samples exceeded standard (%)	100	100	56	24	0

\*TBC: Total Bacterial Count

This suggests, the possibility of the presence of other intestinal pathogens in this dessert. These microorganisms are transmitted via fecal-oral route.

The results show a negligence such as poor sanitation during the preparation or storage of ice cream including dirty premises, using bare hands in preparing the products, etc. Even, the raw milk could be a possible source of contamination. In the manufacturing processes, proper pasteurization is effective in destroying most of pathogenic bacteria. Automatic machines that are widely used for ice-cream production in dairy industry minimize direct hand manipulation and possibility of cross contamination.

The present study shows the sub-standard microbiological quality of open/artisanal ice cream and encourage the consumers to shift to the packed/industrial ice cream with consistently good quality. The study calls for action in two fold:

- Production permissions for artisanal ice cream manufacturing should be given only after a proper inspection against set standards. Periodical controls also, should be conducted to control this kind of production
- Consumers should be made aware of the non-hygienic quality of traditional ice cream and hygienic and other quality parameters of industrial ice cream

#### ACKNOWLEDGEMENT

We acknowledge the Islamic Azad University, Kazeroon Branch who technically supported this project.

#### REFERENCES

Gogov, I., G. Slavchev and T. Peeva, 1984. Cold resistance of *S. aureus* and Staphylococcal enterotoxins A and C2 in ice cream. Vet. Med. Nauki, 21: 46-50. www.ncbi.nlm.nih.gov/pubmed/6531859.  
Iranian Institute of Standard and Industrial Studies, 2001a. Microbial Examination of Milk Ice Cream, No. 2450.

Iranian Institute of Standard and Industrial Studies, 2001b. Total count of microorganisms at 30°C, No. 5272.  
Iranian Institute of Standard and Industrial Studies, 1995a. Enterobacteriaceae (Identification and counting), No. 2461.  
Iranian Institute of Standard and Industrial Studies, 1995b. Coagulase positive *Staphylococcus aureus* (Identification and counting in food), No. 1194.  
Iranian Institute of Standard and Industrial Studies, 1994. *E. coli* (Isolation, Identification and Most-Probable-Number in food), No. 2946.  
Iranian Institute of Standard and Industrial Studies, 1991. *Salmonella* sp. (Search and Identification), No. 1810.  
Jay, J.M., 1992. Modern Food Microbiology. 4th Edn. Chapman and Hall Inc. New York. ISBN: 0442007337.  
Kocak, C., M. Akan and H. Yardimci, 1998. Bacteriological quality of ice cream marketed in Ankara. Ankara Univ. Vet. Fak. Derg., 45: 131-134. dergiler.ankara.edu.tr/dergiler/11/559/6919.pdf.  
Massa, S., G. Poda, D. Cesaroni and L.D. Trovatielli, 1989. A bacteriological survey of retail ice cream. Food Microbiol., 6: 129-134. DOI: 10.1016/S0740-0020(89)80020-6. linkinghub.elsevier.com/retrieve/pii/S0740002089800206.  
M-E Elahi, A. T.M., S. Habib, M.M. Rahman, G.I. Rahman and M.J.U. Bhuiyan, 2002. Sanitary quality of commercially produced ice cream sold in retail stores. Pak. J. Nutr., 1: 93-94. DOI: 10.3923/pjn.2002.93.94. scialert.net/abstract/?doi=pjn.2002.93.94.  
Orallo, G.O., A.H. Pangan and E.C. Cabrera, 1999. Microbiological analysis of Ice cream produced by big scale and small scale manufactures in Metro Manila. Phil. J. Microbiol. Infect. Dis., 28: 99-101. www.psmid.org.ph/vol28/vol28num3.topic4.pdf.  
Tamminga, S.K., R.R. Beumer and E.H. Kampelmaeber, 1980. Bacteriological examination of ice-cream in the Netherlands: Comparative studies on methods. J. Applied Bacteriol., 49: 239-253. DOI: doi.wiley.com/10.1111/j.1365-2672.1980.tb05122.x www.ncbi.nlm.nih.gov/pubmed/6783602.