

## Effects of Subinhibitory Concentrations of Antibiotics and Antibodies on the Adherence of *Escherichia coli* to Human Uroepithelial Cells *In vitro*

<sup>1</sup>Maryam Jahanshahi, <sup>2</sup>Saeed Azad, <sup>2</sup>Bahram Aslanbeigi and <sup>3</sup>Mohammad Rahbar

<sup>1</sup>Department of Microbiology, Faculty of Medicine, Tarbiat Modarres University, Tehran, Iran

<sup>2</sup>Department of Pathology, Khatamol Anbya Hospital, Tehran, Iran

<sup>3</sup>Department of Microbiology, Iranian Reference Health Laboratory, Tehran, Iran

---

**Abstract:** Uropathogenic *Escherichia coli* is an important cause of urinary tract infection. Adherence to uroepithelial cells is a first step for colonization of bacteria. The aim of this study was to determine effects of sub Minimum Inhibitory Concentrations (MICs) of antibiotics and antibodies on the adherence of this organism at *in vitro* condition. Seven strains of *E. coli* isolated from patients with acute pyelonephritis were subject of the study. MICs of trimethoprim, sulphamethoxazole, sulphadiazine and ampicillin for these strains were determined and effect of these antibiotics at the ½ and ¼ MIC on the adhesion of *E. coli* to human urinary tract epithelial cells were studied. Trimethoprim, sulphamethoxazole and sulphadiazine at ½ and ¼ of the MICs decreased the adherence of four out of five *E. coli* strains tested whereas combinations of the compounds did not potentiate the effect. Ampicillin caused a similar effect. Specific pili-antibodies as well as gamma globulin and milk inhibited the adherence but did not work synergistically with ampicillin. In conclusion, present preliminary investigation gives further support to the theory that the effect of sub inhibitory concentration of antibiotics on the growth of bacteria may prevent their attachment to urinary epithelial cells.

**Key words:** Sub MIC, uropathogenic *E. coli*, uroepithelial cells, ampicillin, trimethoprim, sulphadiazine, pili-antibodies

---

### INTRODUCTION

Association between bacteria and mucosal surfaces is considered necessary for colonization (Jones, 1977). The microorganism may bind to different constituents of the mucosa. Binding to epithelial cells is defined as attachment ability to selectively attach to different epithelial surfaces is a determinant of the microflora in each ecological niche (Ellen and Gibbons, 1974). Adhesive capacity may also be essential for the pathogenesis of infection (Jones, 1977; Ellen and Gibbons, 1974; Svanborg-Eden *et al.*, 1978a). as one of several virulence factors co-appearing on the infection strain. The biochemical basis of the adhesion process has been defined most extensively for *E. coli* strains causing Urinary Tract Infection (UTI) and possessing the ability to adhere to human urinary tract epithelial cells (Svanborg-Eden *et al.*, 1979) and erythrocytes (Kallenius *et al.*, 1980). *Bacterial pili* (fimberiae) have been found to bind to glycosphingolipid receptors on the membranes of epithelial cells (Leffler and Svanborg-Eden, 1980; Kallenius *et al.*, 1980). Such ligand mediated adherence of the bacteria can be prevented by interference either with the pili or with the epithelial receptor structure. Specific

bacterial antibodies can bring about an inhibition of the binding. On the mucous membranes particularly secretory IgA (SIgA) but also IgG, possess such activities (Svanborg-Eden *et al.*, 1978a). Analogues of the glycolipid receptor can inhibit attachment. There have previously demonstrated that sub-Inhibitory Concentrations (sub-MICs) of ampicillin and amoxycillin possess adherence-decreasing activities *in vitro* (Sandberg *et al.*, 1979; Svanborg-Eden *et al.*, 1978b; Vidya *et al.*, 2005; Loudbeyre *et al.*, 1993; Wojnicz and Jankowski, 2007). The aim of the present study was to investigate the effect of sub MICs of trimethoprim, sulphadiazine and sulphamethoxazole alone or in combination on the attachment of *E. coli* to uroepithelial cells. Furthermore, antibodies to isolated pili were used to evaluate the effect of ampicillin on the piliation of bacteria.

### MATERIALS AND METHODS

*E. coli* strains possessing good adhesive capacity isolated from the urine of patients with acute urinary tract infection were used. The bacteria, kept in deep agar storage cultures were transferred to lactose-bromothymol blue agar plates.

**Antibiotics and determination of MICs:** Trimethoprim, sulphamethoxazole, sulphadiazine and ampicillin were used. The Minimum Inhibitory Concentration (MIC) of each antibacterial agent for each *E. coli* strain was determined under conditions identical to those employed for growth of bacteria in the adhesion test system. Serial twofold dilutions of the compounds in antibiotic sensitive medium were inoculated with  $10^5$  bacteria from a 4 h broth culture. After incubation for 24 h at 37°C, the lowest concentration of antibiotic without visible turbidity was recorded and considered as the MIC.

The range of MICs of trimethoprim for the seven strains of *E. coli* tested was 0.3-1.25  $\mu\text{g mL}^{-1}$ , of sulphamethoxazole 50-600  $\mu\text{g mL}^{-1}$  and of sulphadiazine 100-600  $\mu\text{g mL}^{-1}$ .

**Adhesion testing:** Adhesion testing was performed as previously described. Uroepithelial cells were obtained from the sediment of fresh urine from a non-bacteriuric woman. The cells were washed, resuspended in phosphate-buffered saline (PBS; PH 7.1, 300 mosm/l) and quantitated by direct light microscopy using Burker chamber. To  $10^5$  epithelial cells were added  $10^8$  bacteria and PBS to total volume of 1 mL after incubation of bacteria and epithelial cells for 60 min, the number of bacteria attached to the cells was counted under a light microscope. Adhesion was expressed as the mean number of bacteria attached to 40 epithelial cells. In each experiment antibiotic-treated bacteria and identically handled controls were compared (Korhonen *et al.*, 1980).

**Treatment of bacteria with subinhibitory concentrations of antibiotics:** Parts of one bacterial colony were transferred from a lactosebromothymol blue agar plate to 3 mL of ASM broth. After 2 h growth at 37 °C without shaking, the antibacterial agent suspended in ASM broth was added to a final concentration of one-half or one-fourth of the MIC. After incubation for 4 h at 37°C, the bacteria were harvested by centrifugation and used for adhesion testing as described above.

**Viable counts:** The number of bacteria registered by viable counts in the adhesion test tube agreed well with

the number counted in the Burker chamber by direct light microscopy (Loubeyre *et al.*, 1993; Wojnicz and Jankowski, 2007; Korhonen *et al.*, 1980).

**Ampicillin and antibodies:** One strain of *E. coli* with good adhesive properties, *E. coli* 3048, untreated or treated with Ampicillin at 1-8 of the MIC was used. The MIC of ampicillin for this strain was 6.25  $\mu\text{g mL}^{-1}$ . Purified pili were obtained from the same strain as recently described by Korhonen *et al.* (1980) and Svanborg-Eden *et al.* (1976). Antiserum to the purified pili was obtained from immunized rabbits. The antiserum was specific for the pili and did not react with the O or K antigen of the strain.

After preincubation of bacteria for 30 min with 0.001, 0.01, 0.05 and 0.1 mL of anti-pili serum, epithelial cells and PBS were added to a total volume of 1 mL and adhesion testing was completed as described above. Bacteria exposed to preimmune serum served as a control. Commercial gamma globulin and pool of human breast-milk served as sources of IgG and IgA antibodies (Svanborg-Eden and Svennerholm, 1978).

## RESULTS

All concentrations of one-half and one-fourth of their MICs, trimethoprim, sulphamethoxazole and sulphadiazine decreased the adhesive capacity of 6-7 *E. coli* strains tested.

The combination of sub-MICs of trimethoprim and either of the sulphonamides did not decrease adherence further compared with each compound alone Table 1. Bacteria exposed to sub-MICs of ampicillin but not of the other antibacterial agents, became elongated, as studied by direct microscopy. The majority of the bacteria that retained their adhesive capacity in presence of ampicillin were not elongated.

Specific antipili antibodies (as well as gamma globulin and milk (data not shown)) inhibited adherence in a dose-response relationship. Pretreatment of bacteria with ampicillin at 1-8 of the MIC diminished the adherence

Table 1: Effect of sub-MICs of trimethoprim, sulphamethoxazole and sulphadiazine alone and in combination on the adherence of seven strains of *E. coli* to human uroepithelial cells. Estimated as the mean number of bacteria on 40 epithelial cells

<i>E. coli</i> strain	Control	Trimethoprim		Sulphameth-oxazole		Sulphadi-azine		Trimethoprim+ Sulphamethoxazole 1/4+ 1/4	Trimethop rim+ Sulph adiazine 1/4+ 1/4
		1/2	1/4	1/2	1/4	1/2	1/4		
1	45	13	13	12	24	38	35	19	25
2	25	0	8	9	16	5	4	0	0
3	25	0	9	15	11	6	0	20	11
4	20	16	14	23	16	15	20	18	28
5	98	6	35	23	100	51	57	15	30
6	33	143	12	0	3	14	14	3	0
7	70	39	28	28	19	35	35	14	4




Fig. 1: Effect of anti-pili antibodies in hyperimmune rabbit serum on the attachment to human uroepithelial cells of *E. coli* 3048, either untreated (×) or treated (O) with one-eighth of the MIC of Ampicillin

further Fig. 1. No decrease in attachment could be demonstrated when bacteria were incubated with preimmune serum only.

### DISCUSSION

The ability of *E. coli* to attach to human urinary tract epithelial cells *in vitro* has been found to correlate with the type of UTI caused by the strains (Korhonen *et al.*, 1980; Svanborg-Eden *et al.*, 1976; Deschner, 1976). Thus, *E. coli* isolated from patients with acute pyelonephritis and acute cystitis adhere more efficiently than strains from patients with asymptomatic bacteriuria (Svanborg-Eden *et al.*, 1976).

We have shown before that sub-MICs of ampicillin and amoxicillin suppress bacterial adherence to uroepithelial cells (Svanborg-Eden *et al.*, 1978a, b; Sandberg *et al.*, 1979). The underlying mechanism explaining this effect may be a distortion of the bacterial cell wall synthesis producing elongated forms of resulting in a lack of adhesion-mediating structures on the surface of the bacteria.

In the present study, using the same *In vitro* adherence model, one-half and one-fourth of the MIC of trimethoprim, sulphadiazine and sulphamethoxazole suppressed the adhesive ability of all except one of the *E. coli* strains tested. Both trimethoprim and sulphonamides are inhibitors of bacterial folate metabolism resulting in disturbances of DNA and RNA synthesis. This may cause deficient production of structures, possibly pili, involved in the adherence process.

Bacteria exposed to sub-MICs of ampicillin and nitrofurantoin became elongated but only ampicillin decreased attachment. This suggests that the altered morphology observed, per se is not responsible for the altered adhesive properties of the bacteria but may be due to unrelated injuries either within or on the microbial cells (Klainer and Perkins, 1972; Smellie *et al.*, 1976).

The attachment of *E. coli* to human uroepithelial cells is inhibited by commercial gamma globulin, human breast-milk containing SigA antibody and IgG and SIgA fractions of urine from patients with acute pyelonephritis (Stamey and Condy, 1975). These antibodies are probably directed against various bacterial surface structures (Svanborg-Eden *et al.*, 1978a). The adherence decreasing effect of commercial gamma globulin seems to be potentiated by sub-MICs of ampicillin (Svanborg-Eden *et al.*, 1978b).

In this study, specific anti-pili antibodies decreased attachment in a dose-response relationship. A synergistic effect between ampicillin and anti-pili antibodies could not be demonstrated but the effects were additive. This finding may be explained by a lower density of pili present on bacteria after ampicillin treatment.

The clinical impact of these *in vitro* findings has not yet been elucidated. Excretion of small amounts of antimicrobials in urine or vaginal fluid can be obtained by low-dose prophylaxis (Cattel *et al.*, 1976; Stamey *et al.*, 1978). A marked reduction in *E. coli* carriage both in the stool and especially on the periurethral area, persisting over months has been demonstrated in patients on long-term low-dose prophylaxis with trimethoprim-sulphamethoxazole (Stamey *et al.*, 1978).

Furthermore, a high percentage of women not prone to UTI and seldom colonized with enterobacteria periurethrally had Cervicovaginal Antibodies (CVA) against their predominant faecal strain of *E. coli* in contrast, a minority of non-bacteriuric, UTI-susceptible subjects had CVA against enterobacteriacease colonizing the introitus.

This suggests that local antibody is one important determinant of bacterial adherence to periurethral cells, possibly by blocking bacterial adhesions (Cattel *et al.*, 1976; Stamey *et al.*, 1978). UTI-prone individuals lacking specific CVA may avoid being colonized with potentially uropathogenic bacteria by receiving low-dose prophylaxis with antimicrobial drugs.

### CONCLUSION

The present preliminary investigation gives further support to the theory that the effect of sub inhibitory concentration of antibiotics on the growth of bacteria may prevent their attachment to urinary epithelial cells.

### ACKNOWLEDGEMENT

The researchers are grateful to Khatamolanbia Hospital for financial support of this study. The researchers declare that there is no conflict of interest.

### REFERENCES

- Cattel, W.R., M.A. McSherry, H.L. Brooks and F.W. O'Grady, 1976. The carriage of *Escherichia coli* on the peri-urethral area and in the faces in patients on long-term low-dose co-trimoxazole therapy. *Clin. Nephrol.*, 6: 506-508.
- Deschner, F., 1976. Inhibition of cell wall synthesis by sulfonamids and trimethoprim. *Chemotherapy*, 22: 12-18.
- Ellen, R.P. and R.J. Gibbons, 1974. Parameters affecting the adherence and tissue tropism of *Streptococcus pyogenes*. *Infect. Immun.*, 9: 85-91.
- Jones, G.W., 1977. The Attachment of the Bacteria to the Surfaces of Animal Cells. In: *Receptors and Recognition Ser. B*, Reissig, J.L. (Ed.). Chapman and Hall, London, pp: 133-176.
- Kallenius, G., R. Mollby, S.B. Svenson, J. Winberg and H. Hultberg, 1980. Identification of a carbohydrate receptor recognized by uropathogenic *Escherichia coli*. *Infect. Suppl.*, 8: S288-S293.
- Klainer, A.S. and R.L. Perkins, 1972. Surface manifestations of antibiotic-induced alterations in protein synthesis in bacterial cells. *Antimicrob. Agents Chemother.*, 1: 164-170.
- Korhonen, T.K., E.L. Nurmiäho and C. Svanborg-Eden, 1980. A new method for the purification of *Escherichia coli* pili. *Infect. Immun.*, 27: 569-575.
- Leffler, H. and Svanborg-Eden, 1980. Chemical identification of a glycosphingolipid receptor for *Escherichia coli* attaching to human urinary tract epithelial cells and agglutinating human erythrocytes. *FEMS Microbiol. Lett.*, 8: 127-134.
- Loubeyre, C., J.F. Desnottes and N. Moreau, 1993. Influence of sub inhibitory concentrations of antibacterial on the surface properties and adhesion of *E. coli*. *J. Antimicrob. Chemotherapy*, 31: 37-45.
- Sandberg, T., K. Stenqvist and C. Svanborg-Eden, 1979. Effects of subminimal inhibitory concentrations of ampicillin, chloramphenicol and nitrofurantoin on the attachment of *Escherichia coli* to human uroepithelial cells in vitro. *Rev. Infect. Dis.*, 1: 838-844.
- Smellie, J.M., R.N. Grunberg, A. Leakey and W.S. Atkin, 1976. Long-term low-dose co-trimoxazole in prophylaxis of childhood urinary tract-infection: Clinical aspects. *Br. Med. J.*, 2: 203-206.
- Stamey, T.A. and M. Condy, 1975. The diffusion and concentration of trimethoprim in human vaginal fluid. *J. Infect. Dis.*, 131: 261-266.
- Stamey, T.A., N. Wehner, G. Mihara and M. Condy, 1978. The immunologic basis of recurrent bacteriuria: Role of cervicovaginal antibody in enterobacterial colonization of the introital mucosa. *Medicine*, 57: 47-56.
- Svanborg-Eden, C., L.A. Hanson, U. Jodal, U. Lindberg and A. Sohl Akerlund, 1976. Variable adherence to normal human urinary epithelial cells of *Escherichia coli* strains associated with various forms of urinary tract infection. *Lancet II*, 308: 490-492.
- Svanborg-Eden, C. and A.M. Svennerholm, 1978. Secretory immunoglobulin A and G antibodies prevent adhesion of *Escherichia coli* to human urinary tract epithelial cells. *Infect. Immun.*, 22: 790-797.
- Svanborg-Eden, C., T. Sandberg, K. Stenqvist and S. Ahlstedt, 1978a. Decrease in adhesion of *Escherichia coli* to human urinary tract epithelial cells *in vitro* by sub-inhibitory concentrations of ampicillin. A preliminary study. *Infection*, 6: S121-S124.
- Svanborg-Eden, C., B. Eriksson, L.A. Hanson, U. Jodal and B. Kaijser *et al.*, 1978b. Adhesion to normal human uroepithelial cells of *Escherichia coli* from children with various forms of urinary tract infection. *J. Pediatre*, 93: 398-403.
- Svanborg-Eden, C., T. Sandberg, K. Stenqvist and S. Ahlstedt, 1979. Effects of sub-inhibitory amounts of ampicillin, amoxicillin and mecillinam on the adhesion of *Escherichia coli* bacteria to human urinary tract epithelial cells: A preliminary study. *Infection*, 7: S452-S455.
- Vidya, K.C., P.S. Mallya and P.S. Rao, 2005. Inhibition of bacterial adhesion by sub inhibitory concentrations of antibiotics. *Indian J. Med. Microbiol.*, 23: 102-105.
- Wojnicz, D. and S. Jankowski, 2007. Effects of subinhibitory concentrations of amikacin and ciprofloxacin on adherence to epithelial cells of uropathogenic *E. coli* strains. *Int. J. Antimicrob. Agents*, 29: 700-704.