

## Revision of the Fish Fauna of the Seyhan and Ceyhan River Basins in Turkey

Fusun Erk'akan and Filiz Ozdemir

Department of Biology, Faculty of Science, Beytepe Campus, Hacettepe University, Ankara, Turkey

---

**Abstract:** This study conducted between April 2008 to March 2009, aimed revision of ichthyofauna of the Seyhan and Ceyhan river Basins, determined that 35 species and one sub-species belonging to *Anguillidae*, *Salmonidae*, *Cyprinidae*, *Nemacheilidae*, *Cobitidae*, *Clariidae*, *Siluridae*, *Cyprinodontidae*, *Poeciliidae*, *Blenniidae* and *Percidae*. In this study, *Alburnus baliki* and *Cobitis levantina* was recorded for the 1st time in the Ceyhan river and *Nemacheilus namiri* in the Seyhan river Basins. Samples of *Capoeta turani* described in the Seyhan river Basin and *Capoeta erhani* described in the Ceyhan river Basin have been caught in large quantities and examined in different locations in these Basins and these species have been accepted as a synonym of *Capoeta barroisi*.

**Key words:** Ichthyofauna, *Alburnus baliki*, *Cobitis levantina*, *Nemacheilus namiri*, *Capoeta barroisi*, Seyhan-Ceyhan river Basins

---

### INTRODUCTION

Turkey is geologically active from the Miosen to the present and many fish genera speciated in this area. In recent years, taxonomic and systematic studies on Turkish Freshwater fishes have been investigated by many researchers and some descriptions have been revised (Erk'akan, 1983a, b; Erk'akan and Kuru, 1983; Erk'akan, 1984; Krupp, 1985; Balik, 1985; Erk'akan and Kuru, 1986a, b; Coad and Sarieyyupoglu, 1988; Krupp and Schneider, 1991; Bogustkaya, 1992, 1997; Ahnelt, 1995; Ahnelt *et al.*, 1995; Ahnelt and Holcik, 1996; Elvira, 1997; Wildekamp *et al.*, 1997, 1999; Erk'akan *et al.*, 1998, 2007, 2008a, b; Bogustkaya *et al.*, 2000, 2006; Freyhof and Ozulug, 2006; Turan *et al.*, 2006, 2008, 2009; Kucuk *et al.*, 2007; Ozulug and Freyhof, 2007; 2008).

Seyhan and Ceyhan river Basins have interesting fish fauna because these basins are junction for some fish species which are also living in the geographically close basins such as Firat from the East, Asi from the South and Manavgat and Tarsus from the West.

Balik (1985) in his study entitled; a taxonomical revision of freshwater fish in South Anatolia, investigated the distribution of some fish species in the Seyhan and Ceyhan rivers.

Kara *et al.* (2010) were established *Anguilla anguilla*, *Salmo trutta macrostigma*, *Cyprinus carpio*, *Achanthobrama* sp., *Alburnus orontis*, *Pseudophoxinus zekayi*, *Squalius kottelati*, *Garra rufa*, *Chondrostoma regium*, *Luciobarbus pectoralis*, *Capoeta angorae*, *Capoeta erhani*, *Cobitis evreni*, *Schistura*

*ceyhanensis*, *Oxynemacheilus* sp., *Silurus glanis*, *Clarias gariepinus*, *Aphanius mento*, *Gambusia affinis*, *Salaria fluviatilis* (*Blennius fluviatilis* ASSO, 1801) in the Seyhan and Ceyhan river Basins.

According to Alagoz, *Acanthalburnus microlepis*, *Capoeta capoeta capoeta*, *Capoeta capoeta umbla*, *Barbus esocinus*, *Rutilus rutilus*, *Rutilus rubilio*, *Rutilus frisii*, *Aphanius chanteri*, *Abramis brama* and *Cobitis bilseli* were found in the Seyhan dam reservoir.

Bostanci in his study entitled; a systematic investigation on the fish fauna of the Seyhan, Ceyhan and Asi rivers determined 37 species and one sub-species belonging to fourteen genus. *C. erhani* by Turan *et al.* (2008) and *C. turani* by Ozulug and Freyhof (2008) were recently described from in the Ceyhan and Seyhan Basins, respectively. However, the small number of samples and without examination of serial material studied by some researchers has prevented observation of the variations of some species and this situation has caused confusion in the freshwater fish fauna in Turkey.

The aim of this study is to eliminate any confusion in the literature by determining species in the Seyhan and Ceyhan Basins involving a number of researchers and revealing deficiencies and mistakes in species descriptions.

### MATERIALS AND METHODS

Employing electroshocker, fish samples were collected between April 2008 to March 2009 from the determined stations on the water body of the Seyhan and

Table 1: Sampling stations in the Seyhan river Basin

Location	Coordinates	Elevation (m)
Seyhan dam reservoir	37°02.031'N; 35°20.029'E	67
Cakit stream I; Pozanti-Adana	37°30.898'N; 34°44.675'E	977
Cakit stream II-Erminlik-Ciftehan	37°30.090'N; 34°49.069'E	881
Camardi stream, Kamisli	37°33.140'N; 34°57.282'E	1112
Pozanti	37°27.828'N; 34°52.228'E	955
Delicay-Adana	37°02.355'N; 34°30.304'E	26,5
Eglence stream-Adana	37°17.499'N; 35°13.412'E	142
Cerkes-Karabogaz-Zamanti stream	38°46.526'N; 36°27.248'E	1500
Gumusgun stream-Pinarbasi-Kahramanmaraş road	38°43.242'N; 36°24.846'E	1453
Buyukgumusgun-Sariz road	38°41.070'N; 36°27.406'E	1650
20 km to Sariz village	38°34.942'N; 36°25.831'E	1740
Goksun-Elbistan road	38°27.507'N; 36°28.307'E	1710
Sariz stream	38°26.924'N; 36°28.532'E	1770
Yamanli village-Tufanbeyli road	38°12.967'N; 36°14.016'E	1620
Tufanbeyli-Adana road	38°12.802'N; 36°14.174'E	1632
Cataloluk stream-Bakirdag	38°12.154'N; 35°50.879'E	1740
Korkun stream-Karaisali-Adana	37°13.095'N; 35°09.100'E	145
Feke-Saimbeyli road	37°51.941'N; 36°03.502'E	560
Degirmendere, Feke-Mansurlu road	37°50.573'N; 35°45.403'E	928
Feke	37°51.537'N; 35°51.490'E	970
Feke Exit-Kozan road	37°46.952'N; 35°54.343'E	515

Table 2: Sampling stations in the Ceyhan river Basin

Location	Coordinates	Elevation (m)
Carnizagili (Avcilar bridge) Kadirli road	37°14.918'N; 35°50.313'E	36.2
Hacilar-Ceyhan Kadirli road	37°18.900'N; 35°53.324'E	35.8
Davutlar village	37°21.318'N; 35°56.396'E	32.7
Kadirli village	37°21.952'N; 36°06.355'E	86.4
Andirin, Celik stream	37°30.730'N; 36°21.507'E	1047.0
Hemite Creek Tecirli-Osmaniye	37°11.154'N; 36°05.138'E	53.4
Sabun stream-Osmaniye-Duzici	37°16.898'N; 36°28.030'E	289.0
Terbuzek-I stream, Goksun	38°04.155'N; 36°27.912'E	1350.0
Terbuzek-II stream, Goksun	38°01.917'N; 36°28.336'E	1344.0
Komur stream-Goksun exit	38°00.524'N; 36°30.704'E	1319.0
Kizilcik stream-Koture village-Afsin road	38°11.701'N; 36°50.620'E	1320.0
Sogutlu Creek-Elbistan	38°13.411'N; 37°13.386'E	1280.0
Yalak stream-Sevilli village-Elbistan	38°15.271'N; 37°31.984'E	1600.0
Korsulu stream-Andirin road	37°38.218'N; 36°37.548'E	757.0
Gecit stream-Andirin road	37°38.320'N; 36°39.969'E	687.0
Firiz stream-Tekir road	37°45.552'N; 36°41.823'E	960.0
Tekir Creek-Dongel cave	37°51.590'N; 36°38.508'E	1248.0

Ceyhan Basin which reflected varied ecological characteristics. Requisite measurements for the taxonomic studies of fish were made with a millimetric ruler and digital calipers with 0.01 mm precision based on the method of Banarescu (1991). Comparison of similarities and differences in the metric and meristic characteristics between fish belonging to the same species caught in the study and fish samples belonging to the same species caught in others parts of Turkey was made by Prof. Dr. Fusun Erk'akan. Coordinates and elevation of the sampling stations were given for the Seyhan river Basin in Table 1 and for the Ceyhan river Basin in Table 2.

## RESULTS

This study aimed to revise the ihtiyofauna of the Seyhan and Ceyhan river Basins. About 29 species and one sub-species belonging to nine families were

determined in the Seyhan river Basin. Of these, 8 species are endemic and 5 species are exotic (Table 3). Total 28 species and 1 sub-species belonging to ten families were determined in the Ceyhan Basin. Of these, 6 species are endemic and 2 species are exotic (Table 4).

### List of fish species found in the Seyhan and Ceyhan river Basins:

Kingdom: Animalia  
 Phylum: Chordata  
 Subphylum: Vertebrata (Craniata)  
 Infraphylum: Gnathostomata  
 Class: Actinopterygii  
 Subclass: Neopterygii  
 Infraclass: Teleostei  
 Superorder: Elopomorpha  
 Order: Anguilliformes  
 Subclass: Anguilloidei  
 Family: Anguillidae  
 Genus: *Anguilla* (Linnaeus, 1758)  
 Species: *Anguilla anguilla* (Linnaeus, 1758)  
 Superorder: Protacanthopterygii

Table 3: The fish fauna of the Seyhan river Basin

Family	Species	Localities	Endemism
Salmonidae	<i>Salmo trutta macrostigma</i> (Dumeril, 1858)	Pinarbasi stream	
	<i>Salmo platycephalus</i>	Soguksu; Pinarbasi-Zamanti, Karagoz;	
	<i>Oncorhynchus mykiss</i> (Walbaum, 1792)	Buyukgumusgun; Sariz	Endemic
Percidae	<i>Sander lucioperca</i> (Linnaeus, 1758)	Seyhan dam reservoir	Exotic
Blennidae	<i>Salaria fluviatilis</i> (Asso, 1801)	Seyhan dam reservoir	Exotic
Cyprinidae	<i>Acanthobrama marmid</i> (Heckel, 1843)	Eglençe stream; Korkun stream	
	<i>Alburnus orontis</i> (Sauvage, 1882)	Seyhan dam reservoir	
	<i>Alburnus adanensis</i> (Battalgazi, 1944)	Seyhan dam reservoir; Feki; Korkun stream;	
	<i>Barbus pectoralis</i> (Heckel, 1843)	Saimbeyli-Goksu	Endemic
	<i>Barbus rajanorum</i> (Heckel, 1843)	Seyhan dam reservoir; Korkun stream; Feki;	
	<i>Capoeta damascina</i> (Valenciennes, 1842)	Eglençe stream	
	<i>Capoeta barroisi</i> (Lortet, 1894)	Seyhan dam reservoir; Zamanti stream;	
	<i>Chondrostoma regium</i> (Heckel, 1843)	Korkun stream	
	<i>Cyprinus carpio</i> (Linnaeus, 1758)	Seyhan dam reservoir; Eglençe stream;	
	<i>Carasius carassius</i> (Linnaeus, 1758)	Delicay	Exotic
	<i>Garra rufa</i> (Heckel, 1843)	Eglençe stream; Seyhan dam reservoir; Korkun	
	<i>Hemigrammacapoeta caudomaculata</i> (Battalgil, 1942)	stream; Feki; Saimbeyli-Goksu	Endemic
	<i>Pseudophoxinus zekayi</i> (Bogutskaya et al., 2006)	Korkun stream	Endemic
	<i>Squalius cephalus</i> (Linnaeus, 1758)	Eglençe stream;	
	<i>Squalius lepidus</i> (Heckel, 1843)	Seyhan dam reservoir; Eglençe stream;	
	<i>Tinca tinca</i> (Linnaeus, 1758)	Cerkes-Karabogaz;	
Siluridae	<i>Silurus glanis</i> (Linnaeus, 1758)	Gumusgun river; Sariz; Tufanbeyli-Yamanli	
Clariidae	<i>Clarias gariepinus</i> (Burchell, 1822)	Village; Cataloluk	
Cyprinodontidae	<i>Aphanius mento</i> (Heckel, 1843)	Stream-Bakirdag; Saimbeyli-Goksu	
	<i>Aphanius danfordii</i> (Boulenger, 1890)	Zamanti stream; Pinarbasi; Seyhan dam	
Poeciliidae	<i>Gambusia holbrooki</i> (Girard, 1859)	reservoir; Saimbeyli-Goksu	
Nemacheilidae	<i>Oxyemacheilus samanticus</i>	Seyhan dam reservoir; Korkun stream	Exotic
	<i>Paracobitis seyhanensis</i>	Pinarbasi; Bakirdag-Zamanti; Karabogaz-Cerkes;	Endemic
	<i>Paracobitis tigris</i> (Heckel, 1843)	Karabogaz-Cerkes;	
	<i>Schistura seyhanicola</i> (Erk'akan et al., 2007)	Eglençe stream; Karabogaz-Cerkes;	
	<i>Schistura evreni</i> (Erk'akan et al., 2007)	Korkun stream	Endemic
		Bakirdag-Zamanti; Karabogaz-Cerkes	Endemic

Table 4: The fish fauna of the Ceyhan river Basin

Family	Species	Localities	Endemism
Anguillidae	<i>Anguilla anguilla</i> (Linnaeus, 1758)	Tecirli village	
Salmonidae	<i>Salmo trutta macrostigma</i> (Dumeril, 1858)	Göksun stream; Sogutlu; hurman stream; Terbuzek I-II;	
Blennidae	<i>Salaria fluviatilis</i> (Asso, 1801)	Davutlar village; Sabun stream; Tecirli;	Exotic
Cyprinidae	<i>Alburnus baliki</i> (Bogutskaya et al., 2000)	Gecit stream; Aksu stream	
	<i>Alburnus orontis</i> (Sauvage, 1882)	Davutlar village	Endemic
	<i>Alburnus adanensis</i> (Battalgazi, 1944)	Davutlar village; Sabun stream; Tecirli; Goksun stream	
	<i>Barbus pectoralis</i> (Heckel, 1843)	Goksun stream; Aksu stream	
	<i>Barbus rajanorum</i> (Heckel, 1843)	Gecit Stream; Tecirli	
	<i>Capoeta damascina</i> (Valenciennes, 1842)	Camizagili stream; Aksu stream	
	<i>Capoeta barroisi</i> (Lortet, 1894)	Gecit stream; Aksu stream; Celik stream	
	<i>Chondrostoma regium</i> (Heckel, 1843)	Göksun stream; Sevdilli; Andirin; Firniz stream;	
	<i>Garra rufa</i> (Heckel, 1843)	Sabun stream Aksu stream	
	<i>Hemigrammacapoeta caudomaculata</i> (Battalgil, 1942)	Gecit stream; Aksu stream; Celik stream; Camizagili stream;	
	<i>Pseudophoxinus zekayi</i> (Bogutskaya et al., 2006)	Davutlar stream; Sabun stream; Tecirli; Goksun stream	
		Davutlar village; Tecirli stream	Endemic
		Davutlar village; Tecirli stream	Endemic

Table 4: Continued

Family	Species	Localities	Endemism
	<i>Squalius cephalus</i> (Linnaeus, 1758)	Celik stream; Andirin	
	<i>Squalius lepidus</i> (Heckel, 1843)	Aksu stream	
Clariida	<i>Clarias gariepinus</i> (Burchell, 1822)	Aksu stream, Davutlar village	
Cyprinodontidae	<i>Aphanius danfordii</i> (Boulenger, 1890)	Camizagili stream; Davutlar; Kadirli	
	<i>Aphanius mento</i> (Heckel, 1843)	Camizagili stream; Davutlar; Kadirli	
Poecilidae	<i>Gambusia holbrooki</i> (Girard, 1859)	Camizagili stream; Davutlar; Kadirli	Exotic
Nemacheilidae	<i>Paracobitis tigris</i> (Heckel, 1843)	Aksu stream; Celik stream	
	<i>Schistura evreni</i> (Erk'akan <i>et al.</i> , 2007)	Gecit stream; Aksu stream; Celik stream; Firmiz stream; Andirin; Tekir Creek	Endemic
	<i>Schistura ceyhanensis</i> (Erk'akan <i>et al.</i> , 2007)	Sevdilli village, Elbistan, Komur stream	Endemic
	<i>Schistura seyhanicola</i> (Erk'akan <i>et al.</i> , 2007)	Kesis stream	Endemic
Cobitidae	<i>Cobitis levantina</i> (Krupp and Moubayed, 1992)	Hemite Creek; Davutlar	

Order: Salmoniformes  
 Family: Salmonidae  
 Genus: *Salmo* (Linnaeus, 1758)  
 Species:  
 • *Salmo platycephalus* (Behnke, 1969)  
 • *Salmo trutta magrostitigma* (Dumeril, 1858)  
 • *Oncorhynchus mykiss* (Walbaum, 1792)  
 Superorder: Euteleostei  
 Superclass: Ostariophysii  
 Order: Cypriniformes  
 Family: Cyprinidae  
 Genus: *Acanthobrama* (Heckel, 1843)  
 Species: *Acanthobrama marmid* (Heckel, 1843)  
 Genus: *Alburnus* (Heckel, 1843)  
 Species:  
 • *Alburnus adanensis* (Battalgil, 1944)  
 • *Alburnus baliki* (Bogustkaya *et al.*, 2000)  
 • *Alburnus orontis* (Sauvage, 1882)  
 Genus: *Barbus* (Cuvier, 1817)  
 Species:  
 • *Luciobarbus pectoralis* (Heckel, 1843)  
 • *Barbus rajanorum* (Heckel, 1843)  
 Genus: *Capoeta* (Cuvier-Valenciennes, 1842)  
 Species:  
 • *Capoeta barroisi* (Lortet, 1894)  
 • *Capoeta damascina* (Valenciennes, 1842)  
 Genus: *Chondrostoma* (Agassiz, 1835)  
 Species: *Chondrostoma regium* (Heckel, 1843)  
 Genus: *Cyprinus* (Linnaeus, 1758)  
 Species: *Cyprinus carpio* (Linnaeus, 1758)  
 Genus: *Garra* (Heckel, 1843)  
 Species: *Garra rufa* (Heckel, 1843)  
 Genus: *Hemigrammocapoeta* (Pellegrin, 1927)  
 Species: *Hemigrammocapoeta culiciphaga* (Pellegrin, 1927)  
 Genus: *Pseudophoxinus* (Heckel, 1843)  
 Species: *Pseudophoxinus zekayi* (Bogustkaya *et al.*, 2000)  
 Genus: *Squalius* (Cuvier, 1817)  
 Species:  
 • *Squalius cephalus* (Linnaeus, 1758)  
 • *Squalius lepidus* (Heckel, 1843)  
 Genus: *Tinca* (Cuvier, 1817)  
 Species: *Tinca tinca* (Linnaeus, 1758)  
 Family: Nemacheilidae  
 Genus: *Oxynemacheilus* (Lacépède, 1790)  
 Species: *Oxynemacheilus samanticus*  
 Genus: *Paracobitis* (Noemacheilus) (Bleeker, 1963)  
 Species:  
 • *Paracobitis seyhanensis*  
 • *Paracobitis (Barbatula) tigris* (Heckel, 1849)  
 Genus: *Schistura* (Clelland, 1838)  
 Species:  
 • *Schistura ceyhanensis* (Erk'akan *et al.*, 2007)  
 • *Schistura evreni* (Erk'akan *et al.*, 2007)  
 • *Schistura seyhanicola* (Erk'akan *et al.*, 2007)

• *Schistura namiri* (Krupp and Schneider, 1991)  
 Family: Cobitidae  
 Genus: *Cobitis* (Linnaeus, 1758)  
 Species:  
 • *Cobitis levantina* (Krupp and Moubayed, 1992)  
 • *Cobitis evreni* (Erk'akan *et al.*, 2008)  
 Class: Siluriformes  
 Family: Clariidae  
 Genus: *Clarias* (Gronovius, 1781)  
 Species: *Clarias gariepinus* (Burchell, 1822)  
 Family: Siluridae  
 Genus: *Silurus* (Linnaeus, 1758)  
 Species: *Silurus glanis* (Linnaeus, 1758)  
 Superclass: Protacanthopterygii  
 Order: Cyprinodontiformes  
 Sub-order: Cyprinodontoidi  
 Family: Cyprinodontidae  
 Genus: *Aphanius* (Nardo, 1827)  
 Species:  
 • *Aphanius mento* (Heckel, 1843)  
 • *Aphanius danfordii* (Boulenger, 1890)  
 Family: Poecilidae  
 Genus: *Gambusia* (Poey, 1854)  
 Species: *Gambusia holbrooki* (Girard, 1859)  
 Class: Perciformes  
 Sub-class: Blennioidei  
 Family: Blenniidae  
 Genus: *Salaria* (Blennius) (Linnaeus, 1758)  
 Species: *Salaria fluviatilis* (Asso, 1801)  
 Family: Percidae  
 Genus: *Mugil* (Linnaeus, 1758)  
 Species: *Mugil cephalus* (Linnaeus, 1758)  
 Genus: *Sander* (Stizostedion) (Linnaeus, 1758)  
 Species: *Sander lucioperca* (Linnaeus, 1758)

## DISCUSSION

In this study, *A. baliki* is recorded for the 1st time from the Camizagili Creek which belongs to Ceyhan river Basin. This species described by Bogustkaya *et al.* (2000) from the Manavgat Basin, Oymapinar. According to their description, number of scales in lateral line series (unpored and pored) varying considerably, 50-65 due to irregular character of scales. Researchers examined nine specimens and treat morphological characters that distinguished *A. baliki* from other *Alburnus* species are the same with this study. *Cobitis levantina* described by Krupp and Moubayed (1992) caught from the Ceyhan river Basin (Hemite Creek and Davutlar village) for the 1st

time in this study. Total 25 specimens agree with in the description of Krupp and Moubayed (1992). Eleven *Nemacheilus namiri* individuals caught from Eglence stream-Seyhan Basin has the same morphometric characteristics and colour pattern given in Krupp and Schneider (1989).

The lateral line scales ranged from 66-76 in the *Capoeta barroisi* samples we examined in the Seyhan and Ceyhan river Basins. Some of the meristic characteristics of *C. barroisi* demonstrated similarities to findings made by Dagli and Erdemli (2008). Samples determined as *C. barroisi* by Sahan and Cengizler (2002) were described as *C. turani* by Ozulug and Freyhof (2008) in the Seyhan river Basin and were described as *C. erhani* by Turan *et al.* (2008) in the Ceyhan river Basin. The lateral line scales in both species, the density of spots on the body, the ossification and length of the last simple dorsal fin ray and the head length being shorter when compared to each other all indicated a distinction. The number of lateral line scales was given between 64-70 in *C. turani* from the Seyhan river Basin by Ozulug and Freyhof (2008), 69-78 in *C. erhani* from the Ceyhan Basin by Turan *et al.* (2008).

However in the study, 25 individuals collected from the Seyhan river Basin had 66-75 lateral line scales and from the Ceyhan river Basin was found to be 64-79 for 40 individuals. Sungur noted that the number of lateral line scales of this species were between 67-89 for the Asi river Basin (type locality of *C. barroisi*) in a study carried out for freshwater fish fauna in the province of Gaziantep. At the same time in this study conducted for the ihtiyofauna of the Seyhan and Ceyhan Basin, the number of spots on the body was observed to be more extensive or less dense in individuals seen in both the Seyhan and Ceyhan basins. Moreover in young individuals in both basins, the length of the last rigid ray of the dorsal fin and the length of the head were shorter than in adult individuals which were found to be longer. In evaluating these findings in terms of zoogeographics, researchers treat *C. erhani* and *C. turani* as a synonyms of *C. barroisi*.

*Capoeta capoeta angorae* described from the Pozanti-Seyhan river Basin by Hanko (1924) based on a single individual has the same variation divergence of the *C. capoeta damascina*. Problems with the level of species and sub-species obtained in systematic studies collected with few individuals are uncovered when evaluated in later studies encompassing larger numbers of individuals. No matter how controversial further argument is in the validity of similar species there is benefit to be drawn from

re-evaluation in the literature. Watanabe (1998) stated that variations between geography and population in such meristic characteristics in freshwater fish as the number of fin rays, lateral line scales and gill rakers on the 1st arch are usually the result of the isolation of populations. Banarescu (1991) also stated that in his researches on the validity of sub-species of freshwater fish, it is necessary to adopt the notion that the circumstances of divergence or overlapping of some meristic characteristics such as the lateral line scale number are not caused by an individuals' subspecies or species but rather by geographical differences arising from clinal variations. The high ecological tolerance of species in the genus *Capoeta* can cause wide variation divergence and populations can display very close or overlapping variation between the metric and meristic characters of each other.

In approximately 2000 years, the Ceyhan river has incorporated with and separated from the Seyhan river six times. Therefore, a species found in the Seyhan river can also be found in the Ceyhan river with local variations. However depending on the speed of evolution of species, it is possible to see some speciation or sub-speciation in populations of species. For instance, species of *Cobitidae* and *Nemacheilidae* living in Anatolia have shown a rapid local speciation.

Bohlen *et al.* (2006) and Erk'akan *et al.* (2008a, b). Differences seen in the Seyhan and Ceyhan river Basins in species belonging to the the *Cobitidae* and *Nemacheilidae* families may be due to the high rate of evolution. In addition, validity of these species by mt-DNA studies has been approved.

In a study conducted by Kara *et al.* (2010) to establish the ihtiyofauna of the middle and upper Ceyhan Basin, samples of *Oxynoemacheilus* sp. from the *Nemacheilidae* family were determined as *Shistura evreni* by us after upon examination of the morphological and diagnostic features of this species.

In the assessment of fish fauna in the Seyhan Dam lake, Alagoz determined some species such as *Acanthalburnus microlepis* and *Capoeta capoeta capoeta* distributed in the Kura-Aras river Basins, *Capoeta capoeta umbla* and *Barbus esocinus* distributed in the Tigris Basin, *Rutilus rutilus*, *Rutilus rubilio* and *Rutilus frisii* distributed in the Eastern black sea and Marmara Basins, *Aphanius chanteri* distributed in the Kizilirmak river and Central black sea Basins, *Abramis brama* distributed in the Black sea Basin, the Marmara and Thrace regions and locally endemic species *Cobitis bilseli* in the Beysehir lake Basin. However from

the perspective of zoogeographical aspects, these species are not found in the fish fauna of the Seyhan river Basin. According to this and other study, it can be concluded that species which had been previously established wrongly have been the cause of significant scientific errors. Bostanci determined 37 species and one sub-species belonging to fourteen families in the Seyhan, Ceyhan and Asi river Basins. From these species, *Barbus barbulus*, *Alburnoides oblongus* and *Hemigrammocapoeta culiciphaga* were new recordings in the fish fauna of Turkey and *Alburnoides bipunctatus* was a new recording for the Mediterranean Basin. Krupp and Schneider (1989) stated that one of the distinguishing features of *H. culiciphaga* is the barbels around the mouth.

However, Bostanci did not recognize barbels around the mouth of *Hemigrammocapoeta culiciphaga* while Krupp and Schneider (1989) and researchers recorded a double set of very short barbells. In addition, Bostanci reported that the lateral line complete and scales were between 30-33 but in other studies (Krupp and Schneider, 1989), it has been recorded that the lateral line is incomplete.

In this study, the lateral line of *H. culiciphaga* living in the Seyhan and Ceyhan Basins was interrupted and the number of scales was found to vary between 5-20. Moreover, researchers did not catch any *Alburnoides oblongus* and *Barbus barbulus* samples from the Ceyhan river Basin. Researchers doubt that these species occurred in this basin may be determined wrongly.

Living in clean, slow-flowing, deep structured stony, gravelly and sandy areas, *Shistura ceyhanensis* is living in the Ceyhan river Basin (Erk'akan *et al.*, 2007). In this study, it was also found in the Seyhan river Basin. In a study by Bostanci, some Nemacheilid specimens determined as *Barbatula panthera*. Based on the meristic, morphometric characteristics and color pattern of these samples, it was established that they are in fact *S. ceyhanensis*.

The type locality of *B. panthera* is the Barada river, in the Damascus Basin and it is endemic for this basin and does not exist in Turkey (Krupp and Schneider, 1989). *S. evreni*, generally prefers clean, cold, oxygen-enriched mountain water. Bostanci determined some Nemacheilid specimens as *N. argyrogramma* but according to morphometric and meristic properties there was identified this species as *S. evreni*.

The type locality of *Noemacheilus tschайssuensis* described by Banarescu and Nalbant is Caysu-Gaziantep-Ceyhan river Basin. However, the coordinates given in the study are all encompassed within the Euphrates Basin and the Ceyhan river does not have any tributary within the

province of Gaziantep. The description of this new species is a synonym of *N. argyrogramma* distributed in the Euphrates Basin. In an oral interview with Teodor Nalbant, this species was verified to be a synonym of *N. argyrogramma*. The metric and meristic characteristics of the *P. seyhanensis* caught by us in the Seyhan river Basin were similar in general to those given by Banarescu. According to Bostanci from the meristic properties of *P. seyhanensis*, the unbranched dorsal fin rays count was II. In the findings, this count was III. In addition, the ratio of standard length for the caudal peduncle length was 14.1% in the findings, Banarescu gave an average of 14.5% and Bostanci gave an average of 18%, respectively. All the other species identified in the study agree with the literature data.

#### ACKNOWLEDGEMENT

Researchers have the thanks to Hacettepe University Research Funding supported this study.

#### REFERENCES

- Ahnelt, H. and J. Holcik, 1996. Distribution of two species of the genus neogobius (Pisces: Gobiidae) in the catchment area of the Southern Caspian Sea. Acta Univ. Carolinae Biol., 40: 99-114.
- Ahnelt, H., 1995. Two new Species of *Kniptowischia* ILJIN, 1997 (Teleostei: Gobiidae) from Western Anatolia. Mitt. Hamb. Zool. Mus. Inst., 92: 155-168.
- Ahnelt, H., P.G. Bianco and H. Schwammer, 1995. Systematics and zoogeography of *kniptowischia caucasica* (Teleostei: Gobiidae) based on new records from the aegean anatolian area. Ichthyol. Explor. Freshwaters, 6: 49-60.
- Balik, S., 1985. Taxonomical revision and the present situations of the inland water from Turkish Thrace. Doga Bilim Derg. A, 9: 2-2.
- Banarescu, P.M., 1991. The subspecies in freshwater ichthyology. Evol. Daptation, 4: 183-189.
- Bogustkaya, N.G., 1992. A revision of species of the genus *pseudophoxinus* (Leuciscinae, Cyprinidae) from asya minor. M. Hamburg Zool. Mus. Inst., 89: 261-290.
- Bogustkaya, N.G., 1997. Contribution to the knowledge of leuciscinae fishes of the Asia Minor. Part 2., An annotated check-list of leuciscinae fishes (Leuciscinae, Cyprinidae) of Turkey with description of a new species and two new subspecies. Mitt. Hamb. Zool. Mus. Inst., 94: 161-186.
- Bogustkaya, N.G., F. Kucuk and E. Unlu, 2000. *Alburnus baliki*, a new species of cyprinid fish from the manavgat river system Turkey. Ichthyol. Explor. Freshwaters, 11: 55-64.

- Bogustkaya, N.G., F. Kucuk and M.A. Atalay, 2006. A description of three new species of the genus *Pseudophoxinus* from Turkey (Teleostei: Cyprinidae: Leuciscinae). *Zoosyst. Rossica*, 15: 335-341.
- Bohlen, J., A. Percides, I. Doadrio and P.S. Economidis, 2006. Vicariance, colonisation and fast local speciation in Asia Minor and the Balkans as revealed from the phylogeny of spined loaches (Osteichthyes: Cobitidae). *Mol. Phylogenet. Evol.*, 39: 552-561.
- Coad, B. and M. Sarieyyupoglu, 1988. *Cobitis elazigensis*, A new species of cobitid fish from anatolia, Turkey. *Jap. J. Ichthyol.*, 34: 426-430.
- Dagli, M. and A.U. Erdemli, 2008. A taxonomical study on the fish of sabun suyu and delicay stream (Kilis, Turkey). *Int. J. Sci. Technol.*, 3: 19-25.
- Elvira, B., 1997. Taxonomy of the genus *Chondrostoma* (Osteichthyes, Cyprinidae) an updated review. *Folia Zool.*, 46: 1-14.
- Erk'akan, F. and M. Kuru, 1986a. A new noemacheilinae loach subspecies from Turkey osteichthyes-Cobitidae). *Turk. J. Biol.*, 10: 106-109.
- Erk'akan F. and M. Kuru, 1986b. A new noemacheilinae loach subspecies from the Lake Van Basin, Turkey (Osteichthyes- Cobitidae). *Turk. J. Biol.*, 10: 160-162.
- Erk'akan, F. and M. Kuru, 1983. Re-discussion of systematical status of *Varicorhinus antalyensis* Battalgil, 1944. *Hacettepe Bull. Nat. Sci. Eng.*, 12: 49-65.
- Erk'akan, F., 1983a. The fish species of the Sakarya basin and their abundance. *Hacettepe Bull. Nat. Sci. Eng.*, 12: 21-38.
- Erk'akan, F., 1983b. Studies on the systematics and bio-ecology of the fishes in the Sakarya Basin, (in Turkish). *Doga Bilim Dergisi Vet. ve Hayvancilik*, 7: 141-153.
- Erk'akan, F., 1984. A new cyprinid fish record for Turkey, *Pseudorasbora parva* from the thrace region, (in Turkish). *Doga Bilim Dergisi*, 3: 350-351.
- Erk'akan, F., F.G. Atalay-Ekmekci and T.T. Nalbant, 1998. Four new species and one new subspecies of the genus *cobitis* (Pisces: Ostariophysi: Cobitidae) from Turkey. *Turk. J. Zool.*, 22: 9-15.
- Erk'akan, F., S.C. Ozeren and T.T. Nalbant, 2008a. *Cobitis evreni* sp. nova-a new spined loach species (Cobitidae) from the southern Turkey. *J. Fish. Int.*, 3: 112-114.
- Erk'akan, F., S.C. Ozeren and T.T. Nalbant, 2008b. Two new species of stone loaches from Turkey (Teleostei: Nemacheilidae). *J. Fish. Int.*, 3: 115-119.
- Erk'akan, F., T.T. Nalbant and S.C. Ozeren, 2007. Seven new species of Barbatula, three new species of Schistura and a new species of Seminemacheilus (Ostariophysi: Balitoridae: Nemacheilinae) from Turkey. *J. Fish. Int.*, 2: 69-85.
- Freyhof, J. and M. Ozulug, 2006. *Pseudophoxinus ninae*, a new species from Central Anatolia, Turkey (Teleostei: Cyprinidae). *Ichthyol. Explor. Freshwaters*, 17: 255-259.
- Hanko, B., 1924. Fische aus klein-Asien. *Ann. Hist. Nat. Mus. Nation Hung.*, 21: 137-158.
- Kara, C., A. Alp and M. Simsekli, 2010. Distribution of fish fauna on the upper and middle basin of ceyhan river, Turkey. *Turk. J. Fish. Aquat. Sci.*, 10: 111-122.
- Krupp, F. and J. Moubayed, 1992. A new species of *cobitis linnaeus*, 1758 from the orontes and litani drainage basins of Syria and Lebanon (Pisces: Osteichthyes: Cobitidae). *Senckenbergiana Biol.*, 72: 13-18.
- Krupp, F. and W. Schneider, 1989. The fishes of the Jordan River drainage basin and Azraq oasis. *Fauna Saudi Arabia*, 10: 347-416.
- Krupp, F. and W. Schneider, 1991. Two new species of *Nemacheilus* Bleeker 1863 from the Orontes River drainage basin of Lebanon, Syria and Turkey. *Senckenbergiana Biol.*, 71: 23-34.
- Krupp, F., 1985. A new species of *Chondrostoma* from the Orontes River drainage basin of Turkey and Syria. *Senckenbergiana Biol.*, 66: 27-33.
- Kucuk, F., I.I. Turna and O. Demir, 2007. Taxonomic characteristics and distribution of *capoeta pestai* (Pietschmann, 1933) (Pisces: Cyprinidae) (In Turkish). *SDU Fen Bilimleri Enst. Dergisi*, 11: 18-25.
- Ozulug, M. and J. Freyhof, 2007. *Alburnus demiri*, A new species of bleak from Western Anatolia, Turkey (Teleostei: Cyprinidae). *Ichthyol. Explor. Freshwaters*, 18: 307-312.
- Ozulug, M. and J. Freyhof, 2008. *Capoeta turani* a new species of barbel from river Seyhan, Turkey (Teleostei: Cyprinidae). *Ichthyol. Explor. Freshwaters*, 19: 289-296.
- Sahan, A. and I. Cengizler, 2002. Determination of some hematological parameters in spotted barb (*Capoeta barroisi*, 1894) and roach (*Rutilus rutilus*, 1758) Living in Seyhan River (Adana City Region). *Turk. J. Vet. Anim. Sci.*, 26: 849-858.
- Turan, D., B.T. Yilmaz and C. Kaya, 2009. *Squalius kottelati*, a new cyprinid species (Teleostei: Cyprinidae) from Orontes River, Turkey. *Zootaxa*, 2270: 53-62.
- Turan, D., M. Kottelat and F.G. Ekmekci, 2008. *Capoeta erhani*, A new species of cyprinid from ceyhan River, Turkey (Teleostei: Cyprinidae). *Ichthyol. Explor. Freshwaters*, 19: 263-270.

- Turan, D., M. Kottelat, S.G. Kirankaya and S. Engin, 2006. *Capoeta ekmekciae*, a new species of cyprinid fish from northeastern Anatolia (Teleostei: Cyprinidae). *Ichthyol. Explor. Freshwaters*, 17: 147-156.
- Watanabe, K., 1998. Meristic variation in the endangered bagrid catfish, *Pseudobagrus ichikawai*. *Ichthyol. Res.*, 45: 99-104.
- Wildekamp, R.H., W.V. Neer, F. Kucuk and M. Unlusayin, 1997. First record of the eastern Asiatic gobionid fish *Pseudorasbora parva* from the asiatic part of Turkey. *J. Fish Biol.*, 51: 858-868.
- Wildekamp, R.H., F. Kucuk, M. Unlusayin and W.V. Neer, 1999. Species and subspecies of the genus *aphanius nardo 1827* (Pisces: Cyprinodontidae) in Turkey. *Turk. J. Zool.*, 23: 23-44.