



Diversity of Endophytic Fungi from Leaves of *Cenetlla asiatica* L. and *Melia azedarach* L.

A.S. Shijila Rani, V. Ambikapathy and A. Panneerselvam

PG and Research Department of Botany and Microbiology, AVVM Sri Pushpam College (Autonomous), 613503 Poondi, Thanjavur, India

Key words: *Cenetlla asiatica*, *Melia azedarach*, leaves, Czapek dextrose agar, endophytic fungi, diversity

Corresponding Author:

A.S. Shijila Rani

PG and Research Department of Botany and Microbiology, AVVM Sri Pushpam College (Autonomous), 613503 Poondi, Thanjavur, India

Page No: 22-25

Volume: 15, Issue 2, 2020

ISSN: 1815-8846

Research Journal of Biological Sciences

Copy Right: Medwell Publications

Abstract: Endophytic fungi are symbiotic association and the harbor of plants. It is living in inner tissues and without causing any apparent symptoms. In the present investigation, the endophytic fungi were isolated from leaves of *Cenetlla asiatica* and *Melia azedarach*. These 2 plants are highly medicinal properties. Totally eight different genera were isolated from both the leaves. Generally, *Absidia repens*, *Alternaria* sp., *Aspergillus flavus*, *A. terreus*, *Chaetomium globosum*, *Curvularia* sp., *Fusarium solani* and *Humicola* sp. The maximum percentage of endophytic fungi and diversity indices were recorded in *Melia azedarach* leaves fragment when compared to *Cenetlla asiatica* leaves.

INTRODUCTION

Endophytes are ubiquitous in nature and are found in every plant. Endophytic fungi are microorganisms found inside plant species^[1] and may play a key role in plant development by controlling endophytic microbes and herbivorous insects or producing growth promoting substances^[2]. Endophytic fungi also have functions related to saprophytic microorganisms and many species are involved in the process of maturation and leaf decomposition^[3, 4]. Endophytic fungi infect and inhabit primarily the aerial tissue of the host plant without causing detectable symptoms. The relationship between endophytes are thought to be symbiotic such as those endophytes to get nutrients and protection from the host but contribute to effective host defense mechanism against pathogens herbivores or abiotic stress^[5, 6].

One of the important role of endophytic fungi is to initiate the biological degradation of dead or dying host plant which is necessary for nutrient recycling^[7]. Endophytic fungi are reported from plants grow in various environmental including tropic^[8] temperate^[9] xerophytic^[10] and aquatic environment^[11]. Medicinal

plants are reported to harbour endophytes^[12] which in turn provide protection to their host from infection agents and also, provide adaptability to survive in adverse environmental condition. It is therefore, important to determine the endophytic fungi diversity of medicinal plants. Recently, the knowledge about endophyte biodiversity is becoming more apparent.

Cenetlla Asiatica (CA) is a very important medicinal herb used in the orient^[13] which is also, becoming popular in the West^[14]. It is a tasteless, odourless plant that thrives in and around water. It has small fan-shaped green leaves with white or light purple-to-pink or white flowers and it bears small oval fruit. The whole plant is used for medicinal purposes^[15]. It is widely used as a blood purifier as well as for treating high blood pressure for memory enhancement and promoting longevity. In Ayurveda, *Cenetlla asiatica* is one of the main herbs for revitalizing the nerves and brain cells.

Melia azedarach is a species of deciduous tree in the Mahogany family, Meliaceae that is an evergreen tree, cultivated in various parts of the Indian subcontinents. Neem has a long history of use in the traditional medical systems of India (Ayurvedic, Unani-Tibb). Leaves have

been used as a natural insecticide to keep with stored food but must not be eaten as they are highly poisonous. A diluted infusion of leaves and trees has been used in the past to induce uterus relaxation. Extracts from neem leaf, seed and bark possess a wide spectrum of antibacterial action^[16,17]. Although, the fruits are poisonous part of the tree, they have been used traditionally for the treatment of a variety of diseases, especially, dermatitis and rubella^[18].

MATERIALS AND METHODS

Sample collection: Collection, identification and authentication of leaf materials of *Melia azedarach* L. and *Centella asiatica* L. were collected from Thanjavur District (the latitude in such as 10.6590°N and longitude in 79.2451°E) Tamil Nadu, India. The plants were identified by taxonomist for St. Joesph's College, Triuchirappalli (Fig. 1).

Isolation of endophytes: The endophytic fungi were isolated from leaves of *Melia azedarach* L. and *Centella asiatica* L. as described by Strobel and Daisy (2003). In first instance, the leaves were cleaned by thorough washing in running tap water, followed by Deionized (DI) water. Clean leaves were cut into small pieces were sterilized in series of solution; 70% ethanol, 1% sodium hypochlorite (v/v) again 70% ethanol for 1 min each. Finally, they were again rinsed with sterile distilled water two times. After surface sterilization, the tissues were dried on blotting sheets and cut into 1 cm square pieces. These sterile small pieces were placed on Czapek Dextrose Agar (CDA) plates containing streptomycin (250 µg mL⁻¹) to inhibit the bacterial growth. The plates were incubated at 25±2°C after wrapping with parafilm and observed daily. The fungal mycelia which started growing from the tissues were sub-cultured on new CDA plates. These endophytes were stored in paraffin at 4°C.

Colony frequency: Colonization Frequency (CF %) was calculated as described by Suryanarayanan *et al.*^[19] and Photita *et al.*^[20]. The colonies appearing on petriplates were sub-cultured into the tube containing Czapek dextrose agar medium for identification. Morphological identification was done according to the standard taxonomic key included colony diameter, texture, color and the dimensions and morphology of hyphae and conidia:

$$\text{Percentage of colony} = \frac{\text{No. of colonies in individual segment}}{\text{Total number of colonies in the sample}} \times 100$$



Fig. 1: Map showing the study site

$$\text{Percentage of CF} = \frac{\text{No. of segment colonized by fungi}}{\text{Total number of segment observed}} \times 100$$

Diversity index: The species diversity of the endophytic fungi was measured using the diversity indices with species richness and relative species abundance. The species evenness which assesses the contribution of the individuals to the community also, calculated.

The diversity indices used for the isolates were the Shannon-Wiener index (H) (formula: [(ni/n) ln(ni/n)]) Simpson's diversity index (1-D) (formula: 1-[D = (ni/n) 2]) and Simpson's dominance index (D) where ni is the number of distinct species (i) and (n) is the abundance of each species in the community. The equitability was calculated by the following formula; E = H/H_{max} where H is the Shannon-Wiener index and H_{max}, is the ln of ni. The diversity analyses were performed the MS Excel program^[21].

RESULTS AND DISCUSSION

In the present study, 11 isolates were obtained from 7 tissue fragments from 2 medicinal plant species. The 10 different types of genera were isolated, the maximum number of colonies of *Absidia repens* and *Aspergillus terreus* were observed in *M. azedarach* and *A. terreus* in *C. asiatica* leaves fragment. The number of colony, percentage of colony and percentage of colony frequency were recorded Table 1.

Endophytic fungal diversity is measured in terms of Shannon, Margalef's index, Menhinick's species richness,

Table 1: Isolation of endophytic fungi from the leaf samples

Name of the fungi	<i>M. azedarach</i>			<i>C. asiatica</i>		
	No. of colony	CP (%)	Colony frequency	No. of colony	CP (%)	Colony frequency
<i>Absidia repens</i>	5	20	71.42	1	6.25	20
<i>Alternaria</i> sp.	1	4	14.28	1	6.25	20
<i>Aspergillus flavus</i>	4	16	57.14	2	12.5	40
<i>A. terreus</i>	5	20	71.42	4	25	80
<i>Chaetomium globosum</i>	1	4	14.28	1	6.25	20
<i>Curvularia geniculata</i>	1	4	14.28	-	0	0
<i>C. lunata</i>	-	0	0	1	6.25	20
<i>Curvularia</i> sp.	2	8	28.57	2	12.5	40
<i>Fusarium solani</i>	3	12	42.85	3	18.75	60
<i>Humicola</i> sp.	2	8	28.57	1	6.25	20
<i>Melanospora</i> sp.	1	4	14.28	-	0	0
Total	25	100	357.14	16	100	320

Table 2: Diversity indices of isolated endophytic fungi

Diversity indices	<i>M. azedarach</i>	<i>C. asiatica</i>
No. of species	25	16
No. of individuals	10	9
Genera	10	8
Margalef's index (d)	6.463	4.039
Mehinick's Species Richness (SR)	5.735	3.671
Pielou's Evenness (J)	0.086	0.081
McIntosh's Evenness (McE)	0.141	0.188
Shannon's Diversity (H)	0.277	0.224
Margalef's Diversity (Dmg)	12.427	14.427
Fisher's alpha	6.17	8.5

Margalef's Diversity, McIntosh's Evenness, Simpson and Pielou's evenness and Fisher's alpha indices. The diversity indices were calculated in the endophytic fungi of both the plants. The highest species richness present in *M. azedarach* when compared to *C. asiatica*. The Margalef's Diversity is 14.427 in *C. asiatica* leaves and 12.427 in *M. azedarach* (Table 2).

Endophytic microorganism, especially, fungi established a symbiotic relationship with plants while living inside the plant and synthesize metabolites that often helps their host to survive and adapt in adverse environment^[22]. In addition, they are the group of organism with very good potential for application in plant improvement and disease control^[23]. A study of endophytic biodiversity of the temporal variation in *Plumeria rubra* leaves was conducted by the Suryanarayanan and Thennarasan^[24].

In the similar study 2 different genera of *Aspergillus* (6.6) and *Nigrospora* (3.3%) the colonization frequency were isolate from *Melia azedarach*. This is slightly less than the above cited report by Suryanarayanan and Thennarasan^[24]. During the present study, *Absidia* sp., *Alternaria* sp., *Aspergillus* sp., *Chaetomium* sp., *Curvularia* sp., *Fusarium* sp., *Humicola* sp. and *Melanospora* sp. were isolated from *Melia azedarach* and *Centella asiatica*. All the species belong to Hyphomycetes class. These class fungi were largely prevalent and showed the resistance too many pathogens. The genus *Aspergillus* found most common in plant.

Similar to the findings reported previously in *Centella asiatica* leaves, the higher colonization of endophytic fungi in mature leaf as compared to young leaf over all the 3 seasons suggests that mature leaf provides more favourable environment for the establishment of endophytic fungi^[25]. Nascimento *et al.*^[26] reported *Phaeoramularia calotropidis* (63.5%) as dominant endophyte in *Calotropis procera* leaves.

The diversity of the endophytic fungal populations isolated from leaves of *C. asiatica* and *Melia azedarach* such as was evaluated by various indices such as Fisher alpha diversity index (α) Shannon index H, Simpson's Diversity index (1-D) and Margalef richness index (Dmg) (Table 2). Similarly, Mehinick's index is similar, conceptually, to Margalef's richness index for analyzing species richness.

CONCLUSION

The assume diversity values from each of the 2 indices showed similar patterns in these results because of the endophytes, Shannon's diversity index (H')^[27] and Simpson's Diversity index (D')^[28] were not used. The species richness (Dmn) was highest in the leaves (4.24) followed by the flowers (2.82) and roots (1.41)^[29].

ACKNOWLEDGEMENT

The researchers thank the principal of A.V.V.M Sri Pushpam College (Autonomous), Poondi, Thanjavur and the Director, Indian Biotrack Research Institute, Thanjavur for extending the lab facilities to carry out the research.

REFERENCES

- Stone, J.K., J.D. Polishook and J.F. White, 2004. Endophytic Fungi. In: Biodiversity of Fungi: Inventory and Monitoring Methods, Mueller, G.M., G.F. Bills and MS. Foster (Eds.). Elsevier Academic Press, Cambridge, Massachusetts, USA., pp: 241-270.

02. Peixoto, N.P., J.L. Azevedo and W.L. Araujo, 2002. Microrganismos endofíticos: Interação com as plantas e potencial biotecnológico. *Biociencia*, 29: 62-76.
03. Promputtha, I., S. Lumyong, V. Dhanasekaran, E.H.C. McKenzie and K.D. Hyde *et al.*, 2007. A phylogenetic evaluation of whether endophytes become saprotrophs at host senescence. *Microb. Ecol.*, 53: 579-590.
04. Sieber, T.N., 2007. Endophytic fungi in forest trees: Are they mutualists?. *Fungal Biol. Rev.*, 21: 75-89.
05. Redman, R.S., K.B. Sheehan, R.G. Stout, R.J. Rodriguez and J.M. Henson, 2002. Thermotolerance generated by plant/fungal symbiosis. *Science*, 298: 1581-1581.
06. Arnold, A.E. and E.A. Herre, 2003. Canopy cover and leaf age affect colonization by tropical fungal endophytes: Ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia*, 95: 388-398.
07. Strobel, G.A., 2002. Microbial gifts from rain forests. *Can. J. Plant Pathol.*, 24: 14-20.
08. Mohali, S., T.I. Burgess and M.J. Wingfield, 2005. Diversity and host association of the tropical endophyte *Lasiodiplodia* revealed using simple sequence repeat markers. *For. Path.*, 35: 385-396.
09. Ganley, R.J., S.J. Brunfeld and G.A. Newcombe, 2004. A community of unknown, endophytic fungi in western white pine. *Nat. Acad. Sci. USA.*, 101: 10107-10112.
10. Suryanarayanan, T.S., S.K. Wittlinger and S.H. Faeth, 2005. Endophytic fungi associated with cacti in Arizona. *Mycol. Res.*, 109: 635-639.
11. Sraj-Krzic, N., P. Pongrac, M. Klemenc, A. Kladnik and M. Regvar *et al.*, 2006. Mycorrhizal colonisation in plants from intermittent aquatic habitats. *Aquat. Bot.*, 85: 331-336.
12. Strobel, G. and B. Daisy, 2003. Bioprospecting for microbial endophytes and their natural products. *Microbiol. Mol. Biol. Rev.*, 67: 491-502.
13. Bown, D., 1995. *Encyclopaedia of Herbs and their Uses*. Dorling Kindersley, London, UK., Pages: 424.
14. Chevallier, A., 1996. *The Encyclopedia of Medicinal Plants*. 1st Edn., DK Publishing Inc., New York, USA., pp: 336.
15. Singh, P. and J.S. Singh, 2002. Recruitment and competitive interaction between ramets and seedlings in a perennial medicinal herb *Centella asiatica*. *Basic Appl. Ecol.*, 3: 65-76.
16. Biswas, K., I. Chattopadhyay, R.K. Banerjee and U. Bandyopadhyay, 2002. Biological activities and medicinal properties of neem (*Azadirachta indica*). *Curr. Sci.*, 82: 1336-1345.
17. Mahfuzul Hoque, M.D., M.L. Bari, Y. Inatsu, V.K. Juneja and S. Kawamoto, 2007. Antibacterial activity of guava (*Psidium guajava* L.) and Neem (*Azadirachta indica* A. Juss.) extracts against foodborne pathogens and spoilage bacteria. *Foodborne Pathog Dis.*, 4: 481-488.
18. Kim, M., S.K. Kim, B.N. Park, K.H. Lee and G.H. Min *et al.*, 1999. Antiviral effects of 28-Deacetylshikimic acid on herpes simplex virus-1 replication. *Antiviral Res.*, 43: 103-112.
19. Suryanarayanan, T.S., G. Venkatesan and T.S. Murali, 2003. Endophytic fungal communities in leaves of tropical forest trees: Diversity and distribution patterns. *Curr. Sci.*, 85: 489-492.
20. Photita, W., S. Lumyong, P. Lumyong and K.D. Hyde, 2001. Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep Pui national Park. *Mycol. Res.*, 105: 1508-1513.
21. Hammer, O., D.A.T. Harper and P.D. Ryan, 2001. PAST: Paleontological statistics software package for education and data analysis. *Paleontol. Electron.*, 4: 1-9.
22. Clay, K. and C. Schardl, 2002. Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *Am. Nat.*, 160: 99-127.
23. Sandhu, S.S., S. Kumar and R.P. Aharwal, 2014. Isolation and identification of endophytic fungi from *Ricinus communis* linn and their antibacterial activity. *Intl. J. Res. Pharm. Chem.*, 4: 611-618.
24. Suryanarayanan, T.S. and S. Thennarasan, 2004. Temporal variation in endophyte assemblages of *Plumeria rubra* leaves. *Fungal Diversity*, 15: 197-204.
25. Gupta, S. and P. Chaturvedi, 2017. Foliar endophytic diversity of *Centella asiatica* (L.) urban in relation to different seasons and leaf age. *Intl. J. Curr. Microbiol. Appl. Sci.*, 6: 468-477.
26. Nascimento, T.L., Y. Oki, D.M.M. Lima, J.S. Almeida-Cortez and G.W. Fernandes *et al.*, 2015. Biodiversity of endophytic fungi in different leaf ages of *Calotropis procera* and their antimicrobial activity. *Fungal Ecol.*, 14: 79-86.
27. Pielou, E.C., 1975. *Ecological Diversity*. John Wiley and Sons Inc., New York, Pages: 165.
28. Simpson, E.H., 1949. Measurement of diversity. *Nature*, 163: 688-688.
29. Katoch, M., S. Phull, S. Vaid and S. Singh, 2017. Diversity, phylogeny, anticancer and antimicrobial potential of fungal endophytes associated with *Monarda citriodora* L. *BMC. Microbiol.*, 17: 1-13.