Kinetic of Carbon Dioxide Production by *Leuconostoc mesenteroides* Grown in Single and Mixed Culture with *Lactococcus lactis* in Skimmed Milk

¹M. Kihal, ²H. Prevost, ²D.E. Henni, ²Z. Benmechernene and ²C. Diviès ¹Laboratoire de Microbiologie Appliquée, Département de Biologie, Faculté des Sciences, Université Oran, Es-senia, Bp 16, 31000, Oran, Algérie ²Laboratoire de Microbiologie-Biotechnologie, ENSBANA, 1 Esplanade Erasme Université de Bourgogne 21000, Dijon, France

Abstract: The effect of mixed culture of *Leuconostoc mesenteroides* subsp. dextranicum and *Lactococcus lactis* subsp. lactis (prot⁺) was investigated to achieve an optimal production of carbone dioxide. Only the strain of *Leuconostoc mesentroides* can produce carbon dioxide from lactose and citrate in milk. The influence of the initial concentration ration between the 2 strains on growth, carbon dioxide, L-lactate, acetic acid production and citrate used was displayed. When the initial inoculum of *Lactococcus lactis* was 2.5 10⁵ cfu mL⁻¹, the growth and evolved CO₂ by *Leuconostoc mesenteroides* (3 10⁷ cfu mL⁻¹) increased. Whereas, high inoculum of *Lactococcus lactis* induce a decrease of growth and CO₂ production by *Leuconostoc mesenteroides*. In mixed culture CO₂ production continued after growth stopped, a partial uncoupling can be observed between growth and CO₂ production. A shift of acetate production was observed in mixed culture and 25.6 mM was obtained, whereas 30.18 mM was obtained at the same time in pure culture of *Leuconostoc mesenteroides*.

Key words: Leuconostoc mesenteroides, Lactococcus lactis, milk, lactic acid, acetic acid, citric acid, growth kinetics, fermented milk

INTRODUCTION

Lactic acid bacteria are now used extensively as starter cultures in the dairy industry and therefore, the optimization of growth conditions appears to be essential for successful industrial applications. Furthermore, studying the effects of some environmental parameters on growth kinetics should provide useful information concerning the physiology of the microorganisms (Carr et al., 2002; Ayad et al., 2004). Strains belonging to the species Lactococcus lactis and Leuconostoc sp. are the most important organism in the manufacture of these products at a moderate temperature (Drosinos et al., 2005). Mixed cultures of these bacteria are commonly used as a starter in manufacture of cheese and fermented milk (Bellangier et al., 1999). Large-scale industrial processes rely on the use of starter cultures that have been selected for their performance during milk fermentation and product formation.

Leuconostoc strains grow associatively with acid producing Lactococcus strains and are employed for their technological properties (mainly aroma and texture). The associative growth between these 2 groups of bacteria

has been studied with respect to citrate metabolism and aroma formation and has been described as a synergistic functional relationship. Thus attention has been devoted to the growth rate, acid production and final population of these bacteria, which also reflect the interactions occurring in mixed strain cultures of *Leuconostoc* and *Lactococcus* (Bellangier *et al.*, 1999; Padmanabhan and Kim, 2002).

The importance of *Leuconostoc* sp. in dairy fermentation is related to their ability to produce carbon dioxide from lactose and citrate. Various methods have been used to measure CO₂ production by microorganisms (Gibson and Abdelmalek, 1945; Kneifel and Gretner, 1992; Mohr *et al.*, 1993; Girard and Boyaval, 1994; Kihal *et al.*, 2006). Holmes *et al.* (1968) demonstrated that cell extract of *Lactococcus lactis* and *Lactobacillus* enhanced gas production of *Leuconostoc mesenteroides* subsp. *cremoris*. During the fermentation of milk by *Leuconostoc* the ratio of CO₂ is critical as it affects not only the flavor and aroma of the fermented product but also on the texture of blue cheese.

The shift in metabolic pathways in response to environmental conditions is well documented in the literature in the case of *Homofermentative* sp. (Bobillo and Marshall, 1992; Kandler, 1983), but little information is available about the behavior of *Heterofermentative* sp. such as *Leuconostoc mesenteroides*. The quantitative aspects and kinetics of evolved CO₂ in mixed culture of *Leuconostoc-Lactococcus* under different culturing conditions have been fairly studied (Beal and Corrieu, 1991; Kihal *et al.*, 1996). In point of view of the recent report demonstrating the importance of evolved CO₂ to the texture of dairy products, a study of this matter seemed appropriate.

The present study was undertaken to determine whether *Leuconostoc mesenteroides* plays a role in carbon dioxide production in pure and mixed culture with *Lactococcus lactis* in milk.

MATERIALS AND METHODS

Bacterial strains: Leuconostoc mesenteroides subsp. dextranicum (L4), Leuconostoc mesenteroides subsp. mesenteroides (19D) and Lactococcus lactis subsp. lactis (SLO₃), which is known for its proteolytic activity, were obtained from Laboratoire de Microbilogie-biotechnologie, ENSBANA, Dijon, France. The identity of species was confirmed by the use of physiological, biochemical and sugar test by API 50 CHL system according to the manufacturer's instructions (API System, Bio-Merieux France) (Huang et al., 1994; Thapa et al., 2006; Perea et al., 2007).

The strains were stored as frozen stock at 20°C in fortified skimmed milk (10% skim milk, 0.25% yeast extract, 0.5% glucose) containing 30% glycerol as appropriate. Working cultures were prepared from stock cultures by 2 consecutive transfers in fresh MRS and M17 broth for *Leuconostoc* and *Lactococcus*, respectively (Kihal, 1996).

Media and conditions cultures: Leuconostoc strains were cultured in MRS broth (De Man et al., 1960) and Lactococcus lactis in M17 broth (Terzaghi and Sandine, 1975). Milk medium (reconstituted skim milk 10%) was also used for the preparation of single and mixed cultures. Skimmed-milk medium was prepared from reconstituted skim milk powder 10% (w v⁻¹) and sterilized by autoclaving at 110°C for 10 min. All cultures were incubated at 30°C for 48 h.

Kinetic growth on milk: The growth rate, lactic acid production and evolved CO₂ were studied on reconstituted skim milk. The reconstituted milk was sterilized at 110°C for 10 min, cooled at room temperature (20°C) and inoculated with the strains at 30°C. All the mother culture of *Leuconostoc* and *Lactococcus lactis* were grown separately in sterile skim milk at 30°C.

Freshly prepared starter culture of *Leuconostoc* was inoculated at 5% initially yielded about 5×10^7 cfu mL⁻¹, however, in the case of *Lactococcus lactis*, the yield was about 2.5×10^5 cfu mL⁻¹ at 0.1% of inoculums. Cultures were incubated at 30°C for 24 h. Every hour, samples were aseptically withdrawn from tubes to determine pH, titratable acidity and evolved CO_2 , the experiments were repeated 3 times.

Determination of pH and total acidity: Measurement of pH was carried out by pH-meter. The total acidity was determined by titrating 10 mL of culture with 0.1 N NaOH, phenolphthalein was used as indicator and total acidity was expressed as mM of lactic acid (Huang *et al.*, 1994; Badis *et al.*, 2004).

Growth kinetic: Cultures samples were collected aseptically at 0 and every 2 h post inoculation until 24 h. Culture sample of 1 mL were submitted to decimal dilutions in sterile tryptone salt solution and agar plate were performed to assess cell count. *Leuconostoc mesenteroides* and *Lactococcus lactis* were enumerated in MRSv and M17, respectively according to the method of Mathot *et al.* (1994). Plates were incubated at 30°C for 48 h. Generation times were calculated in the logarithmic phase of growth.

 ${\bf CO_2}$ production: The evolved ${\bf CO_2}$ was measured by a technique based on the pressure which is created by ${\bf CO_2}$ production by culture in tubes. Evolved ${\bf CO_2}$ was trapped in burette in which it was measured (Kihal *et al.*, 2006). The total amount of ${\bf CO_2}$ produced was released by acidifying with 0.5 mL of HCl 2N (Mohr *et al.*, 1993).

In order to evaluate the linearity of the method, solution of sodium carbonate 50 mM were used to liberate CO_2 in the tube by addition of sulfuric acid (2N). The blank contains 10 ml of sterile milk.

Analytical methods: The ability to produce different lactic acid isomers (L-lactate and D-lactate) was tested by an enzymatic method utilizing Boehringer Mannheim GmbH (Mannheim, Germany), Also, citric acid and acetic acid were carried out by enzymatic methods (Boehringer) (Bjokroth *et al.*, 2000).

RESULTS AND DISCUSSION

Strains characteristics: All the used strains were cocci associated in diplococci and chain, Gram+, catalase-and were can grow under both aerobic and anaerobic conditions. All the strains form on solid medium identical lentil colonies.

The $2\,Leuconostoc$ strains were Heterofermentatives and produce CO2 and D-lactic acid from glucose. The

production of dextrane from saccharose was observed. However, *Leuconostoc mersenteroides* subsp. mesenteroides (19D) was arabinose+ and *Leuconostoc mersenteroides* subsp. *dextranicum* (L4) was arabinose-.

In the case of *Lactococcus lactis* (SLO₃) was homofermentative and produce the isomer L-lactate from glucose. This strain can not use citrate and hydrolyze arginine (Perea *et al.*, 2007).

All this characteristics are in accordance with Carr *et al.* (2002), Stiles and Holzapfel (1997) and Klein *et al.* (1998).

Limite of detection: A high correlation was observed when the volume of CO₂ measured is lower than 80 mM. The method has a good linearity between 0 to 80 mM of CO₂, but the application of the method to the determination of CO₂ content in the sample needs a blank which must be prepared and measured before each culture sample. The lower value of coefficient of variation observed was caused by the preparation of sample than by the measurement itself (Table 1).

By definition, Heterofermentative lactic acid bacteria, ferment glucose to produce equimolecular amounts of lactate, carbon dioxide and acetate or ethanol (Kandler, 1983; Kihal et al., 1996). However, certain modifications in the conditions of culture may result in the prevalence of one of these products. The kinetics of both evolved CO₂ and pH evolution are shown in Fig. 1 and 2. When Leuconostoc mesenteroides was cultured with Lactococcus lactis subsp. Lactis prot (+) in milk, several differences in the fermentation products were observed. No evolved CO₂ was observed in pure culture of Lactococcus lactis.

A proportion of 2.5×10^6 cfu mL⁻¹ of *Lactococcus lactis* decreases the CO₂ production and gives only 12.5 mM of CO₂ in mixed culture after 24 h of incubation, while 2.5×10^5 cfu mL⁻¹ of *Lactococcus lactis* enhanced CO₂ production in mixed culture (Fig. 1). Good growth of *Leuconostoc mesenteroides* was observed by Todorov and Dicks (2005) in the presence of 10% soy milk or molasses. The final volume of CO₂ was 17.4 and 1.5 mM in p ure culture of *Leuconostoc mesenteroides* subsp *dextranicum* and *Lactococcus lactis*, respectively. A volume of 72.86 mM of CO₂ was observed when

Table 1: Illustrate the response of the complete device to the liberation of CO₂ from a solution of sodium carbonate; Mean and standard deviation were obtained from 10 measurements

mL of Na ₂ CO ₃ 50 mM	Mean (mL)	Std. Deviation	Coefficient of variance (%)
8	9.70	0.38	3.2
10	10.91	0.099	0.9
12	14.5	0.51	3.5

Lactococcus lactis was used (2.5×10⁵ cfu mL⁻¹) with Leuconostoc (3×10⁷ cfu mL⁻¹) (Fig. 3). A greater inhibition of evolved CO2 coming from the contribution of total lactic acid in mixed culture was caused by high inoculums of Lactococcus lactis. Bellangier et al. (1999) demonstrated that the proteolytic Lactococcus inhibit the gas production of Leuconostoc sp. In Fig. estimation of results coming from the pure culture relationships applied to a mixed culture experiment are presented. A shift between CO2 production rate in pure and mixed culture could be appreciated. In pure culture of Leuconostoc mesenteroides the production of lactic acid was correlated with evolved CO2 and the coefficient of correlation was high (r = 0.996). The evolution of the curves of CO2 and lactic acid production were almost identical in pure culture of Leuconostoc. Kinetics of pH evolution by culture was presented in Fig. 2.

The lowest titrable acidity and pH evolution were given by the pure culture of *Leuconostoc*. As starter cultures *Leuconostoc mesenteroides* may offer the potential to carry out fermentations that have a less acidic character which is mentioned earlier and may be desirable from a sensory standpoint. In mixed culture, the final acidity could be substantially reduced by partial conversion of sugar to ethanol and CO₂. A high ratio of volatile/non volatile acids might result in better flavor quality. The lactic acid production in skim milk was chosen as an index of starter performance by Beal and Corrieu (1991).




Fig. 1: Kinetics of CO₂ production by Leuconostoc mesenteroides subsp. dextranicum (■), Lactococcus lactis subsp. lactis (♦) in pure culture in milk and in mixed culture of the 2 strains (•)




Fig. 2: Kinetics of pH evolution by *Leuconostoc* mesenteroides subsp dextranicum (■), *Lactococcus lactis* subsp. lactis (♦) in pure culture in milk and in mixed culture of the 2 strains (•)




Fig. 3: Growth kinetics of Leuconostoc mesenteroides subsp dextranicum (□,■) Lactococcus lactis subsp. lactis (♠, ♦) in pure culture in milk and in mixed culture of the 2 strains

In the first phase, the specific growth rate, lactic acid production rate and evolved CO₂ rate exhibited a constant relationship. After the first 10 h an inhibition of growth, lactic acide production and evolved CO₂, were observed.

Maximum CO_2 production rate (V_{max}) showed a decrease when the inoculum level of *Lactococcus lactis* increases by a factor of 10 from 2.5×10^5 - $2.5 \varsigma \times 10^6$ cfu mL⁻¹ in mixed culture with *Leuconostoc mesenteroides* $(3 \times 10^7 \text{ cfu mL}^{-1})$. This parameter of Vmax could be used to compare different mixed cultures. *Leuconostoc mesenteroides* subsp. *dextranicum* had a maximum rate of CO_2 production of 3 mM h⁻¹ in pure culture. In the presence of $2.5 \times 10^5 \text{ cfu mL}^{-1}$ of *Lactococcus lactis* Vmax of CO_2 production was higher (4 mM h⁻¹) and was obtained quicker in mixed culture than in pure culture,

although, the higher rate of CO₂ production was maintained for a shorter period in pure culture of *Leuconostoc*. *Leuconostoc mesenteroides* did not produced acetoine and diacetyl, the lack of acetoine production at neutral pH is thought to be due to inhibition of acetolactate synthase by several intermediates of sugar metabolism (Cogan, 1987). The pH value of pure culture of *Leuconostoc mesenteroides* decreased from 6.7-5.6 after 10 h, whereas, in mixed culture and pure culture of *Lactococcus lactis*, the pH progressively dropped to 4.4 after 14 h, a value at which growth of both strains ceased. This resulted in the reduction of CO₂ production (Fig 2).

An interesting way to estimate independently Leuconostoc mesenteroides and Lactococcus lactis concentration during mixed cultures is to use the relationships obtained for pure culture with the associated D (-) lactic acid for Leuconostoc mesenteroides and L (+) lactic acid for Lactococcus lactis. Using curves measurements of D (-) and L (+) lactic acid concentration indicate the evolution of the concentration of each strain in mixed cultures which was obtained from the relationships. A linear correlation were established between evolved CO₂ and D (-) lactic acid production by Leuconostoc mesenteroides

The estimated results coming from the pure culture relationships applied to mixed culture experiment are presented in (Fig. 4). A shift between μ max for Leuconostoc and Lactococcus could be appreciated. This shows that Lactococcus lactis grew faster than Leuconostoc mesenteroides due to a more rapid assimilation of nutrients and their proteolytic activity. The same phenomenon is reflected by the specific lactic acid production curves (Gonzalo et al., 1994; Robert et al., 2006). Lactococcus lactis can not consume citric acid (Fig. 5). In mixed culture and pure culture of Leuconostoc mesenteroides, citric acid began to be consumed at the beginning of fermentation, but a 12 mM were entirely consumed within 7 h. As shown in (Fig. 5). Whereas, Lee (2005) has suggested that mixed culturing of Lactobacilli may be more effective than single culturing of Lactobacillus for improving lactic acid production.

The amount of acetic acid formed from citrate was higher in pure and mixed culture of *Leuconostoc mesenteroides*. The kinetics of acetate production and citric acid consumption were similar and a peak of production at early sampling times followed by deceleration of acetate production rate.

Growth of *Leuconostoc mesenteroides* on lactose was stimulated by citrate but no growth on citrate alone was observed. The growth stimulation of *Leuconostoc mesenteroides* in the presence of citrate can be expained by the action of citrate as an external electron acceptor,




Fig. 4: Kinetics of lactic acid production during incubation of *Leuconostoc mesenteroides* subsp dextranicum (□), *Lactococcus lactis* (◊) in pure culture in milk and D-lactic acid (■) and L-lactic acid (♦) production in mixed culture




Fig. 5: Kinetic of citrate consumption (a) and acetate production (b) in culture pure in milk by Leuconostoc mesenteroides subsp dextranicum (□) Lactococcus lactis subsp lactis (◊) and in mixed culture (■)

resulting in more acetate (and ATP) production and less ethanol production during the *Heterofermentative* lactose conversion. This phenomenon has been described by Schmitt *et al.* (1990) and Zurera-Cosano *et al.* (2006).

CONCLUSION

A greater inhibition of evolved CO₂ in *Leuconostoc* mesenteroides coming from the high production of lactic acid in mixed culture by high inoculums of *Lactococcus* lactis (SLO₃ prot⁺) was observed. The growth and CO₂ production by *Leuconostoc* mesenteroides was related to know the precise ratio of initial inoculum of 2 strains milk for good performance in dairy industries.

REFERENCES

Ayad, E.H.E., S. Nashat, N. El-Sadek, H. Metwaly and M. El-Soda, 2004. Selection of wild lactic acid bacteria isolated from traditional Egyptian dairy products according to production and technological. Criteria Food Microbiol., 21: 715-725.

Badis, A., B. Guetarni, Moussa Boudjema, D.E. Henni and M. Kihal 2004. Identification and technological properties of lactic acid bacteria isolated from raw goat milk of 4 Algerian races. Food Microbiol., 21: 579-588.

Beal, C. and G. Corrieu, 1991. Influence of pH, temperature and inoculum composition on mixed cultures of *Streptococcus thermophilus* 404 and *Lactobacillus bulgaricus* 398. Biotechnol. Bioeng., 38: 90-98.

Bellangier, P., J. Richard and C. Foucaud, 1999. Associative growth of lactococcus lactis and Leuconostoc mesenteroides strains in milk. J. Dairy Sci., 80: 1520-1527.

Bjokroth, K.J., R. Geisen, U. Schillinger, N. Weiss, P. DeVos, W.H. Holzapfel, J.H. Korkeala and P. Vandamme, 2000. Characterization of *Leuconostoc* gasicomitatum sp. Associated with spoiled raw tomato Marinated Broiler meat strips packaged under modified atmosphere conditions. Applied Environ. Microbiol., 9 (66): 3764-3772.

Bobillo, M. and V.M. Marshall, 1992. Effect of acidic pH and salt on acid end-products by Lactobacillus plantarum in aerated, glucose-limited continuous culture. J. Applied Bacteriol., 73: 67-70.

Carr, F.J., D. Chill and N. Maida, 2002. The lactic acid bacteria: A literature survey. Critical Rev. Microbiol., 28 (4): 281-370.

Cogan, T.M., 1987. Co-metabolism of citrate and glucose by *Leuconostcoc* sp.: Effet on growth, substrates and products. J. Applied Bacteriol., 63: 551-558.

De Man, M.C., M. Rogosa and M.E. Sharpe, 1960. A medium for the cultivation of lactobacilli. J. Applied Bacteriol., 23: 130-135.

Drosinos, E.H., M. Mataragas, P. Nasis, M. Galiotou and J. Metaxopoulos, 2005. Growth and bacteriocin production kinetics of *Leuconostoc mesenteroides* E131. J. Applied Microbiol., 99: 1314-1323.

- Gibson, T. and Y. Abdelmalek, 1945. The formation of carbon dioxide by lactic acid bacteria and *Bacillus licheniformis* and a cultural method of detecting the process. J. Dairy Res., 14: 35-44.
- Girard, F. and P. Boyaval, 1994. Carbon dioxide measurement in swiss-types cheeses by coupling extraction and gas chromatography. Le Lait., 74: 389-398.
- Gonzalo, A., E. Latrille, C. Beal and G. Corrieu, 1994. On line estimation of biological variables during pH controlled lactic acid fermentations. Biotechnol. Bioeng., 44: 1168-1176.
- Holmes, B., W.E. Sandine and P.R. Elliker 1968. Some factors influencing carbon dioxide production by Leuconostoc citrovorum. Applied Microbiol., 1: 56-61.
- Huang, D.Q., H. Prevost, M. Kihal and C. Diviès, 1994. Instability of plasmid encoding for B-galactosidase in *Leuconostoc mesenteroides* subsp mesenteroides. J. Basic Microbiol., 34 (1): 23-30.
- Kandler, O., 1983. Carbohydrate metabolism in lactic acid bacteria. Antonie Van Leewenhoek, 49: 209-224.
- Kihal, M., H. Prevost, M.E. Lhotte, D.Q. Huang and C. Diviès, 1996. Instability of plasmid-encoded citrate permease in *Leuconostoc*. Lett. Applied Microbiol. (Mars), 22: 219-223.
- Kihal, M., 1996. Etude de la production du dioxyde de carbone par Leuconostoc mesenteroides, élément d'application en technologie fromagère type fromage bleu. Thèse de Doctorat d'état. Université d'Oran Algérie, pp. 167.
- Kihal, M., D.E. Henni, H. Prevost and C. Diviès, 2006. New manometric method for measuring carbon dioxide production by dairy starter culture: A case of *Leuconostoc mesenteroides*. Afr. J. Biotechnol., 5 (4): 378-383.
- Klein G., A. Pack, C. Bonaparte and G. Reuter, 1998. Taxonomy and physiology of probiotic lactic acid bacteria. Int. J. Food Microbial., 41: 103-125.
- Kneifel, W. and R. Gretner, 1992. A simple method for estimating the carbon dioxide product of mesophilic starter cultures. Milchwissenschaft, 11: 108-111.
- Lee, K., 2005. Comparison of fermentative capacities of lactobacilli in single and mixed culture in industrial media. Process Biochemis., 40: 1559-1564.
- Mathot, A.G., M. Kihal, H. Prevost and C. Diviès, 1994. Selective Enumeration of *Leuconostoc* on Vancomycine Agar Media. Int. Dairy J., 5 (4): 459-469.

- Mohr, S.A., W.E. Zottola and G.A. Reineccius, 1993. The use of gas chromatography to measure carbon dioxide production by starter cultures. J. Dairy Sci., 76: 3350-3353.
- Padmanabhan, P.A. and D.S. Kim, 2002. Production of insoluble dextran using cell-bound dextransucrase of Leuconostoc mesenteroides NRRL B-523. Carbohydrate Res., 337: 1529-1533.
- Perea, M. Velez, K. Hermans, T.L.A. Verhoeven, S.E. Lebeer, J. Vanderleyden and S.C.J. De Keersmaecker, 2007. Identification and characterization of starter lactic acid bacteria and probiotics from Columbian dairy products. J. Applied Microbiol., 32: 1-9.
- Robert, H., V. Gabriel, D. Lefebvre, P. Rabier, Y. Vayssier and C.F. Fontagne, 2006. Study of the behaviour of *Lactobacillus plantarum* and *Leuconostoc* starters during a complete wheat sourdough breadmaking process. LWT, 39: 256-265.
- Schmitt, P., C. Diviès and C. Merlot, 1990. Origin of end products from the metabolism of glucose and citrate by *Leuconostoc mesenteroides* subsp. cremoris. Applied Microbiol. Biotechnol., 36: 679-683.
- Stiles, M.E. and W.H. Holzapfel, 1997. Lactic acid bacteria of foods and their current taxonomy. Int. J. Food Microbiol., 36: 1-29.
- Thapa, N., J. Pal and J.P. Tamang, 2006. Phenotypic identification and technological properties of lactic acid bacteria isolated from traditionally processed fish products of the Eastern Himalayas. Int. J. Food Microbiol., 107: 33-38.
- Terzaghi, B.E. and W.E. Sandine, 1975. Improved medium for lactic streptococci and their bacteriophages. Applied Microbiol., 29: 807-813.
- Todorov, S.D. and L.M.T. Dicks, 2005. Production of bacteriocin ST33LD, produced by *Leuconostoc mesenteroides* subsp. *mesenteroides*, as recorded in the presence of different medium components. World J. Microbiol. Biotechnol., 21: 1585-1590.
- Zurera-Cosano, G., R.M., Garcý a-Gimeno, R. Rodrýguez-Perez and C. Hervas-Martýnez, 2006. Performance of response surface model for prediction of *Leuconostoc mesenteroides* growth parameters under different experimental conditions. Food Control, 17: 429-438.