

## Microsporidia in Stools from Cancer Patients

R. Angela and K. Suresh

Department of Parasitology, Faculty of Medicine,  
 University of Malaya, 50603 Kuala Lumpur, Malaysia

### RAPID COMMUNICATION

Microsporidia are widespread obligate intracellular parasites which have been recently more akin to degenerate fungi than to protozoa (Gill and Fast, 2006). They are found widely widespread in invertebrates and vertebrates but was rarely seen in humans until the AIDS epidemic (Schwartz *et al.*, 1996). *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis* are the most common species associated in opportunistic infections seen in immunosuppressed individuals such as transplant-recipients (Kelkar *et al.*, 1997; Orenstein *et al.*, 2005), AIDS patients and occasionally in immunocompetent individuals in the form of travellers' diarrhea (Field, 2002).

There have been studies associating parasitic infections in cancer patients such as cryptosporidium (Noureldin *et al.*, 1999; Baqai *et al.*, 2005; Bialek, 2005), visceral leishmaniasis (Sah *et al.*, 2002; Bialek, 2005), liver flukes (Hughes *et al.*, 2006), *Blastocystis hominis* (Noureldin *et al.*, 1999; Tasova *et al.*, 2000), *Strongyloides stercoralis* (Reddy *et al.*, 1983; Dini *et al.*, 1987; Bialek, 2005) and *Toxoplasma gondii* (Klastersky, 1985; Bialek, 2005). In Malaysia there were reports of extra-hepatic cholangiocarcinoma co-existing with biliary ascariasis which were seen in 2 cases involving rural Malay women (Lim and Selliah, 1994) and pediatric cancer patients receiving chemotherapy with parasitic infections (Menon *et al.*, 1999). Besides scanty reports of microsporidia in cancer patients (Yazar *et al.*, 2003, Botero *et al.*, 2003) there has been no report of microsporidia seen in cancer patients in Malaysia despite its occurrence in HIV population. The present study is pivotal in establishing the occurrence of microsporidia in cancer patients and to assess if the organism is opportunistic when compared to the prevalence of microsporidia in normal population.

There hundred and eleven and 144 fresh stool specimens were collected from the oncology clinics of three hospitals and from a normal healthy population respectively over a period of 2 years. All specimens were randomly obtained from confirmed cancer patients undergoing chemo-and/or radiotherapy, s who visited the

oncology clinic. The normal health population were renadomly selected from a community during a local medical camp organized by volunteers. From the cancer patient population, 36.0% (112/311) were males and 64.0% (199/311) were females. \_\_\_\_\_ health population? Specimens were concentrated with a water-ether sedimentation method as described by Van Gool *et al.* (1994). A thin smear was made before being stained with modified trichrome stain as described by Weber *et al.* (1992). The result was examined by light microscopy under oil immersion. At least 100 fields were examined, in duplicate, before a specimen was declared negative, as recommended by Bendall and Chiodini (1993).

21.9% (68/311) were found to be positive for microsporidia. 69.2% (47/68) were from females, while 30.88% (21/68) were from males. The difference in infection between genders was not statistically significant (female vs. male = 47/199 vs. 21/112, p-value = 1). 7.36% (5/68) of the positive specimens also had co- or multiple infections with other gastrointestinal parasites such as *Blastocystis hominis*, *Giardia* sp. *Dientamoeba fragilis*, *Ascaris lumbricoides* and *Trichuris trichura*, while 92.64% (63/68) were single infections. 29.41% (20/68) of these patients had been newly diagnosed and/or had not undergone any treatment, while 70.59% (48/68) had already undergone chemo-and/or radiotherapy at the time the specimens were collected.

The organisms were visible against a clear background as ovoid or ellipsoid spores, measuring 1.0-1.5  $\mu\text{m}$ , that were brightly outlined in dark pink, with the polar zones staining slightly darker than the centres. While a light pink central vacuolar zone was seen in most of the spores that were detected, an extremely faint, belt-like stripe was observed in other spores, girdling the equatorial region (Fig. 1 and 2). This is an identifying characteristic of *Microsporidia* sp. the polar filament by which the parasite uses to infect other cells with its sporoplasm.

Only 4.41% (3/68) of the positive specimens were diarrhetic, suggesting that most microsporidial infections are asymptomatic. One patient had undergone 11 cycles of therapy for colorectal cancer, while the other had




Fig. 1: Fresh stool specimen from a cancer patient with microsporidiosis. Spores are numerous and widely scattered throughout the specimen. A faint polar filament is visible in one of the spores (arrow). X scale




Fig. 2: Positive stool specimen showing ellipsoid-shaped spores with dark pink outlines and polar filament (arrow)

undergone 2 cycles of treatment for a cancer that was not specified in the patient data. The remaining patient had just been diagnosed with breast cancer and had not begun treatment.

While the majority of positive specimens were from individuals who had undergone therapy, the difference between the pre-and post-treatment groups was not statistically significant (infected pre-treatment vs. infected post-treatment = 20/103 vs. 48/208; p-value = 1). However, when the findings were compared with the results of screening in a normal healthy population (a control group, results not shown), the difference was statistically significant (normal vs. cancer = 5/144 vs.

68/217; p-value = 0.001). It has been reported that immunosuppressive treatment increases the probability of acquiring parasitic infections, generally with a high degree of severity (Smith *et al.*, 1988; Manheimer and Soave, 1994; Rotterdam and Tsang, 1994). This is because the immunocompromised cellular or humoral responses that have undergone qualitative and/or quantitative alterations (due to immunosuppressive therapy) that impede them from acting efficiently against the infections, which is manifested in the deterioration of their overall condition (Botero *et al.*, 2003). This finding concurs with previous studies that microsporidia are opportunistic as evidenced by a larger proportion of patients after chemotherapy having a higher numbers of spores (Connolly *et al.*, 1988; Smith *et al.*, 1988; Souza-Dias *et al.*, 1988; Manheimer and Soave, 1994).

CD4+ T-lymphocyte counts were not established in this study, but previous studies showed that cancer patients infected with microsporidia had CD4+T-lymphocyte levels of less than 50 cells mm<sup>-3</sup> (Botero *et al.*, 2003). As the specimens were collected in random fashion, a more controlled study involving the collection of stool specimens from a patient from the moment of diagnosis, before and after each cycle of treatment, right up to the end of the treatment, as well as from a follow-up visit to the clinic would yield more thorough data on such opportunistic infections.

To the best of our knowledge, this is the first report of microsporidial infection in Malaysian cancer patients and the study confirms that it is imperative to include microsporidia in the screening of opportunistic pathogens in stools especially from cancer and other immunosuppressed patients as the spores could contribute to complications.

## REFERENCES

- Baqai, R., S. Anwar and S.U. Kazmi, 2005. Detection of cryptosporidium in immunosuppressed patients. J. Ayub. Med. Coll. Abbottabad, 17: 38-40.
- Bendall, R.P. and P.L. Chiodini, 1993. New diagnostic methods for parasitic infections. Curr. Opin. Infect. Dis., 6: 318-323.
- Bialek, R., 2005. Parasitic diseases in pediatric cancer patients. Klin. Padiatr., pp: S85-90.
- Botero, J.H., A. Castaño, M.N. Montoya, N.E. Ocampo, M.I. Hurtado and M.M. Lopera, 2003. A preliminary study of the prevalence of intestinal parasites in immunocompromised patients with and without gastrointestinal manifestations. Rev. Inst. Med. Trop. S. Paulo., 45: 197-200.

- Connolly, G.M., M.S. Dryden, D.C. Shanson and B.G. Gazzard, 1988. Cryptosporidial diarrhea in AIDS and its treatment. *Gut.*, 29: 593-597.
- Dini, D., F. Rivasi, M. Federico, C. Casolari, F.A. Barbieri, 1987. Case of Hodgkin's disease complicated by *Strongyloides stercoralis* infestation. *Minerva Med.*, 78: 49 - 53.
- Field, A.S., 2002. Light microscopic and electron microscopic diagnosis of gastrointestinal opportunistic infections in HIV-positive patients. *Pathology*, 34: 21-35.
- Gill, E.E. and N.M. Fast, 2006. Assessing the microsporidia-fungi relationship: Combined phylogenetic analysis of eight genes. *Gene.*, 375: 103 -109.
- Hughes, N.R., C. Pairojkul, S.G. Royce, A. Clouston and P.S. Bhathal, 2006. Liver fluke-associated and sporadic cholangiocarcinoma: An immunohistochemical study of bile duct, peribiliary gland and tumour cell phenotypes. *J. Clin. Pathol.*, 59: 1073 -1078.
- Kelkar, R., P.S. Sastry, S.S. Kulkarni, T.K. Saikia, P.M. Parikh and S.H. Advani, 1997. Pulmonary microsporidial infection in a patient with CML undergoing allogeneic marrow transplant. *Bone Marrow Transplant.*, 19: 179-182.
- Klustersky, J., 1985. Infections in immunocompromised patients. I. Pathogenesis, etiology and diagnosis. *Clin. Ther.*, 8: 90-99.
- Lim, K.G. and S.P. Selliah, 1994. Biliary ascariasis and extrahepatic cholangiocarcinoma: A report of two cases. *Sing. Med. J.*, 35: 400-402.
- Manheimer, S.B. and R. Soave, 1994. Protozoal infections in patients with AIDS. Cryptosporidiosis, isosporiosis, cyclosporiasis and microsporidiosis. *Infect. Dis. Clin. Infect. North Amer.*, 8: 483- 498.
- Menon, B.S., M.S. Abdullah, F. Mahamud and B. Singh, 1999. Intestinal parasites in children with cancer. *J. Trop. Pediatr.*, 45: 241-242.
- Noureldin, M.S. A.A. Shaltout, E.M. El Hamshary and M.E. Ali, 1999. Opportunistic intestinal protozoal infections in immunocompromised children. *J. Egypt Soc. Parasitol.*, 29: 951-961.
- Orenstein, J.M., P. Russo, E.S. Didier, C. Bowers, N. Bunin and D.T. Teachey, 2005. Fatal pulmonary microsporidiosis due to *Encephalitozoon cuniculi* following allogeneic bone marrow transplantation for acute myelogenous leukemia. *Ultrastruct. Pathol.*, 29: 269 - 276.
- Reddy, K.R., A.R. Laurain and E. Thomas, 1983. Strongyloidiasis. When to suspect the wily nematode. *Postgrad. Med.*, 74: 273 -275, 279-282.
- Rotterdam, H. and P. Tsang, 1994. Gastrointestinal disease in the immunocompromised patient. *Hum. Pathol.*, 25: 1123-1140.
- Sah, S.P., S. Rijal, P.P. Bhadani, S. Rani and S. Koirala, 2002. Visceral leishmaniasis in two cases of leukemia. *SEA. J. Trop. Med. Pub. Health*, 33: 25 - 27.
- Schwartz, D.A., I. Sobottka, G.J. Leitch, A. Cali and G.S. Visvesvara, 1996. Pathology of microsporidiosis. Emerging parasitic infections in patients with acquired immunodeficiency syndrome. *Arch. Pathol. Lab. Med.*, 120: 173-188.
- Smith, P.D., H.C. Lane and V.J. Gill *et al.*, 1988. Intestinal infection in patients with the Acquired Immunodeficiency Syndrome (AIDS). Etiology and response to therapy. *Ann. Int. Med.*, 108: 328- 333.
- Souza-Dias, R., A.C. Mangini and D. Torres *et al.*, 1988. Cryptosporidiosis among patients with acquired immunodeficiency syndrome (AIDS) in the county of São Paulo, Brazil. *Rev. Inst. Trop. Med. S. Paulo.*, 30: 310-312.
- Tasova, Y., B. Sahin, S. Koltas and S. Paydas, 2000. Clinical significance and frequency of *Blastocystis hominis* in Turkish patients with hematological malignancy. *Acta Med. Okayama*, 54: 133-136.
- van Gool, T., E.U. Canning and J. Dankert, 1994. An improved practical and sensitive technique for the detection of microsporidian spores in stool samples. *Trans. R. Soc. Trop. Med. Hyg.*, 88: 189-190.
- Weber, R., R.T. Bryan and R.L. Owen, 1992. Improved light-microscopical detection of microsporidia in stool and duodenal aspirates. *N. Eng. J. Med.*, 326: 161-165.
- Yazar, S., B. Eser, S. Yalcin, I. Sahin, A.N. Koc, 2003. A case of pulmonary microsporidiosis in an Acute Myeloblastic Leukemia (AML)-M3 patient. *Yonsei Med. J.*, 44: 146-149.