

## Anti-Proliferation of MDA-MB-231 Human Breast Tumour Cells by Arsenic Trioxide via Induction of Apoptosis

<sup>1</sup>Lin-Li Zhou, <sup>1,2</sup>Judy Yuet-Wa Chan, <sup>1</sup>Jun Wang and <sup>1,2</sup>Kwok-Pui Fung

<sup>1</sup>Department of Biochemistry, <sup>2</sup>Institute of Chinese Medicine,  
The Chinese University of Hong Kong, Shatin, N.T., Hong Kong, SAR, China

**Abstract:** Arsenic Trioxide (As<sub>2</sub>O<sub>3</sub>) has been explored for its use as medicine in both Western and Chinese societies. In 1990s, As<sub>2</sub>O<sub>3</sub> was reported to treat patients with acute promyelocytic leukemia (APL). Until recently, research on other solid tumour cells was emerged. In the present study, the mechanism of As<sub>2</sub>O<sub>3</sub> treatment of human breast tumour MDA-MB-231 cells was investigated. It was found that As<sub>2</sub>O<sub>3</sub> inhibited the cell proliferation of MDA-MB-231 cells in a time- and dose-dependent manner. Mechanistic study indicated that the inhibition was induced via cell cycle arrest and apoptosis. As<sub>2</sub>O<sub>3</sub> induced apoptosis via both extrinsic and intrinsic apoptotic pathways by regulating the pro- or anti-apoptotic molecules. Moreover, As<sub>2</sub>O<sub>3</sub>-induced cell cycle arrested at G<sub>2</sub> phase in MDA-MB-231 cells. The study revealed that As<sub>2</sub>O<sub>3</sub> was a potent candidate for further investigation for combating against human breast tumour including the late stage breast tumour.

**Key words:** Arsenic trioxide, breast tumour, MDA-MB-231, apoptosis

### INTRODUCTION

Breast tumour is the fifth most common tumours in the world. Estrogens have been found to be the primary stimulant of the breast tumour. Mechanistic study demonstrated that estrogen promoted cell proliferation and prevented apoptosis of breast tumour cells (Lippman *et al.*, 1976; Umans *et al.*, 1984; Hajek *et al.*, 1997). The anti-estrogen drug for breast tumour treatment that has been used most often is tamoxifen. Tamoxifen is also used to treat metastatic breast tumour and to prevent the development of breast tumour in women at high risk. Tamoxifen works as an anti-estrogen to treat human breast tumour and prevents the induction and growth of estrogen-receptor (ER) positive carcinogen-induced rat mammary carcinomas (Jordan, 1976). Tamoxifen is effective in pre-menopausal women but is ineffective in ER negative patients (i.e., advanced breast tumour patients). However, tamoxifen may cause some severe side effects, including blood clots (thrombosis), endometrial tumour (tumour of the uterine lining), abnormal growth of uterine tissue (endometriosis), stroke and fertility issues (Osborne, 1998; Clemons *et al.*, 2002). Alternative treatment for breast tumour is urgently needed.

In early 1970s, a group of researchers from Harbin Medical University in Northeast China found that intravenous administration of arsenic trioxide (As<sub>2</sub>O<sub>3</sub>) with relatively small doses (10 mg day<sup>-1</sup>) was effective in treating patients with acute promyelocytic leukemia (APL) (Zhang *et al.*, 1998; Li *et al.*, 1988; Sun *et al.*, 1992). Then, Sun *et al.* (1992) reported promising results of 32 cases of APL treated with As<sub>2</sub>O<sub>3</sub>. Later, a trial performed at the Shanghai Second Medical University indicated that complete remission could be achieved in 14 of 15 investigated patients that had relapsed after prior treatment with all-trans retinoic acid (ATRA) or conventional chemotherapy (Shen *et al.*, 1997). In Western countries, the population clinical efficacy of As<sub>2</sub>O<sub>3</sub> was proven by studies performed by Soignet *et al.* (1998, 2001). Recently, the drug Trisenox™ was formally approved by the State Drug Administration in China (1999) and the Food and Drug Administration of the United States (2000). Apart from APL, As<sub>2</sub>O<sub>3</sub> were reported to inhibit growth and promote apoptosis in other hematological tumour cell lines (Konig *et al.*, 1997). As<sub>2</sub>O<sub>3</sub> also exhibits anti-proliferative effect and induces apoptosis on solid tumour cells such as prostate tumour, renal tumour, cervical tumour and hepatoma (Ling *et al.*, 2002; Li *et al.*, 2004). We recently found that As<sub>2</sub>O<sub>3</sub> is

effective in suppressing growth of ER $\alpha$  positive human breast tumour cell line MCF-7 (Chow *et al.*, 2004a, b), which mimic the early stage of breast tumour.

In the study, we further investigate the effects of As<sub>2</sub>O<sub>3</sub> on another human breast tumour cell line, MDA-MB-231, which is ER $\alpha$  negative, mimicking the late stage of human breast tumour. The aim of this study is to explore the potential anti-tumour effect of As<sub>2</sub>O<sub>3</sub> in MDA-MB-231 cells and to study the underlying mechanisms.

## MATERIALS AND METHODS

**Cell culture and chemicals:** MDA-MB-231 cell line was purchased from American Type Culture Collection (ATCC, Rockville, MD, USA). The cells were maintained in RPMI 1640 medium, which was supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin. All of these reagents were purchase from Invitrogen, USA. The cells were maintained at 37°C in a humidified atmosphere of 5% carbon dioxide (CO<sub>2</sub>).

Arsenic Trioxide (As<sub>2</sub>O<sub>3</sub>) was purchased from Sigma Chemical Company. Stock As<sub>2</sub>O<sub>3</sub> solution was prepared by dissolving powder As<sub>2</sub>O<sub>3</sub> in boiled phosphate buffered saline (PBS) at 10 mM. Boiling was continued until As<sub>2</sub>O<sub>3</sub> powder was completely dissolved. The stock solution was sterilized using 0.22  $\mu$ M syringe filter (Millipore) and stored at -20°C.

**Cell viability assay:** Total 1 $\times$ 10<sup>4</sup> cells/well were seeded onto a 96-well plate in RMPI 1640 medium. After 24 h incubation at 37°C, the cells were treated with various concentrations of As<sub>2</sub>O<sub>3</sub> in a fresh medium. After 24, 48 or 72 h, the medium in each well was removed and washed with PBS. Negative control was prepared by treating cells with me dium alone. Then, 30  $\mu$ L of MTT solution (5 mg mL<sup>-1</sup>) was added to each well and the plate was incubated at 37°C for 3 h. After incubation, MTT solution was discarded and 100  $\mu$ L DMSO was added to dissolve the crystals in the cells. Then, absorbance at 540 nm was measured using an ELISA plate reader (BIO-RAD). Percentage cell survival of each treatment was calculated as:

$$\text{Percent cell survival} = (\text{O.D. treatment} / \text{O.D. negative control}) \times 100\%$$

**Detection of DNA fragmentation:** Total 5 $\times$ 10<sup>5</sup> cells/well were seeded onto 60 mm culture dish and incubated at 37°C, 5% CO<sub>2</sub>. After appropriate treatment with As<sub>2</sub>O<sub>3</sub>, cells were lysed with 400  $\mu$ L of DNA lysis buffer and vortexed until no cell debris was left. Twenty microliter of 10 mg mL<sup>-1</sup> proteinase K was added and incubated at 37°C for 2 h. Then, 150  $\mu$ L of saturated NaCl was added

and the sample was shaken vigorously. The mixture was centrifuged at 6500 $\times$  g for 15 min. The supernatant was collected; 1 mL of cold ethanol was added and centrifuged again at 15000 $\times$ g for 20 min. After rinsing with cold 75% ethanol, the pellet was dried at room temperature. Finally, 20  $\mu$ L of RNase A solution in Tris-EDTA buffer (0.2 mg mL<sup>-1</sup>) was added to each sample and further incubated at 37°C for 90 min. Equal amount of sample was electrophoresed in 1.5% agarose gel and visualized by ethidium bromide staining.

**Cell cycle analysis:** Total 3 $\times$ 10<sup>5</sup> cells/well were seeded onto 6-well culture plate and incubated at 37°C, 5% CO<sub>2</sub>. Twenty-four h later, the medium was discarded and appropriate concentrations of As<sub>2</sub>O<sub>3</sub> were added. After incubation for 72 h, the cells were harvested and washed with PBS. The cells were then fixed with 1 mL of 70% ethanol at 4°C overnight. After fixation, the cells were centrifuged at 3000  $\times$  g for 5 min. to remove the ethanol. Cells were then resuspended in 0.46 mL freshly prepared propidium iodide solution containing propidium iodide (43  $\mu$ g mL<sup>-1</sup>) and RNase A (1 mg mL<sup>-1</sup>) and incubated in dark at 37°C for 30 min. After incubation, the cells were analyzed by FACSsort flow cytometer (Becton Dickinson) (Chow *et al.*, 2004a).

**Detection of apoptotic cells by Annexin V-PI staining:** The detection of apoptosis was performed by using BD Pharmingen™ Annexin V-FITC kit (Bio-Gene Technology Ltd.). Total 5 $\times$ 10<sup>5</sup> cells were seeded in a 60 mm culture dish. Twenty four hour later, medium was discarded and As<sub>2</sub>O<sub>3</sub> was added with fresh medium. Then after 72 h, cells were harvested and washed with PBS. And 1 $\times$ 10<sup>5</sup> cells were used for the assay. Ten micro liters of 10 $\times$ binding buffer, 10  $\mu$ L of PI, 3  $\mu$ L of Annexin V-FITC conjugate and 77  $\mu$ L of dH<sub>2</sub>O were added to a sample and incubated in the dark at room temperature for 15 min. After incubation, 400  $\mu$ L of 1  $\times$  binding buffer was added to the sample and the sample was analyzed by FACSsort flow cytometer (Becton Dickinson) as soon as possible. A computer program, WinMDI, was used for data analysis (Chow *et al.*, 2004a).

**Western blot analysis:** Total 1 $\times$ 10<sup>6</sup> cells were seeded on 100 mm culture dish with 10 mL RPMI 1640 medium. Twenty-four hour later, appropriate concentrations of As<sub>2</sub>O<sub>3</sub> were added. After 72 h, cells were collected and washed twice with PBS. Then, 100  $\mu$ L of lysis buffer was added and allowed to stand on ice for 2 h. After that, samples were boiled for 10 min and centrifuged at 13600  $\times$  g for 10 min at 4°C. Finally, the supernatant was collected and protein concentration of each sample was determined

by BCA assay. Equal amount (50 µg) of each protein sample was dissolved in 12.5% SDS-PAGE. The protein was then transferred to PVDF membrane by an electro-blotter (Bio-Rad). The membrane was blocked with 10% non-fat milk for 2 h. Then, appropriate amount of each of primary antibody was added and incubated at 4°C overnight. After that, the membrane was washed with TBS-T for 3 times and probed with secondary antibody (conjugated with horseradish peroxidase) at room temperature for 1 h. The membrane was washed again with TBS-T for 3 times and finally the signal was detected using Enhanced Chemiluminescence (ECL) detection kit and developed on X-ray film (Chow *et al.*, 2004a).

## RESULTS DISCUSSION

**Anti-proliferative effects of As<sub>2</sub>O<sub>3</sub> on MDA-MB-231 cells:** MDA-MB-231 cells were treated with various concentrations of As<sub>2</sub>O<sub>3</sub> for 24, 48 and 72 h, respectively. After treatment, MTT assay was performed. The results were shown in Fig. 1. IC<sub>50</sub> values when MDA-MB-231 cells were treated with As<sub>2</sub>O<sub>3</sub> for 24, 48 and 72 h were found to be 20, 18.4 and 12.1 µM, respectively.

**DNA fragmentation induced by As<sub>2</sub>O<sub>3</sub> treatment:** MDA-MB-231 cells were incubated with various concentrations of As<sub>2</sub>O<sub>3</sub> for 72 h. After treatment, the DNA contents were extracted and separated on 1.5% agarose gel. Laddering pattern was observed when the MDA-MB-231 cells were treated with 6.25, 12.5 and 25 µM As<sub>2</sub>O<sub>3</sub>, respectively, for 72 h (Fig 2).

**Increase in sub-G<sub>1</sub> phase by As<sub>2</sub>O<sub>3</sub> treatment:** Cell cycle analysis was performed after MDA-MB-231 cells were




Fig 1: The anti-proliferative effect of As<sub>2</sub>O<sub>3</sub> on MDA-MB-231 cells. MDA-MB-231 cells were treated with various concentrations of As<sub>2</sub>O<sub>3</sub> for 24, 48 or 72 h. Data was expressed in mean±standard deviation of four replicates

treated with different concentrations of As<sub>2</sub>O<sub>3</sub> for 72 h. The results showed an increase in sub-G<sub>1</sub> phase upon As<sub>2</sub>O<sub>3</sub> treatment indicating that As<sub>2</sub>O<sub>3</sub> might inhibit the growth of MDA-MB-231 cells by inducing apoptosis. Increase in S and G<sub>2</sub>/M phase observed from the results indicated that cell cycle arrest occurred in As<sub>2</sub>O<sub>3</sub>-treated MDA-MB-231 cells (Fig 3).

**Increase in percentage of apoptotic cells by As<sub>2</sub>O<sub>3</sub> treatment:** From Fig 4, the percentage of apoptotic cells as shown in the lower right quadrant of the dot plot increased from 6.5-12.5% after treatment with 25 µM of As<sub>2</sub>O<sub>3</sub> for 72 h.

**Regulation of apoptosis-related proteins by As<sub>2</sub>O<sub>3</sub> treatment:** The MDA-MB-231 cells were treated with various concentrations of As<sub>2</sub>O<sub>3</sub> for 72 h. Then, Western bolt analysis of apoptosis-related proteins was performed. There is an obvious decrease in expression level of bcl-2 while there is no significant change in expression level of bax protein (Fig 5a and b). Results of Fig 5 also showed an increase of cytochrome c released to cytosol while the expression level of pro-caspase 3, pro-caspase 8 and pro-caspase 9 decreased after treatment with As<sub>2</sub>O<sub>3</sub> (Fig 5c-f). Expression level of FasL was up-regulated while that of p53 was down-regulated in the MDA-MB-





Fig 2: DNA fragmentation induced by As<sub>2</sub>O<sub>3</sub> on MDA-MB-231 cells. DNA prepared from MDA-MB-231 cells after treated with As<sub>2</sub>O<sub>3</sub> for 72 h. Lane 1 is negative control without As<sub>2</sub>O<sub>3</sub> treatment. Lanes 2, 3, 4 and 5 were treatment with 3.125 µM, 6.25 µM, 12.5 µM and 25 µM of As<sub>2</sub>O<sub>3</sub>, respectively. M is 100 basepair marker. Laddering patterns were observed at concentrations 6.25 µM, 12.5 µM and 25 µM. This figure showed a representative of three independent trials



As <sub>2</sub> O <sub>3</sub> (μM)	Percentage of cell population (%)			
	Sub-G <sub>1</sub>	G <sub>0</sub> /G <sub>1</sub>	S	G <sub>2</sub> /M
0	7.94	63.65	11.77	17.25
3.125	8.03	56.34	17.03	19.26
6.25	9.33	49.03	19.97	22.57
125	9.83	47.57	18.56	25.16
25	14.59	36.04	21.80	28.79

Fig. 3: Cell cycle analysis of MDA-MB-231 cells treated with As<sub>2</sub>O<sub>3</sub> for 72 h. (Upper panel) 3×10<sup>5</sup> cells were seeded in each well of 6-well plates. After treated with various concentrations of As<sub>2</sub>O<sub>3</sub> for 72 h, the cells were stained with PI. The populations of cells in sub-G<sub>1</sub>, G<sub>0</sub>/G<sub>1</sub>, S and G<sub>2</sub>/M phases were determined by FACSsort flow cytometer and analysed by WinMIDI (Lower panel). The percentage of MDA-MB-231 cells distributions in different cell cycle were summarized in the table. This figure showed a representative of three independent trials

231 cells treated with As<sub>2</sub>O<sub>3</sub> when compared with the untreated cells (Fig. 5g and h).

As shown by the cell cycle analysis by PI staining (Fig. 3), there was an increase in cell population in G<sub>2</sub>/M phase after MDA-MB-231 cells were treated with different concentration of As<sub>2</sub>O<sub>3</sub>, indicating that As<sub>2</sub>O<sub>3</sub> induced a cell cycle arrest at G<sub>2</sub>/M phase. The expression level of cyclin B was examined by Western blotting. The result showed that there was a marked increase in expression level of cyclin B with increasing concentrations of As<sub>2</sub>O<sub>3</sub>. For cyclin E, there is no significant change in the protein expression level (Fig. 5i and j).

In addition to the significant anti-tumour effect on acute promyelocytic leukemia (APL), As<sub>2</sub>O<sub>3</sub> has been studied on its potential use in other types of leukemia as well as solid tumours. In the previous study of our group, As<sub>2</sub>O<sub>3</sub> was found to be able to inhibit the proliferation of ERα positive MCF-7 cells *in vitro* (Chow *et al.*, 2004a). Further mechanistic study indicated that As<sub>2</sub>O<sub>3</sub> induced

apoptosis and cell cycle arrest on MCF-7 cells and these effects were found to be related to the regulation of ERα signaling pathway (Chow *et al.*, 2004a). Patients of breast tumour at late stage always have their breast tissues contain ERα negative cells. To study the anti-tumour effect of As<sub>2</sub>O<sub>3</sub> on late stage breast tumour, MDA-MB-231 cell line, an ERα negative breast tumour cell line, was used to mimic the late stage of breast tumour in this study. Our results indicated that the As<sub>2</sub>O<sub>3</sub> could inhibit MDA-MB-231 cell growth in a dose and time dependent manner (Fig. 1).

Induction of apoptosis and cycle arrest are the most common mechanisms studied on the chemotherapeutic drugs in recent years. As<sub>2</sub>O<sub>3</sub> was reported to induce apoptosis in ATRA-resistant APL cells (Shao *et al.*, 1998; Kinjo *et al.*, 2000). Apart from apoptosis, regulation of cell cycle is another important mechanism involved in the anti-tumour effect of As<sub>2</sub>O<sub>3</sub> in solid tumour cells (Park *et al.*, 2003; Wu *et al.*, 2004). In our study, As<sub>2</sub>O<sub>3</sub> at




Fig. 4: Percentage of apoptotic cells detected by PI and Annexin V-FITC staining. Cells ( $3 \times 10^5$ ) were seeded in each well of 6-well plates. After treated with  $As_2O_3$  for 72 h, the cells were stained with PI and Annexin V-FITC conjugate. A shift of cell population to the right bottom quadrant was observed after  $As_2O_3$  treatment. This figure showed a representative of three independent trials

concentration more than  $3.125 \mu M$  induced apoptosis in MDA-MB-231 cells. The apoptosis was preliminarily proven by the typical DNA ladder (Fig. 2). To further confirm the result, we examined the externalization of PS by Annexin-V and PI staining. An increase in apoptotic cell population was observed after  $As_2O_3$  treatment (Fig. 4). Together with the detection of increase in sub- $G_1$  phase during cell cycle analysis (Fig. 3), it was suggested that  $As_2O_3$  could inhibit the proliferation of MDA-MB-231 cells via induction of apoptosis. Meanwhile, the increase of cell population in  $G_2/M$  phase indicated that  $As_2O_3$  induced  $G_2/M$  cell cycle arrest in MDA-MB-231 cells (Fig. 3).

In previous study,  $As_2O_3$  has been found to induce apoptosis in different human tumour cell lines by regulating the expression level of apoptosis related proteins. However, detailed mechanisms involved were quite different. Chow *et al.* (2004a) reported that  $As_2O_3$ -induced apoptosis in ER $\alpha$  positive breast tumour cell line MCF-7 was due to the collapse of mitochondrial membrane potential, up-regulation of tumour suppressor gene p53 and regulation of ER $\alpha$  signaling pathway. As the MDA-MB-231 cells are known to be ER $\alpha$  negative, the effect of  $As_2O_3$  in these cells does not involve ER $\alpha$ . Also, the features of MCF-7 cells and MDA-MB-231 cells are

quite different. Caspase 3, a key factor for initiation of DNA fragmentation was not expressed in MCF-7 cells but expressed in MDA-MB-231 cells. In our study, the involvement of extrinsic apoptotic pathway in the apoptosis in MDA-MB-231 cells treated by  $As_2O_3$  was confirmed by the findings that the expression level of Fas receptor Ligand (FasL) was increased, caspase 8 was activated and the ratio of bax/bcl-2 was increased. The increased level of cytochrome c released to cytosol and activation of caspase 9 suggested the involvement of intrinsic apoptotic pathway. p53 was involved in both cell cycle arrest and apoptosis. p53 was found to be up-regulated and induced apoptosis through regulating bax protein (Miyashita and Reed, 1995). From our study, the expression level of p53 was decreased which indicated that p53 did not directly trigger apoptosis in MDA-MB-231 cells. Since the function of p53 is cellular repair from DNA damage, lacking of p53 may cause prolonged DNA damage in the MDA-MB-231 cells. DNA damage may then activate the intrinsic apoptotic pathway by down-regulating the expression level of bcl-2 (Fig. 5).

The induction of cell cycle arrest on various tumour cell lines by  $As_2O_3$  is mostly related to the arrest in  $G_0/G_1$  and  $G_2/M$  phase. Cyclin B1 associates with cdk1 and remains inactive in cytosol until it is activated by




Fig. 5: Expression level of apoptosis and cell cycle arrest related proteins in MDA-MB-231 cells were examined by Western blot analysis. Antibodies probing various proteins were used (A-J) and  $\beta$ -actin was used to normalize the protein contents in each sample. Lane 1 is negative control without  $As_2O_3$  treatment. Lanes 2, 3, 4 and 5 were treatment with 3.125, 6.25, 12.5 and 25  $\mu M$  of  $As_2O_3$ , respectively. This figure showed a representative of three independent trials

phosphorylation and cyclinB-cdk1 induced  $G_2/M$  phase arrest by prevention of the dephosphorylation of cyclinB-cdk1 complex (Halloran and Fenton, 1998). In our study, the changes in expression levels of cyclin-B1 and p53 indicated that  $As_2O_3$  induced  $G_2/M$  phase cell cycle arrest via prevention of dephosphorylation of cyclinB-cdk1 complex. Cyclin E-cdk2 regulates the transition from  $G_1$  to S phase and p53 tumour suppressor gene inhibits the cyclin E-cdk2 activity after DNA damage (Di Leonardo *et al.*, 1994). Our results showed that there was a significant decrease in percentage of MDA-MB-231 cells in  $G_1$  phase, together with the decrease of p53 expression and no significant change of cyclin E expression in  $As_2O_3$  treated MDA-MB-231 cells (Fig. 5), indicating that the p53 may not be involved in  $As_2O_3$ -induced cell cycle arrest.

For thousands of year,  $As_2O_3$  was considered as a toxic agent in both Chinese and Western society. Pharmacokinetic study performed by Shen *et al.* (1997) during the treatment of APL suggested that it is safe to

inject  $As_2O_3$  with a dose of 10 mg day<sup>-1</sup> because arsenic was rapidly eliminated in the plasma and continuous administration of  $As_2O_3$  did not result in the accumulation of arsenic in plasma. The growth inhibition of  $As_2O_3$  on normal human 184B5 breast cells was studied to investigate the toxic effect of  $As_2O_3$  on normal human breast tissue. When incubation with  $As_2O_3$  at IC<sub>50</sub> concentration on MDA-MB-231 cells at 48 and 72 h were applied, the percentages of survival of 184B5 cells were over 70% for both 48h and 72h  $As_2O_3$  treatment (data not shown). Our data on nude mice also showed that  $As_2O_3$  at dosage 0.06 mg/kg/day to 0.12 mg/kg/day administered on every other days for 14 days has no apparent toxicity to the animal (data not shown).

## CONCLUSION

In conclusion,  $As_2O_3$  was found to be effective in treating breast tumour cells including late stage breast tumour cells which are  $Er\alpha$  negative and administration

of As<sub>2</sub>O<sub>3</sub> was found to be safe to normal breast cells and on nude mice. Our results indicated that As<sub>2</sub>O<sub>3</sub> is a potential candidate to be further investigated as an anti-tumour drug for human breast tumours.

#### ACKNOWLEDGEMENT

This study was supported by a Direct Grant of Research Grant Committee, Hong Kong (Project code: 2041182) and a grant from South China National Research Centre for Integrated Biosciences in Collaborates with Zhongshan University (Project code: 1902006).

#### REFERENCES

- Chow, S.K., J.Y. Chan and K.P. Fung, 2004a. Inhibition of cell proliferation and the action mechanisms of arsenic trioxide (As<sub>2</sub>O<sub>3</sub>) on human breast cancer cells. *J. Cell. Biochem.*, 93: 172-187.
- Chow, S.K., J.Y. Chan and K.P. Fung, 2004b. Suppression of cell proliferation and regulation of estrogen receptor alpha signaling pathway by arsenic trioxide on human breast cancer MCF-7 cells. *J. Endocrinol.*, 182: 325-337.
- Clemons, M., S. Danson and A. Howell, 2002. Tamoxifen ('Nolvadex'): A review. *Tumour Treatment Rev.*, 28: 165-180.
- Di Leonardo, A., S.P. Linke, K. Clarkin and G.M. Wah, 1994. DNA damage triggers a prolonged p53-dependent G1 arrest and long-term induction of Cip1 in normal human fibroblasts. *Genes Dev.*, 8: 2540-2551.
- Hajek, R.A., A.D. Robertson, D.A. Johnson, N.T. Van, R.K. Tcholakian, L.A. Wagner, C.J. Conti, M.L. Meistrich, N. Contreras, C.L. Edwards and L.A. Jones, 1997. During development, 17 $\alpha$ -estradiol is a potent estrogen and carcinogen. *Environ. Health Perspec.*, 105: 577-581.
- Halloran, P.J. and R.G. Fenton, 1998. Irreversible G2-M Arrest and Cytoskeletal Reorganization Induced by Cytotoxic Nucleoside Analogues. *Tumour Res.*, 58: 3855-3865.
- Jordan, V.C., 1976. Effect of tamoxifen (ICI 46,474) on initiation and growth of DMBA-induced rat mammary carcinoma. *Eur. J. Tumour.*, 12: 419-424.
- Kinjo, K., M. Kizaki, A. Muto, Y. Fukuchi, A. Umezawa, K. Yamato, T. Nishihara, J. Hata, M. Ito, Y. Ueyama and Y. Ikeda, 2000. Arsenic trioxide (As<sub>2</sub>O<sub>3</sub>)-induced apoptosis and differentiation in retinoic acid-resistant acute promyelocytic leukemia model in hGM-CSF-producing transgenic SCID mice. *Leukemia*, 14: 431-438.
- Konig, A., L. Wrazel, R.P. Warrell, R. Rivi, P.P. Pandolfi, A. Jakubowski and J.L. Gabilove, 1997. Comparative activity of melarsoprol and arsenic trioxide in chronic B-cell leukemia lines. *Blood*, 90: 562-570.
- Li, X., X. Ding and T.E. Adrian, 2004. Arsenic trioxide caused redistribution of cell cycle, caspase activation and GADD expression in human colonic, breast and pancreatic tumour cells. *Tumour Invest.*, 22: 389-400.
- Li, Y.S., T.D. Zhang, C.H.W. Li, X.L. Zhao, Z.H.R. Wei, W. Tan, R.L. Li and Y.Y. Mao, 1988. Traditional Chinese and Western Medicine in the treatment of 27 patients with malignant lymphoma. *Chin. J. Oncol.*, 10: 61.
- Ling, Y.H., J.D. Jiang, J.F. Holland and R. Perez-solar, 2002. Arsenic trioxide produced polymerization of microtubules and mitotic arrest before apoptosis in human tumour cell lines. *Mol. Pharmacol.*, 62: 529-538.
- Lippman, M., G. Bolan and K. Huff, 1976. The effects of estrogens and anti-estrogens on hormone-responsive human breast tumour in long-term tissue culture. *Tumour Res.*, 36: 4595-4601.
- Miyashita, T. and J.C. Reed, 1995. Tumour suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell*, 80: 293-299.
- Osborne, C.K., 1998. Tamoxifen in the treatment of breast tumour. *N. Engl. J. Med.*, 339: 1609-1618.
- Park, W.H., Y.H. Cho, C.W. Jung, J.O. Park, K. Kim, Y.H. Im, M.H. Lee, W.K. Kang and K. Park, 2003. Arsenic trioxide inhibits the growth of A498 renal cell carcinoma cells via cell cycle arrest or apoptosis. *Biochem. Biophys. Res. Commun.*, 300: 230-235.
- Shao, W., M. Fanelli, F.F. Ferrara, R. Riccioni, A. Rosenauer, K. Davison, W.W. Lamph, S. Waxman, P.G. Pelicci, F.L. Coco, G. Avvisati, U. Testa, C. Peschle, C. Gambacorti-Passerini, C. Nervi and W.H. Miller, 1998. Arsenic trioxide as an inducer of apoptosis and loss of PML/RAR protein in acute promyelocytic leukemia cells. *J. Natl. Tumour Inst.*, 90: 124-133.
- Shen, Z.X., G.Q. Chen, J.H. Ni, X.S. Li, S.M. Xion, Q.Y. Qiu, J. Zhu, W. Tang, G.L. Sun, K.Q. Yang, Y. Chen, Z.W. Fang, Y.T. Wang, J. Ma, P. Zhang, T.D. Zhang, S.J. Chen and Z.Y. Wang, 1997. Use of arsenic trioxide (As<sub>2</sub>O<sub>3</sub>) in the treatment of acute promyelocytic leukemia (APL): II. clinical efficacy and pharmacokinetics in relapsed patients. *Blood*, 9: 3354-3360.

- Soignet, S.L., P. Maslak, Z.G. Wang, S. Jhanwar, E.L.J. Calleja Dardashti, D. Corso, A. DeBlasio, J. Gabrilove, D.A. Scheinberg, P.P. Pandolfi and R.P. Warrell, 1998. Complete remission after treatment of acute promyelocytic leukemia with arsenic trioxide. *N. Engl. J. Med.*, 339: 1341-1348.
- Soignet, S.L., S.R. Frankel, D. Douer, M.S. Tallman, H. Kantarjian, E. Calleja, R.M. Stone, M. Kalaycio, D.A. Scheinberg, P. Steinherz, E.L. Sievers, S. Coutre, S. Dahlberg, R. Ellison and R.P. Warrell, 2001. United States multicenter study of arsenic trioxide in relapsed acute promyelocytic leukemia. *J. Clin. Oncol.*, 19: 3852-3860.
- Sun, H.D., L. Ma, X.C. Hu and T.D. Zhang, 1992. Ai-Lin I treated 32 cases of acute promyelocytic leukemia. *Chin. J. Integrated Chin. West Med.*, 12: 170-171.
- Umans, R.S., R.R. Weichselbaum, C.M. Johnson and J.B. Little, 1984. Effects of estradiol concentration on levels of nuclear estrogen receptors in MCF-7 breast tumour cells. *J. Steroid. Biochem.*, 20: 605-609.
- Wu, X., Z. Chen, Z. Liu, H. Zhou, Y. You, W. Li and P. Zou, 2004. Arsenic trioxide inhibits proliferation in K562 cells by changing cell cycle and surviving expression. *J. Huazhong Univ. Sci. Technology Med. Sci.*, 24: 342-344, 353.
- Zhang, W., K. Ohnishi, K. Shigeno, S. Fujisawa, K. Naito, S. Nakamura, K. Takeshita, A. Takeshita and R. Ohno, 1998. The induction of apoptosis and cell cycle arrest by arsenic trioxide in lymphoid neoplasms. *Leukemia*, 12: 1383-1391.