

Parasitic Infections among Restaurant Workers in Tabriz (East-Azerbaijan Province) Iran

¹Y. Garedaghi and ²S. Safar Mashaei

¹Department of Pathobiology, Faculty of Veterinary Medicine,

²Tabriz Young Research Club, Islamic Azad University, Tabriz Branch, Tabriz, Iran

Abstract: To identify intestinal parasites among restaurant workers in Tabriz (East Azerbaijan province), Iran in 2009. Stool specimens were collected and examined from total of 300 restaurants workers at Veterinary Faculty, Islamic Azad University of Tabriz branch. Three types of techniques were used: direct examination, saline sedimentation and formol-ether concentration. The positivity in majority of them was single infection whereas 3 cases were double infection that constituted 1% of the prevalence. The prevalence was 14.4% for *Entamoeba histolytica*/dispor and 5.9% for *Giardia lamblia* while it was 4.8% for *Hymenolepis nana*. Other intestinal parasites including *Ascaris lumbricoides*, *Ancylostoma duodenale* were also detected. Additionally, the blood parasite *Schistosoma mansoni* was also detected in 2 cases. The double infection was only with *E. histolytica*/dispor and *Giardia*. The infection with these parasites were also accompanied by abdominal troubles diarrhea, constipation, nausea and vomiting. These results lead to understand that sanitary measurements are not effective and these hazardous situation facilitate parasitic agent's distribution among clients. The effectiveness of current pre-employment screening policy must be annual and systematic surveillance is needed in addition to health education.

Key words: Parasitic infections, restaurants workers, abdominal troubles, nausea, vomiting, Iran

INTRODUCTION

Survey on intestinal parasites among restaurant workers was done in all over the world (Fallah and Amirshakery, 2006; Ali *et al.*, 1992; Khan *et al.*, 1987). Shojaei *et al.* (2006) the effect of one simple hand washing of Iranian restaurant workers in decrease of food microbial infection by hands were shown (Shojaei *et al.*, 2006). The aim of this study is to determination of contamination rate to intestinal parasites among restaurant workers (foodhandlers) in Tabriz city (center of East Azerbaijan province) was done. This group of restaurant workers for direct contact with different classes of society have very importance role in distribution of parasitic agents.

MATERIALS AND METHODS

This survey is one cross-sectional study and from May 2009 to Sep 2010 in Tabriz city for determination of contamination rate to intestinal parasites among restaurants workers was done. In the beginning, A total of 300 restaurant workers were submitted to answer the survey questionnaire, one designed form consist of valid hygiene certificate, symptoms of parasitic disease,

education level, workers literacy, etc. were delivered to restaurants workers for completing. The restaurants workers were examined by one clinical doctor and then some fresh fecal sample were received of them. About 270 (90%) of restaurants workers cooperated. All fecal samples of restaurants workers were examined by 3 currency fecal examination method (direct examination, saline sedimentation and formol-ether concentration) in parasitology laboratory of Veterinary Faculty of Islamic Azad University Tabriz branch (Garcia and Bruckner, 1988).

RESULTS AND DISCUSSION

Table 1 shows the prevalence rate of intestinal parasites among restaurant workers in Tabriz city. In this

Table 1: Percentage and prevalence of intestinal parasites in stool examination of restaurants workers

Type of parasites	Parasites (%)	Number infected	Prevalence (%)
<i>Entamoeba histolytica</i> /dispor	50	39	4/14
<i>Giardia lamblia</i>	20/5	16	6
<i>Hymenolepis nana</i>	16/6	13	5
<i>Schistosoma mansoni</i>	2/5	2	8/0
<i>Ancylostoma duodenale</i>	6/4	5	8/1
<i>Ascaris lumbricoides</i>	3/8	3	1/1
Total	100	78	1/29

Corresponding Author: Y. Garedaghi, Department of Pathobiology, Faculty of Veterinary Medicine, Islamic Azad University, Tabriz Branch, Tabriz, Iran

Table 2: Restaurant workers complaints versus percentage and parasitic infections

Symptoms	Number	Percentage	<i>Hymenolepis nana</i>	<i>Giardia lamblia</i>	<i>Entamoeba histolytica</i> /dispor	Ascaris	Ancylostoma	<i>Schistosoma mansoni</i>
Abdominal pain	31	23/4	0	0	17	0	0	0
Constipation	9	6/8	6	0	0	1	1	1
diarrhea	49	37/1	0	14	35	0	0	0
Fever	37	28	0	7	24	2	2	1
Distention	20	15/1	0	13	7	0	0	0
Nausea and vomiting	41	31	4	7	25	2	2	1
No symptoms	12	9	2	0	5	2	1	1

Table 3: Education levels of restaurant workers

Education level	Number	Percentage
Illiterate	46	1/17
Primary school	162	60
Secondary school	62	9/22
Total	270	100

table also type of intestinal parasites and percent of them has been illustrated. *Entamoeba histolytica* with 50% show maximum infestation rate. In this survey only 2 persons of workers infected with intestinal *Schistosoma mansoni* were observed that this 2 person belong to other regions of Iran which this parasites in this region is current. Table 2 shows infected person with abdominal disorders and relation between main parasites with abdominal troubles in Tabriz city, i.e., *E. histolytica* and *Giardia*. Nausea and vomiting in all of these infections were observed. Table 3 shows education level of Tabriz restaurants workers. Parasitic infections have very importance in human life because this agent causes a dangerous disease in human. World Health Organization (WHO) has been reported which diseases causes by infected foods is one of largest hygienic problems in the world (Sheth and Dwivedi, 2006). Multilateral identification of parasites especially distribution agents of them (eggs and cysts) is secure way for determining of parasites percent and prevalence rate in societies. Restaurant workers are very susceptible population for distribution of parasites infections because these persons have direct contact with restaurant customers. Incidentally the infected person with intestinal parasites disease have very dangerous role for general hygiene of society especially if this person in shopping center is being done therefore this person are counted as reservoir and source for parasitic infections (Khan *et al.*, 1987).

Intestinal parasites have been reported in restaurant workers of very country in all over the world (Abahussain, 2005; Fallah and Amirshakery, 2006; Ibrahim *et al.*, 1993; Ali *et al.*, 1992; Kalantan *et al.*, 2001; Khan *et al.*, 1987; Nolla and Contos, 2005). In Brazil 44.9%, in Saudi Arabia 31.4% and in Yamane 28.7% has been reported. In Tabriz city intestinal parasites of restaurant workers 33% have been reported (Fallah and Amirshakery, 2006). Prevalence rate of intestinal parasites in restaurant workers of Tabriz city in comparison with previous

studies is low and in this study many of parasites in restaurant workers were not identified. *Entamoeba histolytica*/dispor and *Giardia lamblia* with maximum infestation rate in this study were reported and most clinical signs of these parasites are abdominal disorders and diarrhea.

Incidentally Nausea and vomiting in all of these infections was seeing (Garcia and Bruckner, 1988; Chatterjee, 2006). Presence of *Ancylostoma duodenale* ensures that soil transmitted helminthes infection are widely distributed in tropical and subtropical areas. Especially in poor hygienic populations. Presence of *Schistosoma mansoni* in workers of Tabriz city restaurants is more in workers belong to other regions of Iran, in fact impolitic migration in civic region causes easy transmission of human intestinal parasites in society and in coming for non-drug treatment of them, this persons as reservoir for intestinal parasites with great longevity in Tabriz city will be done (Jong, 2002). The majority of infected restaurant workers are symptomatic but they were not proclaiming their suffering for unknown personal reasons so they may be considered as infections carriers.

The infected restaurant workers with clinical signs or without clinical signs are risk factor for public Health of society. Contaminated food plays major role in the occurrence of diarrhea disease (Sheth and Dwivedi, 2006). In this survey all restaurant workers with diarrhea had been infecting to *Entamoeba histolytica*/dispor and *Giardia*, these two parasites with direct contact and customary manner transmission to customers of restaurants and with one simple hand washing by workers decreased from 72.7-32% (Shojaei *et al.*, 2006). In fact the aim of this study was determination of contamination rate to intestinal parasites among restaurant workers that they have all valid employment certificates (these persons had been working in restaurants). This will help the authorities to review their laws of employment. By attention to increase of food origin diseases, complete health of food has not been attention by hygienic managers and restaurant workers (Kafarstein and Abdussalam, 1999).

It is to be known that hygienic measurements concerning the restaurant workers as food handlers must be considered in the first category of measurement control and the periodical supplement of health performance must

be obligatory for each worker. Most restaurant workers are illiterate or have primary school education (Table 3) and without any attention to hygienic standards transmission the parasites agents to other persons and this problem will be causes increase in risk of hygiene and health in society. It is the responsibility of the ministry of Health to motivate all the control rules in restaurants, school canteens and all cites of human propagation in order to minimize the distribution of intestinal parasitic agents.

CONCLUSION

In this study, food safety education is a critical prerequisite and health education in general should be increased to raise awareness of the society about intestinal parasites problems (Abahussain, 2005; Kalantan *et al.*, 2001; Nolla and Contos, 2005; Sheth and Dwivedi, 2006) so researchers are in need for constant epidemiological surveillance through periodical surveys parallel with development of healthcare towards the problem of parasitic infections.

ACKNOWLEDGEMENT

This study was financially supported by a grant from Deputy Research of Islamic Azad University-Tabriz Branch, Tabriz, Iran. The researchers declare that they have no conflicts of interest.

REFERENCES

- Abahussain, N.A., 2005. Prevalence of intestinal parasites among expatriate workers in Al-Khobar, Saudi Arabia. *Middle East J. Family Med.*, 3: 17-21.
- Ali, S.I., K. Jamal and S.M.H. Qadri, 1992. Prevalence of intestinal parasites among food handlers in Al-Madinah. *Ann. Saudi Med.*, 12: 63-66.
- Chatterjee, K.D., 2006. *Parasitology (Protozoology and Helminthology) in Relation to Clinical Medicine*. 12th Edn., Chatterjee Medical Publishers, Calcutta, India.
- Fallah, E. and S.H. Amirshakery, 2006. Survey prevalence of intestinal parasites infections in food handlers in Tabriz. 9th Iranian Nutrition Congress, Tabriz University of Medical Sciences. (Persian).
- Garcia, L.S. and D.A. Bruckner, 1988. *Diagnostic Medical Parasitology*. Elsevier Science Publishing, New York.
- Ibrahim, O.M., A. Bener and A. Shalabi, 1993. Prevalence of intestinal parasites among expatriated workers in Al-Ain, United Arab Emirates. *Ann. Saudi Med.*, 13: 126-129.
- Jong, E., 2002. Intestinal parasites. *Primary Care*, 29: 857-877.
- Kaferstein, F. and M. Abdussalam, 1999. Food safety in the 21st century. *Bull. World Health Organ.*, 77: 347-351.
- Kalantan, K.A., E.A. Al-Faris and A.A. Al-Taweel, 2001. Pattern on intestinal parasitic infection among food handlers in Riyadh, Saudi Arabia. *Saudi Soc. Family Community Med.*, 8: 1-12.
- Khan, Z.A., A.A. Al-Jama and I. Madan, 1987. Parasitic infection among food handlers in Dammam and Al-Khbar, Saudi Arabia. *Ann. Saudi Med.*, 7: 47-50.
- Nolla, A.C. and G.A. Contos, 2005. Relationship between intestinal parasites in food handlers and epidemiological factors in the city of Florianopolis, Santa Catarina, Brazil. *Cad Saud Publica*, 21: 641-645.
- Sheth, M. and R. Dwivedi, 2006. Complementary foods associated diarrhea. *Indian J. Pediatr.*, 73: 61-64.
- Shojaei, H., J. Shooshtaripoor and M. Amiri, 2006. Efficacy of simple hand-washing in reduction of microbial hand contamination of Iranian food handlers. *Food Res. Int.*, 39: 525-529.