



Frequency of Detectability of Larynx Pathology among Patients with Chronic Diseases of the Gastrointestinal Tract

¹S. Taukeleva, ¹D. Toguzbaeva, ²S. Zharmenov, ³A. Mustafin and ²A. Iglíkova

¹Department of Otorhinolaryngology¹, Almaty State Institute of Advanced Medical, Almaty, Kazakhstan

²Kazakh Medical University of Continuing Education, Almaty, Kazakhstan

³Department of Otolaryngology and Eye Diseases, JSC Medical University Astana, Kazakhstan

Key words: Pachydermia, pathology of throat, gastrointestinal tract disorder, interrelation, video image endoscopy

Corresponding Author:

S. Taukeleva

Department of Otorhinolaryngology, Almaty State Institute of Advanced Medical, Almaty, Kazakhstan

Page No.: 123-125

Volume: 14, Issue 5, 2020

ISSN: 1815-9346

Research Journal of Medical Sciences

Copy Right: Medwell Publications

Abstract: The conducted clinical and instrumental study revealed that the aggregate group of patients with chronic diseases of the throat differs by increased frequency of both isolated and combined disorders of the gastrointestinal tract. If a patient has such symptoms as hoarseness, tickling, scratching, burning, or feelings of dryness in the throat, a frequent variant of gastrointestinal tract disorders is gastroesophageal reflux disease with pharyngolaryngeal reflux. There are such severe pathologies of the throat as pachydermia of the vocal cords and the nodules of vocal cords among more frequent disorders of gastrointestinal tract diseases. There is a high probability of accompanying pathology of the throat in the form of pachydermia and nodules of the vocal cords with patients with gastrointestinal tract disorders for conducting a fibrorhinolaryngoscopy in combination with a fibergastrososcopy.

INTRODUCTION

The long-term course of functional diseases of throat, especially for people in the voice and speech professions, leads to an organic pathology of the vocal apparatus. The mucous membrane of the throat, larynx and gastrointestinal tract represents a nonseparable whole therefore, in many cases laryngitis develops simultaneously or subsequent to chronic gastritis or cholecystopancreatitis. One reason such symptoms occur is pharyngolaryngeal reflux, leading to refluxate entering the area located above the upper esophageal sphincter^[1-3]. Lately, the manifestations of gastroesophageal reflux disease outside the esophagus, especially from part of the bronchopulmonary system and the ENT organs have attracted attention. Despite their frequent combination, the

cause-effect relations of gastrointestinal tract disorders with the pathology of the throat have not been studied sufficiently^[4-6].

The uncertainty of the problem related to adequate diagnostics of gastrointestinal tract disorders with patients with vocal dysfunction of the throat pathology gave rise to the goal and determined this work. Purpose: to study the interrelation between gastrointestinal tract disorders of patients with different vocal dysfunctions^[7-9].

Tasks of study:

- To study the interrelation between gastrointestinal tract disorders and throat diseases and vocal dysfunction
- To define the priority of throat diseases and vocal dysfunction diagnostic methods based on complex clinical and instrumental studies

- To conduct a comparative analysis of the frequency of occurrence of gastrointestinal tract disorders options depending on the pathology of the throat

MATERIALS AND METHODS

The clinical and instrumental examination of 99 patients (62 male and 37 female) aged between 50 and 60 years (mean±SD: 8.0±2.6 years) with chronic diseases of the throat with different disorders of gastrointestinal tract or combination of these nosologies was held. The gastroenterological group included 39 patients with disorders of the upper gastrointestinal tract according to physical, endoscopic and X-ray examinations. The criterion of inclusion into the otorhinolaryngologic group was the presence of chronic diseases of the throat in combination with vocal dysfunction. This group included 60 patients followed up by otolaryngologists with pachydermia of the vocal cords (PVC, n = 38) and “singing nodules” vocal cords nodules (VCN, n = 22)^[10].

Fibrogastrroduodenoscopy was used to evaluate the condition of the mucous membrane of the esophagus, stomach and duodenum as well as motor defects in the area of the esophagogastric junction. The radiographic contrast study of the esophagus made it possible to diagnose gastroesophageal refluxes of different degrees of manifestation^[11].

All patients were examined by means of a Karl Storz video endoscope using a camera with a 90° angle of vision. Analysis of vocal function was done using a speech analyzer computer program. The patient’s speech was recorded to study vocal function. The patients called name, surname and patronymic, year of birth and date of study, and then pronounced the vowel sounds, “A”, “I” and “Y” in a drawling manner. Analysis of the signal was held by one of the options in the software program, as this program is able to reveal the abnormalities in vocal range without specification of such acoustic parameters. This made it possible to evaluate the presence of deviations in vocal function condition and to reveal the necessity of more detailed analysis^[12].

RESULTS AND DISCUSSION

In the otorhinolaryngologic group an endoscopic study established the diagnosis of chronic pathology of the throat. In the course of the study, the endoscopic signs of pharyngolaryneal reflux were also defined: hyperemia and edema of the mucous membrane of arytenoid cartilages and interarytenoid area, hyperemia of back sections of vocal folds, inflammable rollers in the underfolds section and pachydermia. In the course of study, 93% (n = 92) of patients from both groups complained of hoarseness, tickling, irritation,

scratching, burning and a feeling of dryness in the throat. The terms of the disease were between 1 month and 20 years^[13, 14].

Statistical processing of data was done on a personal computer using Statistica 6.1 and Microsoft Excel 2003 programs according to medical statistics recommendations. Following the results of study, the following was revealed: an isolated disorder of upper sections of gastrointestinal tract, a combination of gastrointestinal tract disorders and pathologies of throat, and isolated disease of the throat. The patients with a combination of gastrointestinal tract disorders and throat pathologies (n = 56, 56.6%) were the majority of the examined patients (N = 99). The isolated diseases of throat and gastrointestinal tract were observed rarely (n = 14, 14.1%; и, n = 14, 14.1%, accordingly)^[15].

The analysis of the degree of throat pathology influence upon the detectability of gastrointestinal tract disorders was also of interest. The accumulative group of the patients with throat pathology included 60 patients followed up by otolaryngologists with the abovementioned ENT diseases and 40 patients with initial diagnosis of reflux disease.

Pachydermias (n = 33) and nodules of vocal folds (n = 7) in combination with dysphonia were diagnosed in the patients with an initial diagnosis of reflux disease. Pachydermia was located on the vocal folds in 21 (63.7%) patients, vestibular folds in 8 (24.2%) patients and in the interarytenoid area or near to arytenoid cartilages in 4 (12.1%) patients. The following was revealed by means of endoscopy: hyperemia, edema of mucous membrane of arytenoid cartilages and interarytenoid area, hyperemia of back sections of vocal folds, inflammable rollers in under folds section, and pachydermia (n = 33, 82.5%). Pachydermias had a warty structure; they were similar to outgrowths or patches. Coloring of the pachydermia varied from whitish-grey to bright yellow or pink. Thus, the frequency of revealing the specified pathological states of the throat in the patients with gastroenterological practice was 66.7%.

The symptomatic analysis of the examined patients established certain influences of reflux disease options and gastrointestinal tract disorders upon the clinical implications of pathology. There was regurgitation and vomiting (42.9 and 21.4%, respectively) with isolated disorders of the gastrointestinal tract. There was also vocal dysfunction among the symptoms outside the esophagus in the majority of patients with an isolated disease of throat or with the combined reflux disease (80.0% and 66.7%, respectively). The signs of end oesophagitis (77.8%) without evidence of frequency differences depending on the reflux disease option were revealed in the majority of patients who underwent fiber optic esophagogastroduodenoscopy. Reflux disease with esophagitis signs (64.8%) was the prevailing

pathology of the esophagus and without evident inflammatory changes of the mucous membrane (22.2%)^[16].

The conducted acoustic analysis of the voice (functionally limited under technical parameters) revealed vocal dysfunction in all patients but with different degrees of manifestation. The radiographic contrast study of one-third of patients did not reveal the signs of contrast entering the esophagus (33.3%). The same study poorly reflected the level of reflux. The gastroesophageal reflux of 3-4 degrees was insignificant more often with the combined option of reflux disease (26.9%) compared with gastroesophageal reflux disease (14.3%)^[17].

It is expedient to consider the revealed association of gastrointestinal tract disorder with diseases of the throat from the point of view of cause-and-effect relationships. In particular, disorders of the upper sections of the gastrointestinal tract are able to play a part in the provoking factor of throat pathologies and to consequently damage the mechanisms of the clearance of the mucous membrane of respiratory tract upper divisions^[18].

The revealed patterns are likely to have an interdependent nature and may reflect different pathogenetic mechanisms of ENT pathology or degrees of its influence upon gastrointestinal tract disorders.

CONCLUSION

The frequent option of gastrointestinal tract disorders is gastroesophageal reflux with pharyngolaryngeal reflux. The accumulative group of the patients with chronic diseases of the throat differs by increased frequency of isolated and combined disorders of the gastrointestinal tract. Presence of severe pathology of throat, for example, pachydermia of vocal folds and nodules of vocal folds, is the most often pathology in gastrointestinal tract diseases.

A high probability of an accompanying pathology of the throat in the form of pachydermia and nodules of vocal folds in patients with gastrointestinal tract disorders could be due to the prevailing culture in Central Asia as fatty and fried spicy food is the main part of the diet of the local population of patients.

REFERENCES

01. Agadzhanian, N.A., U.S. Vasilenko and A.I. Smirnova, 2005. [The effect of phonational load on parameters of cardiorespiratory system in hypotonic dysphonia]. Bull. Otorhinolaryngology, 4: 15-17.
02. Agaphonova, T.N., 2002. Psychoemotional status at patients with functional impairment of voice. Russ. Otorhinolaryngology, 1: 7-8.
03. Arnold, G.E. and L.W. De Santo, 1983. Adductor spastic dysphonia: Three years after recurrent laryngeal nerve resection. Laryngoscope, 93: 1-8.
04. Bohme, G., 1980. Funktionelle Dysphonie. Ther. Umsch., 37: 1043-1046.
05. Carding, P.N., I.A. Horsley and G.J. Docherty, 1998. The effectiveness of voice therapy for patients with non-organic dysphonia. Clin. Otolaryngology Allied Sci., 23: 310-318.
06. Fant, G., K. Ishizaka, J. Lindqvist and Sundberg, 1972. Speech analysis and speech production. Royal Transmission Laboratory, Quarterly Progress and Status Report, KTH Royal Institute of Technology, Stockholm, Sweden.
07. Husson, R., 1955. [Vocal cord vibration physiology (In French)]. Acad. Des. Sci., 2: 242-244.
08. Isshiki, N., H. Okamura and M. Morimoto, 1967. Maximum phonation time and air flow rate during phonation: Simple clinical tests for vocal function. Ann. Otolaryngology Laryngology, 76: 998-1007.
09. Koufman, J.A. and P.D. Blalock, 1982. Classification and approach to patients with functional voice disorders. Ann. Otolaryngology Laryngology, 91: 372-377.
10. Lavin, N., P.A. Singer and J.M. Hershman, 1994. Manual of Endocrinology and Metabolism. Lippincott Williams & Wilkins, New York, USA., pp: 519-657.
11. Luchsinger, R., 1948. [For stroboscopic symptoms (In German)]. Pract. Oto-Rhino-Laryngologica, 10: 209-214.
12. Monday, L.A., 1983. Clinical evaluation of functional dysphonia. J. Otolaryngology, 5: 307-310.
13. Morrison, M.D., H. Nichol and L.A. Rammage, 1986. Diagnostic criteria in functional dysphonia. Laryngoscope, 96: 1-8.
14. Newman, D. and K. Forbes, 1993. The effects of danazol on vocal parameters is an objective prospective study needed?. Med. J. Aust., Vol. 158, No. 8.
15. Sopko, J., 1983. [Larynx changes due to functional dysphonia (In German)]. ENT. Ear Nose Throat Specialists, 31: 335-342.
16. Stoicheff, M.L., 1983. The present status of adductor spastic dysphonia. J. Otolaryngology, 12: 311-314.
17. Vassilenko, U.S. and G.F. Ivanchenko, 1977. Functional status of throat from the point of view of age. Ear Throat Nose, 23: 75-79.
18. Zraitskiy, L.A., V.A. Tripos and L.A. Tripos, 1984. Practical Phoniatics. Meridian International School Kiev Ukraine, Kiev, Ukraine, Page: 186.