

## Evaluation of *in vivo* Antiplasmodial Activity of Ethanolic Leaf Extract of *Lasianthera Africana*

<sup>1</sup>E. Okokon Jude, <sup>2</sup>S. Antia Bassey, <sup>3</sup>A. Essiet Grace and <sup>4</sup>Lucky L. Nwidu

<sup>1</sup>Department of Pharmacology and Toxicology, Faculty of Pharmacy, University of Uyo, Uyo, Nigeria

<sup>2</sup>Department of Chemistry, University of Uyo, Uyo, Nigeria

<sup>3</sup>Department of Pharmacology and Toxicology, College of Medical Sciences  
University of Calabar, Calabar, Nigeria

<sup>4</sup>Department of Pharmacology and Toxicology, Faculty of Pharmacy, Niger Delta University  
Wilberforce Island, Bayelsa State, Nigeria

**Abstract:** The *in vivo* antiplasmodial activity of the ethanol leaf extract of *Lasianthera africana* grown particularly for the leaf in Niger Delta region of Nigeria was evaluated in *Plasmodium berghei* infected mice. *Lasianthera africana* (1000-3000 mg.kg.day<sup>-1</sup>) exhibited significant (p<0.05) blood schizonticidal activity both in 4- day early infection test and in established infection with a considerable mean survival time though not comparable to that of the standard drug, chloroquine, 5 mg.kg.day<sup>-1</sup>. The leaf extract possesses significant (p<0.05) antiplasmodial activity, which can be exploited in malarial chemotherapy.

**Key words:** Antiplasmodial activity, ethanol leaf extract, *Lasianthera africana*

### INTRODUCTION

From time immemorial plants have served as food and medicine to man. Vegetables and leaves of some shrubs, domesticated or wild, are used by the Ibibios of Niger Delta region of southern Nigeria in the preparation of their soup daily. Some of these edible plants are equally medicinal and are used in the therapy of some diseases, majority of which have been reported to contain vital chemical compounds of medicinal importance.

*Lasianthera africana* (P. Beauv.) is a perennial glabrous shrub of the family Icacinaceae whose height may reach from 61 to 136 cm and is widely distributed in the tropical rain forest (Hutchinson and Dalziel, 1973). There are four ethnovarieties distinguished by their taste, leaf colour and ecological distribution. The leaves are consumed as vegetable in southern Nigeria. Ethnobotanically, *L. africana* is used as antacid, analgesic, antispasmodic, laxative, antipyretic, antiulcerogenic, antidiabetic and antimalarial. *L. africana* has been reported to be bacteriostatic (Itah, 1997) fungicidal (Itah, 1996) and antidiabetic (Ekanem, 2006). The aim of the present study, was to evaluate the nutraceutical potential of the dark green variety against *Plasmodium berghei* infection in mice.

### MATERIALS AND METHODS

**Plant materials:** Fresh leaves of *Lasianthera africana* were collected in August, 2006 from a garden in Uruan,

Akwa Ibom State, Nigeria. The plant was identified and authenticated by Dr. Margaret Bassey, a taxonomist in the Department of Botany, University of Uyo, Uyo, Nigeria. Herbarium specimen was deposited at Faculty of Pharmacy Herbarium. The fresh leaves (2kg) of the plant were dried on laboratory table for 2 weeks and reduced to powder. The powder 100g was macerated in 95% ethanol (300 mL) for 72 h. The liquid filtrate obtained was concentrated in vacuo at 40°C. The yield was 0.98% w/w. The extract was stored in a refrigerator at 4°C until used for experiment reported in this study.

**Phytochemical screening:** Phytochemical screening of the extract was carried out employing standard procedures (Harbone, 1993; Trease and Evans, 1989).

**Animals:** Albino Swiss mice (21-28g) of either sex were obtained from the University of Uyo animal house. They were maintained on standard animal pellets and water ad libitum. Permission and approval for animal studies were obtained from the College of Health Sciences Animal Ethics committee, University of Uyo.

**Parasite inoculation:** The chloroquine-sensitive *Plasmodium berghei berghei* was obtained from National Institute of Medical Research, Lagos, Nigeria and maintained in mice. The inoculum consisted of 5x10<sup>7</sup> *P. berghei berghei* parasitized erythrocytes per mL. This was prepared by determining both the

percentage parasitaemia and the erythrocytes count of the donor mouse and diluting the blood with isotonic saline in proportions indicated by both determinations. Each mouse was inoculated on day 0, intraperitoneally, with 0.2 mL of infected blood containing about  $1 \times 10^7$  *P. berghei berghei* parasitized red blood cells.

**Determination of LD<sub>50</sub>:** The LD<sub>50</sub> of the extract was determined using albino mice by Intraperitoneal (I.P.) route using the method of Lorke (1983).

**Evaluation of schizontocidal activity on early infection (4-day test):** Schizontocidal activity of the extract was evaluated using the method described by Knight and Peters (1980). Each mouse was inoculated on the first day (day 0), intraperitoneally, with 0.2 mL of infected blood containing about  $1 \times 10^7$  *P. berghei berghei* parasitized erythrocytes. The animals were divided into five groups of five mice each and orally administered, shortly after inoculation with 1000, 2000 and 3000 mg.kg.day<sup>-1</sup> doses of the *Lasianthera africana* extract, chloroquine 5 mg.kg.day<sup>-1</sup> and an equivalent volume of distilled water (negative control) for four consecutive days, (day 0 to day 3). On the fifth day (day 4), thin films were made from the tail blood of each mouse and the parasitaemia level was determined by counting the number of parasitised erythrocytes out of 200 erythrocytes in random fields of the microscope. Average percentage chemosuppression was calculated as

$$100 \left[ \frac{(A-B)}{A} \right]$$

Where A is the average percentage parasitaemia in the negative control group and B, average percentage parasitaemia in the test group.

**Evaluation of schizontocidal activity established infection (Curative or Rane test):** Evaluation of curative potential of the extract was done using a method similar to that described by Ryley and Peters (1970). The mice were injected intraperitoneally with standard inoculum of  $1 \times 10^7$  *P. berghei berghei* infected erythrocytes on the first day (day 0). Seventy-two hours later, the mice were divided into five groups of five mice each. The groups were orally administered with *Lasianthera africana* leaf extract (1000, 2000, 3000 mg.kg.day<sup>-1</sup>), chloroquine (5 mg.kg.day<sup>-1</sup>) was given to the positive control group and an equal volume of distilled water to the negative control group. The drug/extract was given once daily for 5 days. Thin films stained with Giemsa stain were prepared from tail blood of each mouse daily for 5 days to monitor the parasitaemia level. The mean survival time for each group was determined arithmetically by finding the average survival time (days) of the mice (post inoculation) in each group over a period of 28 days (day 0 to day 27).

**Statistical analysis:** Data obtained from the study were analyzed statistically using Student's test and values of  $p < 0.05$  were considered significant.

## RESULTS

**Acute toxicity:** The mice were treated intraperitoneally with a single dose of 1.5 g.kg<sup>-1</sup> of either *Lasianthera africana* leaf extract after being starved for 24h. The route was chosen because of its sensitivity and rapid results. *Lasianthera africana* (1.5 g.kg<sup>-1</sup>) produced no physical signs of toxicity in the animals 24 h after administration except writhing within the first minute of administration.

**Phytochemical screening:** Phytochemical screening of the ethanolic leaf extract of *Lasianthera africana* revealed the presence of compounds like alkaloids, terpenes, flavonoids, anthraquinones, saponins, cardiac glycosides and phlobatannins.

**4-day test:** Ethanolic leaf extract of *Lasianthera africana* produced a dose dependent chemosuppressive effect at various doses employed in this study. The chemosuppression were 41.15, 51.14 and 60.05 % for 1000, 2000 and 3000 mg.kg.day<sup>-1</sup> doses. The chemosuppression produced by the extract were significant ( $p < 0.05$ ) compared to control and uncomparable to that of the standard drug (chloroquine 5 mg.kg.day<sup>-1</sup>) with a chemosuppression of 88.2% (Table 1).

**Curative test:** On established infection, it was observed that there was a daily increase in parasitaemia of the control group. However, there was a daily reduction in the parasitaemia levels of the extract treated group as well as that of positive control (chloroquine).

On day 7, the average percentage parasitaemia for the groups were 30, 29.3, 26, 7.0 and 81 % for 1000, 2000, 3000 mg.kg.day<sup>-1</sup> of the extract, chloroquine and control groups respectively (Fig. 1). The mean survival time (m. s. t) of the extract treated groups were significantly ( $p < 0.05$ ) longer than that of control and was uncomparable to that of the standard drug, chloroquine. The values are given in Table 2.

## DISCUSSION

*Lasianthera africana*, a vegetable used by the Ibibios of the Niger Delta Region of Nigeria, was evaluated for its antiplasmodial potentials in *Plasmodium berghei* infected mice. Phytochemical screening and acute toxicity test of the ethanolic leaf extract were carried out.

Table 1: Antiplasmodial activity of *Lasianthera africana* leaf extract during 4-day test

Drug/Extract	Dose (mg.kg.day <sup>-1</sup> )	Average (%) parasitaemia	Average (%) suppression
<i>Lasianthera africana</i> extract	1000	25.66±0.94*	41.15
	2000	21.33±1.24*	51.14
	3000	17.33±1.25*	60.05
Chloroquine (standard)	5	5.13±0.38*	88.2
Distilled water (control)	0.2 mL	43.6±3.16	-

Data are expressed as mean ± S. D for five animals per group. p<0.05 when compared to control

Table 2: Mean survival time of mice receiving various doses of ethanolic leaf extract of *Lasianthera africana*

Drug/Extract	Dose (mg.kg.day <sup>-1</sup> )	Mean survival time (day)
<i>Lasianthera africana</i> extract	1000	12.3±1.53*
	2000	14.7±3.84*
	3000	18.4±0.95*
Chloroquine (standard)	5	30.0±0.00*
Distilled water (control)	0.2 mL	9.51±2.04

Data are expressed as mean ± S. D for five animals per group. p<0.05 when compared to control




Fig. 1: Effect of *Lasianthera africana* leaf extract on established infection (curative test)

The leaf extract which had negligible acute toxicity contains alkaloids, terpenes, flavonoids, tannins, saponins and cardiac glycosides. Thus revealing an enormous medicinal value of this leaves. The leaf extract demonstrated a significant ( $p<0.05$ ) antiplasmodial activity in both early and established infections, which can be attributable to the phytochemical components of the extract like alkaloids, flavonoids and terpenes which already had been implicated in antiplasmodial activity of various plants (Philipson and Wright, 1991; Christensen and Kharazmi, 2001). The antiplasmodial activity of the leaf extract was uncomparable to that of the standard drug, chloroquine, due to the crude nature of the extract. Despite this moderate antiplasmodial activity of the crude leaf extract, diets have been considered very important because plants consumed as food are ingested in relatively large amount and more regularly than that of same plant use in rituals and cosmetics (Etkin and Ross, 1991; Etkin, 1994). Accumulation of these medicinal constituents of these leaves in the body may culminate in a higher antiplasmodial activity. Moreso,

plants compounds which merely slow down or temporary arrest the growth of the parasite (plasmodistatic) as well as those which act as immune stimulant or helping to alleviate symptoms and reverse some pathological result of malaria infection are reported to potentiate malaria resistance and antiplasmodial activity in immune individuals living in endemic areas (Kirby, 1997). In this study, considerable antiplasmodial activity of the leaf extract was observed. Consumption of these leaves in medicinal and dietary context may have cumulative effect and may be responsible for the reported low parasitaemia and rates of malaria in adult malaria patients in some areas of the Niger Delta region of Nigeria (Ezedinachi *et al.*, 1992; Okokon and Ezedinachi, 2002). Etkin (1997) had observed a similar case in a malaria endemic community in Northern Nigeria.

## CONCLUSION

The results of this study demonstrated that the leaves of *Lasianthera africana* possess considerable antiplasmodial activity. Their consumption in diets can promote malaria resistance. Therefore, it would be interesting if the active principle is isolated, identified and characterised.

## ACKNOWLEDGEMENT

The authors are grateful to Mr. Nsikan Malachy for his technical assistance.

## REFERENCES

- Bassey, M.E. and J.U. Ekpo, 2004. Ethnobotany of some lesser known wetlands plants from Akwa Ibom State, Nigeria. *Liv. Sys. Sus.*, pp: 1-4.
- Christensen, S.B. and A. Kharazmi, 2001. Antimalarial natural products. Isolation, characterization and biological properties. In: *Bioactive compounds from natural sources: Isolation, characterization and biological properties.* (Ed.) Tringali, C. London, Taylor and Francis, pp: 379-432.
- Ekanem, A., 2006. Antidiabetic activity of ethanolic leaf extract and fractions of *Lasianthera africana* on alloxan diabetic rats. M. Sc. Thesis. University of Uyo, Nigeria.

- Etkin, N.L. and P.J. Ross, 1991. Should we set a place for diet in ethnopharmacology? J. Ethnopharmacol., 32: 25-36.
- Etkin, N.L., 1997. Antimalarial plants used by Hausa in Northern Nigeria. Tropical Doctor, 27: 12-16.
- Etkin, N.L., 1994. Consuming a therapeutic landscape: A muticontextual framework for assessing the health significance of human-plant interactions. J. Home Consumer Hort., 1: 61-81.
- Ezedinachi, E.N.U., A.A. Alaribe, M. Meremikwu and G.C. Ejezie, 1992. New Trend in chloroquine efficacy in the treatment of malaria: Significance of low (scanty) parasitaemia in an endemic area with emerging chloroquine-resistant *P. falciparum*. Central African J. Med., 38: 303-306.
- Harbone, J. B., 1983. Phytochemical Methods. A guide to Modern Techniques of plant Analysis. London, Chapman and Hall.
- Homburger, F., 1989. *In vivo* testing in the study of toxicity and safety evaluation. In: A Guide to General Toxicology. Marquis J.K. (Ed). (2nd Edn.), Karger, New York.
- Hutchinson, J. and J.M. Dalziel, 1973. Flora of west tropical Africa. (2nd Edn.), Crown Agents for Overseas Government. Administration, 1: 638.
- Itah, A.Y., 1997. Bactericidal and bacteriostatic effect of edible leafy vegetable extract on growth of canned food borne bacteria. Trans. Nig. Soc. Bio. Conserv., 6: 103-111.
- Itah, A.Y., 1996. Screening of plant parts for fungicidal properties. Trans. Nig. Soc. Bio. Conserv., 4: 26-40.
- Iwu, M.M., 1983. Report of a sponsored project by the institute of African studies. University of Nigeria, Nsukka.
- Knight, D.J. and W. Peters, 1980. The antimalarial action of N-benzyloxy dihydrotriazines. The action of cycloguanil (BRL50216) against rodent malaria and studies on its mode of action. Ann. Trop. Med. Parasitol., 74: 393-404.
- Kirby, G.C., 1997. Plants as a source of antimalarial drugs. Tropical Doctor, 27: 7-11.
- Lorke, D., 1983. A new approach to practical acute toxicity test. Arch. Toxicol., 54: 275-286.
- Okokon, J.E. and E.N.U. Ezedinachi, 2002. Plasma total chloroquine level in relation to *Plasmodium falciparum* density in adult malaria patients in Calabar, Nigeria. Global J. Med. Sci., 1: 41-47.
- Philipson, J.D. and C.W. Wright, 1991. Antiprotozoal compounds from plants sources. Planta Medica, 57: 553-559.
- Ryley, J.F. and W. Peters, 1970. The antimalarial activity of some quinone esters. Ann. Trop. Med. Parasitol., 84: 209-222.
- Trease, A. and W.C. Evans, 1989. Trease and Evans Pharmacology. (13<sup>th</sup> Edn.), London. BailliereTindal.