

Withdrawal Times of Intramammary Antibiotics in Camel Milk

O.M. Elemam, S.A. Omer, I.F. Albokhadaim and A.M. Homeida
Camel Research Centre, King Faisal University, P.O. Box 1757, Al-Ahsa, Saudi Arabia

Abstract: The study was conducted to determine withdrawal periods of intramammary preparations Curaclax (ampicillin+ceplulospirin), oxymast (oxytetracycline) and spectrazol (cefroxime) in camel and cow milk. Withdrawal periods in milk were 4 days in cows versus 5 days in camels for curaxlox and 4 days in cows versus 6 days in camel for oxmast and spectrazol. The significant ($p<0.05$) difference in withdrawal periods between camel and cow emphasizes the need to establish specific withdrawal periods for other drugs in the camel.

Key words: Curaclax, ampicillin, ceplulospirin, oxymast, spectrazol, periods of intamammary

INTRODUCTION

The antibacterial treatment of mastitis in lactating animals is of considerable regulatory concern because of the possibilities of antibacterial residues in milk (Sandholm *et al.*, 2009). Intramammary (imm) infusion has been recognized as the route for treating mastitis (Soback, 1988). Over 26% of milk samples collected 96 h post-treatment was positive for antibiotic residues (Berruga *et al.*, 2003).

The possible presence of antibacterial residues in milk poses a risk for consumers because these may cause allergic reactions, intestinal dysbiosis or even the emergence of resistant bacteria (Dewdney *et al.*, 1991). Failure to observe the correct withdrawal time was cited as the most common reason for violative drug residue levels in a study performed by the FDA in the 1970s (Bevill, 1989) and continued to be the most common cause of the residue violations in the 1990s.

The withdrawal period may be defined as the period that us required following the last medication in order to bring the concentration of the drug to below a tolerable value.

The actual withdrawal time appearing on antibiotic label is also a function of experimental design that the manufacture uses in the research studies. Such labels contain no information about withdrawal times in camels and veterinarians have to extrapolate withdrawal times of drugs for dairy camels from other species.

This is scientifically unacceptable since, the camel differs in its physiological, anatomical and pharmacological characteristics from other animals (Abdalla and Abdalla, 1979; Al-Dughaym *et al.*, 1998).

This study was carried out to determine the withdrawal period of intramammary antibiotics in camel milk.

MATERIALS AND METHODS

Animals and treatment: About 6 majahem lactating dromedary camels and 6 jersey dairy cows were used in this study, ranging in age from 4-6 years. The animals were housed in stalls and given free choice alfalfa hay and water. The entire content of the intramammary infusion syringe containing antibiotics, formulated for treatment of bovine mastitis was injected into each udder. The preparations used were oxymast (oxytetracycline, 500 mg, Bimeda, UK), curaxlox (amicillim 75 mg + cloxacillin 200 mg, Norbrook, UK) and spectrazol (cefuroxime 250 mg, Schering-Plough, Germany).

Collection of milk samples: Milk sampling was performed at 12 h interval. All samples were frozen immediately after they were taken. The samples were subsequently thawed and diluted 50:50 with distilled water and mixed well to reach the correct viscosity.

Antibacterial assay: Antibacterial concentration in milk was determined using an agar diffusion method with cut wells. *Bacillus stearothermophilus* (ATCC 10142) was used to test penicillins (Al-Nazawi and Homeida, 2005). *Bacillus subtilis* (ATCC 663) was used to test tetracyclines (Al-Nazawi and Homeisa, 2002).

Statistical test: ANOVA followed by Turkey's multiple range test was used to test the data. A $p<0.05$ was considered significant.

Antibacterial screening test: The screening tests used was brilliant black reduction test (BRT; AIM-An Lytik in Milch production-Und-Vertriebs UmbH; Munchen, Germany). Briefly, the test is based on detecting inhibition of the growth of spores of *Geobacillus stearothermophilus* var. calidolactis C. The test medium is a mixture of nutrients, test bacteria, a colored indicator (brilliant-black) and supplements.

When milk sample contains no residue or these are under the detection limits, the spores germinate and grow and their metabolic activity makes the indicator change color. In the presence of antibacterial residues, no bacterial activity will be detected and the indicator color will remain unchanged. The color change can be assessed visually or photometrically at wavelengths of 450 and 620 nm.

RESULTS AND DISCUSSION

Recently much attention has been paid to the utilization of the inherent properties of camel milk for human health as hyperglycemic, antidiabetic, anticholesterolaemic and anticancer (Mal *et al.*, 2000, 2001; Sahani *et al.*, 2005). Therefore, concern over camel milk antibiotic residues are economic as well as public health related.

Table 1 shows the results of depletion of intramammary antibiotics from camel udder. The Maximum Residue Limit (MRL) and safe clearance levels for antibiotic residues in milk fixed by Codex Alimentarius, European Countries (EC) regulations and United States food and drug administration for ampicillin, cloxacillin, cephalosporins and oxytetracycline are 4, 30, 100 and 100 mg kg⁻¹, respectively (Honkanen-Buzalski and Reybroeck, 1997). In the present experiment, the MRLs were reached after 132 h for ampicillin and cloxacillin, suggesting that these periods represent the withdrawal periods for the 4 antibiotics in camel milk.

The withdrawal periods of the 4 antibiotics are significantly (p<0.05) longer than those in goats (Karzis *et al.*, 2007), cows (Bangen *et al.*, 1992; Debackere, 1995) and sheep (Sierra *et al.*, 2009). Animal models of infection have shown that bacterial killing may be described as a function of either concentration dependent or time-dependent (Sharma *et al.*, 2002).

Bacteriostatic antibiotics and beta-lactams are time dependent. Furthermore, the camel exhibits certain characteristics that differ from other domestic animals. They have low glomerular filtration rate (Etzion and Yagil, 1985), long nephron (Abdalla and Abdalla, 1979) oval erythrocytes (Etzion and Yagil, 1985) and low drug metabolizing enzyme activities (El-Sheikh *et al.*, 1986, 1991; Al-Shiekh *et al.*, 1988) which are expected to delay the elimination of drugs in the camel.

Table 1: Effect of intramammary administration of curaclox, oxytetracycline and spectrazol on concentration of antibiotic in camel milk

Time (h) after antibacterial administration	Antibacterials			
	Curaclox (ampicillin)	Curaclox (cloxacillin)	Oxytetracycline	Spectrazol (cefruxime)
12	102.0±12.1	1225±65.0	980±44.0	8600±120.0
24	93.0±11.6	980±52.0	810±40.0	5250±110.0
26	81.0±9.10	820±44.0	720±30.0	3800±102.0
48	52.0±6.30	710±32.0	580±15.0	2100±90.0
60	36.8±5.10	605±25.0	480±14.0	1400±70.0
72	24.1±3.10	450±15.0	420±13.0	920±65.0
84	18.6±2.10	261±12.0	310±12.0	750±35.0
96	10.1±2.00	140±12.0	280±11.0	510±25.0
108	8.2±1.60	81±8.5	210±10.0	420±18.0
120	5.1±1.20	45±6.1	160±10.0	350±13.0
132	3.1±0.6*	26±3.1*	105±8.2	200±11.0
144	2.2±0.50	19±2.1	94±8.3*	95±10.0
156	1.6±0.20	12±2.0	70±7.1	120±9.5
168	1.2±0.20	9±1.5	53±5.1	90±7.6

* Maximum Residue Limits (MRL)

Table 2: Detection limits of brilliant reduction test in milk of camels and cows treated with intramammary antibiotics preparations

Time (h) after antibacterial administration	Antibacterials					
	curaclox (ampicillin+ cloxacillin)		Oxytetracycline		Spectrazol (cefruxime)	
	Camel	Cow	Camel	Cow	Camel	Cow
12	+	+	+	+	+	+
24	+	+	+	+	+	+
26	+	+	+	+	+	+
48	+	+	+	+	+	+
60	+	+	+	+	+	+
72	+	+	+	+	+	+
84	+	+	+	+	+	+
96	+	-	+	-	+	+
108	+	-	+	-	+	+
120	+	-	+	-	+	+
132	-	-	+	-	+	-
144	-	-	-	-	-	-
156	-	-	-	-	-	-
168	-	-	-	-	-	-

+ indicates presence of antibiotic and indicates absence of antibiotic in camel milk

The BRT results (Table 2) were negative at times and levels corresponding to MRL for the 4 antibiotics which further confirms the observation that withdrawal periods in the camel to be significantly longer than in cows. The values were 4 and 5 days for the camel for ampicillin and cloxacillin 4 days for the cow and 6 days for the camel for oxytetracycline and cefruxime.

CONCLUSION

In conclusion, BRT has been used to detect antibiotic residues in milk of cows, sheep and goats in many places (Sierra *et al.*, 2009). The difference in withdrawal periods in the camel from that in other ruminants emphasizes the need to establish newer withdrawal periods of drugs in camels.

ACKNOWLEDGEMENT

The researchers are thankful to the Deanship of Scientific Research, King Faisal University for financial support.

REFERENCES

- Abdalla, M.A. and O. Abdalla, 1979. Morphometric observations on the kidney of the camel (*Camelus dromedaries*). *J. Anat.*, 129: 45-50.
- Al-Dughaym, A.N., A.L. Al-Afalaq and A.M. Homeida, 1998. Review of adverse effects of some drugs in camels. *Rev. Elev. Med. Paystrop.*, 51: 81-86.
- Al-Nazawi, M.H. and A.M. Homeida, 2002. Disposition kinetics of oxytetracycline after intravenous injection in the Arabian camel. *J. Camel Pract. Res.*, 9: 5-8.
- Al-Nazawi, M.H. and A.M. Homeida, 2005. Residues of sulphadimidine and its metabolism N4-acetyl in camel milk. *Int. J. Pharmacol.*, 1: 249-251.
- Al-Shiekh, A., B.H. Ali, A.M. Homeida, T. Hassan and H.J. Hapke, 1988. The activities of amino pyrine N-demethylase, aniline 4-hydroxylase and UDP-glucouronyl transferease in tissue of camels, desert sheep and Nubian goats. *Gen. Pharmacol.*, 19: 713-717.
- Bangen, M., E. Skjerve, K. Grave and N.E. Soli, 1992. Prescribing of drugs for food-producing animals in Norway. Information about withdrawal times. *J. Vet. Pharmacol. Therapeutics*, 15: 180-187.
- Berruga, M.I., M. Yamaki, R.L. Althaus, M.P. Molina and A. Molina, 2003. Performances of antibiotic screening tests in determining the persistence of penicillin residues in ewe's milk. *J. Food Prot.*, 66: 2097-2102.
- Bevill. R.F., 1989. Sulfonamide residues in domestic animals. *J. Vet. Pharmacol. Ther.*, 12: 241-252.
- Debackere, M., 1995. Pharmacokinetics and pharmacodynamics of antimicrobials in relation to their residues in milk. Proceedings of the Symposium on Residues of Antimicrobial Drugs and other Inhibitors in Milk, Aug. 28-31, Kiel, Germany, pp: 41-53.
- Dewdney, J.M., L. Maes, J.P. Raynaud, F. Blanc and J.P. Scheid *et al.*, 1991. Risk assessment of antibiotic residues of beta-lactams and macrolides in food products with regard to their immunoallergic potential. *Food Chem. Toxicol.*, 29: 477-483.
- El-Sheikh, H.A., B.H. Ali, A.M. Homeida and T. Hassan, 1986. The cytochromes p-450 concentration in microsomes of liver, kidney and duodenal mucosa of the camel, sheep and goats. *Vet. Human Toxicol.*, 25: 527-529.
- El-Sheikh, H.A., B.H. Ali, A.M. Homeida, T. Hassan and H.J. Hapke, 1991. Activities of the glutathione-S-transferase and ethoxycoumarin-O-demethylase in tissue of camel, sheep, goats and rats. *Comp. Biochem. Physiol.*, 98: 293-297.
- Etzion, Z. and R. Yagil, 1985. Tritiated water metabolism during dehydration and rehydration in the camel. *J. Applied Physiol.*, 56: 217-220.
- Honkanen-Buzalski, T. and W. Reybroeck, 1997. Antimicrobials, Monograph on Residues and Contaminants in Milk and Milk Products. International Dairy Federation, Brussels.
- Karzis, J., E.F. Donkin and M. Petzer, 2007. The influence of Intramammary antibiotics treatment, presence of bacteria, stage of lactation and parity in dairy goats as measured by California milk cell test and somatic cell counts. *Ondersteproot J. Vet. Res.*, 74: 161-167.
- Mal, G., D.S. Sena, V.K. Jain and M.S. Sahani, 2001. Therapeutic utility of camel milk as nutritional supplement in chronic pulmonary tuberculosis. *Livest. Int.*, 7: 4-8.
- Mal, G., D.S. Sena, V.K. Jain, N.M. Singhvi and M.S. Sahani, 2000. Role of camel milk as an adjuvant nutritional supplement in human tuberculosis patients. *Livest. Int.*, 4: 7-14.
- Sahani, M.S., R.P. Agarwal, F.C. Tuteja, S.K. Ghouri, A.R. Singh and D.S. Sena, 2005. Hypoglycemic activity of camel milk in streptozotocin induced hyperglycemia in rats. *Indian J. Anim. Sci.*, 75: 1365-1367.
- Sandholm, M., L. Kaartinen and S. Pyorala, 2009. Bovine mastitis-why does antibiotics therapy not always work: An overview. *J. Vet. Pharmacol. Therapeutics*, 13: 248-260.