

Isolated Left Ventricular Noncompaction with Normal Systolic Function in a Middle Age Woman

¹Samiei, N. and ²M. Moshkani Farahani

Department of Echocardiography and General Cardiology, Shahid Rajaei Cardiovascular Medical center, Iran University of Medical Sciences, Tehran

Abstract: Isolated Left Ventricular Non-compaction (LVNC) is a rare congenital cardiomyopathy, which is characterized by excessively prominent trabecular meshwork and deep intratrabecular recesses. Common clinical presentations include: systolic and diastolic dysfunction, systemic embolism, ventricular arrhythmia. In this case we report a case of 42-year-old woman with history of embolic events and with the diagnosis of LVNC with preserved left ventricular systolic function.

Key words: Left ventricular noncompaction, systolic diastolic dysfunction

CASE REPORT

A 44 year-old woman was referred to our echo lab evaluating the source of retinal emboli. She was suffering from retinal emboli (ischemic events) from 3 years ago and transthoracic and transesophageal echocardiography was done for her for two times before this admission, but no positive finding was made. Her vital signs were stable and cardiac auscultation was normal. Other physical examination findings were unremarkable. Electrocardiography showed normal sinus rhythm, normal PR and QRS duration. Chest X ray was within normal. Two dimensional echocardiography showed normal left ventricular size with normal systolic function (LVEF=50-55%) with prominent trabeculations in the LV apex and posterolateral walls. Communication between these recesses and LV cavity was evident with color flow imaging and noncompacted/compacted layer ratio=2.1 (Fig 1).

Echocardiography findings were consistent with criteria of LVNC. Our patient had history of embolic events that the last one occurred 2 weeks before this admission.

DISCUSSION

Isolated ventricular noncompaction (LVNC) is a rare disorder of endomyocardial morphogenesis^[1,2]. It is characterized by prominent trabeculations with deep intratrabecular recesses and impaired ventricular systolic function in the absence of associated congenital heart disease^[3]. The disorder is believed to occur because of arrest of compaction of the loose interwoven mesh of myocardial fibers normally seen during development.



Fig. 1: Multiple trabeculation and deep recesses

Genetic mutations have been identified as a possible cause of LVNC^[4]. Most patients with LVNC present with congestive heart failure and the diagnosis is made by echocardiography. For clear cut echocardiographic criteria: 1) Absence of coexisting cardiac abnormalities 2) A two layered structure of the left ventricular wall with the end systolic ratio of noncompacted to compacted layer >2 3) finding this structure in the apical and mid ventricular areas 4) Blood flow directly from the ventricular cavity into deep intertrabecular recess^[5].

Magnetic Resonance Imaging (MRI) may also be helpful and in Alhabshan study MRI provided better delineation of the extent of the abnormal trabeculation in patients with noncompaction of the left ventricular myocardium. It was particularly useful when the myocardial involvement was subtle^[6]. There are few

reports of isolated left ventricular noncompaction with preserved systolic function^[7]. Our patient presented with history of recurrent embolic events with no positive clinical finding in two previous echocardiographies that was done for her. But the last echocardiography showed LVNC with preserved systolic function. LVNC is associated with a poor prognosis and most deaths are due to ventricular arrhythmia and heart failure, this patient was in middle age with preserved systolic function and no history of arrhythmia. Here are some questions that should be answered that whether preserved systolic function in LVNC can dislodge small thromboses that are made in deep recesses more than patients with depressed LV function and these patients are much more prone to embolic events than arrhythmia and heart failure. There are no specific modalities of treatment for LVNC, but systemic anticoagulation should be administered routinely for these patients.

REFERENCES

1. Oechslin, E.N., C.H. Attenhofer Jost, J.R. Rojas, P.A. Kaufmann and R. Jenni, 2000. Long term follow up of 34 adults with isolated left ventricular noncompaction: A distinct cardiomyopathy with a poor prognosis. *J. Am. Coll Cardiol.*, 35: 493-500.
2. Pignatelli R.H., C.J. McMahan, W.J. Dreyer, S.W. Danfield, J. Price and J.W. Belmont *et al.*, 2003. Clinical characterization of left ventricular noncompaction in children, A Relatively Common form of Cardiomyopathy. *Circulation*, 108: 2672-2678.
3. Chin, T.K., L.J. Perloff, R.G. Williams, K. Jue and R. Mohrmann, 1990. Isolated noncompaction of left ventricular myocardium: A Study of Eight Cases. *Circulation*, pp: 82-507.
4. Mc Mahan, C.J., A.C. Chang, R.H. Pignatelli, W.C. Miller-Hance, B.K. Eble, J.A. Towbin and S.W. Danfield, 2004. Left ventricular noncompaction cardiomyopathy in association with trisomy . *Pediatr Cardiol.*, pp: 18.
5. Jenni, R., E. Oechslin and J. Schneider, *et al.*, 2001. Echocardiographic and pathoanatomical characteristics of isolated left ventricular noncompaction: A Step Towards Classification as a Distinct Cardiomyopathy. *Heart*, 86: 666-671
6. Ichida, F., Y. Hamamichi and T. Miyawaki, *et al.*, 1999. Clinical features of isolated noncompaction of the ventricular myocardium. *J. Am. Coll. Cardiol.*, 34: 233-240.
7. Aras, D., O. Tufekcioglu, S. Topaloglu and K. Ergun, 2005. Preserved systolic function with isolated left ventricular noncompaction in an elderly patient. *Eur. J. Echocardiography*, pp: 9-30.