

# aJava

Asian Journal of Animal and Veterinary Advances



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)

## Plant as a Source of Natural Antiviral Agents

<sup>1</sup>Muhammad Nouman Sohail, <sup>2</sup>Fiaz Rasul, <sup>1</sup>Asia Karim, <sup>1</sup>Uzma Kanwal and <sup>3</sup>Idress Hamad Attitala

<sup>1</sup>Asian Network for Scientific Information, Faisalabad, Pakistan

<sup>2</sup>Faisalabad Institute of Research, Science and Technology, University PARK, Near Abbas Pur, Faisalabad, Pakistan

<sup>3</sup>Department of Botany, Faculty of Science, Omar Al-Mukhtar University, El-Bayda, Libya

*Corresponding Author: Muhammad Nouman Sohail, Asian Network for Scientific Information, Faisalabad, Pakistan*

### ABSTRACT

Viruses are one of the main hazards for both humans and animals. They enter in the living body and redirect body's metabolism to produce large copies of their genome and proteins. Diseases caused by these viruses are difficult to tackle with the help of currently available antiviral drugs. So the aim of this study was to explore the plants with reported antiviral activity, to get understanding for better control of these viruses. Herpes virus, Human Immunodeficiency Virus (HIV), influenza and hepatitis virus were at top among all studied viruses. Prominent modes of action against these viruses were inhibition of viral entry and its replication in host cell. Against RNA viruses plants mainly targeted their Reverse Transcriptase (RT) enzyme (like HIV) or protease (mostly found against hepatitis C virus). A range of active compounds have been identified which could be the potential antiviral agents for future drug development. Some plants like *Allium sativum*, *Daucus maritimus*, *Helichrysum aureonitens*, *Pterocaulon sphacelatum* and *Quillaja saponaria* emerged to have broad spectrum antiviral activity. Detail study of their phytochemicals and mode of action against these viruses could be help full for more effective control of hazardous viruses.

**Key words:** Antiviral, herpes virus, influenza, hepatitis, *Allium sativum*, *Dauucus maritimus*

### INTRODUCTION

Virus "a piece of bad news wrapped in a protein coat" has been defined by Peter Medawar (Oldstone, 1993). It appears as the perfect definition after considering the list of top ten causes of death in low, middle and high income countries. Lower respiratory infections, diarrhoeal diseases and HIV/AIDS are the common death causes among low and middle income countries (WHO, 2011b). All of these three health disorders are directly or indirectly caused by viruses. Except lower respiratory infections none of the above mentioned factors are prevalent among high income countries. It clearly indicates that how severely these viral diseases are affecting the people health in low and middle income/developing countries.

Our planet contains nearly  $10^{31}$  viruses and their ubiquity also invaded the marine environment, where in every 200 liter of water nearly 5000 viral genotypes are present (Breitbart and Rohwer, 2005; Suttle, 2005). Moreover viruses are moving between the environments and they are present almost everywhere e.g. deep sea, polar ice, alkaline, hot and saline waters and more than 2000 m deep in terrestrial environment. There are almost 20 families of viruses that actually

infect humans (Harvey *et al.*, 2006) and some of them also cause diseases in animals (Mahzounieh *et al.*, 2006). The diseases they cause in human include chickenpox, influenza, skin rash, hepatitis, bronchiolitis, acquired immunodeficiency syndrome, liver infection and many others. Virus particles enter in the living system and if they overwhelm the body's immune system then it is almost impossible to stop their spread in body. They direct the host metabolic pathway for the sake of their repeated replication; this makes their treatment difficult. But fortunately, it is now well known that viruses are unique in their mode of replication, which can be easily targeted (Selisko *et al.*, 2007; Syed *et al.*, 2010). They use specific enzymes to infect and replicate, whose inhibition could arrest their metabolism. For example, the proteolytic enzyme promotes virus maturation by separating the viral polyprotein precursor, whose inhibition will stop its maturation (Wapling *et al.*, 2007). So the virus metabolism or replication can be stopped by specific inhibitors.

Today many synthetic antiviral drugs e.g. moroxydine, ganciclovir, valganciclovir, valaciclovir etc. are used, which inhibit the virus replication via different mechanisms (Biron, 2006; Czeizel *et al.*, 2006). But difficulty in drug treatment arises due to their low efficiencies, cytotoxicity and development of viral resistance against them. Another antiviral treatment; vaccination, can be applied but they are still under development, as they often provide incomplete protection against virus and their reliability needs more research (Pervez, 2000b; Subbarao and Joseph, 2007). Thus the treatment through antiviral synthetic drugs and vaccines need more scientific investigation. Nature provides another, more reliable source of antiviral agents; viz. plants phytochemicals; almost 40% of currently available drugs are direct or indirect derivatives of plants. A number of ethnobotanical studies aiming to identify potential therapeutic plants for more effective control of health issues demonstrate the importance of plant species in health care system (Shinwari and Khan, 2000; Heneidy and Bidak, 2004; Appidi *et al.*, 2008; Ky *et al.*, 2009; Ansari and Inamdar, 2010; Makambila-Koubemba *et al.*, 2011). Plants are rich source of phytochemicals like alkaloids, anthocyanins, carotenoids, flavonoids, isoflavones, lignans, monoterpenes, organosulfides, phenolic acids, saponins and many more (Al-Yahya, 2005; Hassan *et al.*, 2006; Anitha and Ranjitha Kumari, 2006; Akomo *et al.*, 2009; Rahman *et al.*, 2009; Amabeoku and Kinyua, 2010; Ndjonka *et al.*, 2010). These phytochemicals have been proved to be responsible for their antimicrobial (Sampathkumar *et al.*, 2008; Krishnan *et al.*, 2010), antihypertensive (Amalia *et al.*, 2008), anti-diabetic (Qureshi *et al.*, 2009), antioxidant (Momtaz and Abdollahi, 2010), hepatoprotective (Mahalakshmi *et al.*, 2010; Ansari *et al.*, 2011), cardioprotective (Ojha *et al.*, 2008; Fard *et al.*, 2008) and other therapeutic activities. Thus this study is aimed to analyze the previously reported antiviral plants and identify potential mode of action and compounds that are responsible for their antiviral activity. Better understanding of natural antiviral agent's mode of action and identification of responsible compounds will be helpful to provide a new insight for the development of new antiviral drugs for more effective viral control.

**Basic viral structure and its mode of action:** Viruses are organic objects, which are metabolically inactive outside the host body but become active on their entry into the host cell (Dupre and O'Malley, 2009). These are mainly composed of proteins and nucleic acid; the proteins majorly contribute to their specific shape and form a coat called capsid (Andersson, 2010). Thus viruses are of various shapes e.g. simple, helical, icosahedral or complex and some viruses are surrounded by a lipid bilayer, derived from host membrane, which is called as envelope (Geng *et al.*, 2007; Raja *et al.*, 2003). Some capsid proteins are also associated with virus nucleic acid and called as nucleocapsid, while nucleic acid proteins, are the direct part of the nucleic acid, known as nucleoproteins. The nucleic acid of virus is either made up of DNA or RNA, is the basic

source of information required for the regulation of its metabolic activities. These DNA and RNA can be further divided into two types depending upon the number of strands i.e. single stranded or double stranded DNA/RNA (Firth *et al.*, 2010; Pichlmair *et al.*, 2006). The single stranded RNA viruses can be further distinguished depending upon the sense of strand as some RNA viruses have positive-sense RNA (+VE ssRNA) and some viruses have negative-sense RNA (-VE ssRNA) (Gorbelenya *et al.*, 2006). The shape of nucleic acid (DNA/RNA) is also an important source of differentiation, because all the viruses did not contain same-shape nucleic acid (Gao and Hu, 2007). It can be either in circular, linear or coiled form.

Virus (either DNA or RNA) life cycle can be divided into some predefined stages; adhesion, adsorption (entry), replication, maturation and release, which involve some enzymes and proteins. For example, the process of virus entry is carried out by cell surface proteins; HCV entry involves claudin-1, occludin, tetraspanin as main receptors proteins (Burlone and Budkowska, 2009). Its entry is also mediated by some other lipoproteins and an enzyme; lipoprotein lipase. On the other hand the influenza virus infection is mediated by protease enzyme, which activates the viral surface protein haemagglutinin (Zambon, 2001). The protease enzyme is also important in the expression of viral proteins; it splits the proteins into groups depending upon their structural and nonstructural functions (Appel *et al.*, 2006). But the RNA viruses need two additional enzymes for their survival; reverse transcriptase and integrase, former transcribes the viral RNA into DNA at the time of replication (Briones *et al.*, 2010; Sluis-Cremer and Tachedjian, 2008). While the second enzyme is used to incorporate the viral DNA into host genome, furthermore it is also needed for proper uncoating of virus core proteins. Thus virus is needy of enzymatic and non-enzymatic proteins, which can be targeted to stop their replication and infection.

**Antiviral plants:** In this review a total of 105 plant species have been identified that were reported for their potential antiviral activities (Fig. 1). Maximum number of plants were reported for their activity against herpes viruses, indicating that herpes viruses were the highly studied viruses with respect to antiviral plants. After herpes virus, HIV, influenza and hepatitis were among other viruses that were addressed in most of the studies in order to discover the plant with antiviral properties. In next sections a brief description with respect to the available antiviral plants against some important viruses has been provided.

**Plants with antiviral activity against herpes virus:** An enveloped double stranded DNA virus with linear genome; it belongs to the family Herpesviridae, which is recently reclassified to separate the mammal's virus from other non-mammalian viruses (Davison *et al.*, 2009). It is also




Fig. 1: Mostly studied viruses found during current review

known as human herpesvirus and Varicella-zoster virus. It is highly infectious and prevalent disease especially in developing countries with almost 50 % prevalence in adults, disease symptoms are often unnoticeable (WHO, 2001). It is a sexually transmitted disease and highly prevalent in females as they often get infected with it before the age of fertility (Ziyaeyan *et al.*, 2007). It also causes febrile rash and graft-versus-host disease in humans, who got hematopoietic stem cell transplantation (Pichereau *et al.*, 2011). It is also the casual agent of genital herpes disease; vaccines against it are currently under developmental stages (Soleimanjahi *et al.*, 2007).

Total 57 plants were identified with antiviral activity against herpes virus (Table 1). Binns *et al.* (2002) reported that h-hexane extracts of *Echinacea purpurea* showed in vitro antiviral activity against herpes virus in another study polysaccharide and cichoric acid were found among the active phytochemicals of various plant parts extracts (Vimalanathan *et al.*, 2005). Another compound of *Phyllanthus urinaria* from its acetone extract (hippomanin A) showed its activity against herpes simplex virus type 2 (Yang *et al.*, 2007); in another study Cheng *et al.* (2011) identifies a compound (excoecarianin) from this herb with same activity. Anthraquinones has been identified as potential compound responsible for antiviral properties of *Rhamnus frangula*, *Rhamnus purshianus* and *Rheum officinale* (Sydiskia *et al.*, 1991). This organic compound has also been reported in many studies to be responsible for positive health attributes of many medicinal plants (Kumar *et al.*, 2007; Hussein *et al.*, 2010; Karou *et al.*, 2011; Mbaya and Ibrahim, 2011; Sonibare *et al.*, 2011). Kurokawa *et al.* (1999) reported the antiviral activity of *Rhus javanica* against herpes virus and found moronic acid as potential active agent from its herbal extract. Plants mainly showed activities like restriction of entry host into cell (Weber *et al.*, 1992; Zandi *et al.*, 2007), reduced viral replication (Duarte *et al.*, 2001; Chiang *et al.*, 2003; Alche *et al.*, 2003) and partial destruction of viral envelope (Sydiskia *et al.*, 1991). Active compound present in most of the plants were anthraquinones, terpenes, quercetin, lectins and phenolics (Amoros *et al.*, 1987; Sydiskia *et al.*, 1991; Chiang *et al.*, 2003; Ooi *et al.*, 2004; Kan *et al.*, 2009). *Allium sativum* is a medicinal plant with a lot of health benefits (Ishtiaq *et al.*, 2007; Sukandar *et al.*, 2010; Abdelaziz and Kandeel, 2011; Weber *et al.*, 1992) found its extract effective against this virus. The extract of *Aloe vera* another important medicinal plant (Alqasoumi *et al.*, 2008; Semalty *et al.*, 2010) has been reported to be effective against this virus by inhibiting the viral entry and replication into the host cell(Zandi *et al.*, 2007).

**Plants with antiviral activity against HIV:** HIV is highly infectious enveloped virus of family Retroviridae; it has liner ssRNA positive sense genome and cause high morbidity. During year 2004, HIV infection was one of the leading factors responsible for almost 2.65 million deaths in low income countries (Patton *et al.*, 2009). It is also prevalent in high income countries, as in United States almost 55400 new HIV cases were observed each year from 2003-06 (Hall *et al.*, 2008). It increases the chances of bacteria and other viruses infections, which poorly effects the health; transmission of HIV from mother to child could lead to the death (Corbett *et al.*, 2003; Ahsan *et al.*, 2004; Ilboudo *et al.*, 2007; Abongo *et al.*, 2008). Its chemo-treatments are scarce, because of their potential side effects on human body, non-significant efficiencies and increased divergence in HIV genome (Fokunang *et al.*, 2006; Jaffary *et al.*, 2007; Kagone *et al.*, 2011). But a nutrient rich food can reduce the HIV caused decreased weight and help in improving the patient's health (Oguntibeju *et al.*, 2007).

In present study 26 plant species were found effective against HIV. Terpenoids, lectins, alkaloids and flavonoids were common among the active compound of these plants (Table 2). Roja and Heble (1995) reported the positive activity of an alkaloid (castanospermine) isolated from seeds

Table 1: Detail description of studies focusing on antiviral plant activities against herpes virus

Plant name	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<i>Achyrocline flaccida</i>	n/a	Aqueous extracts	n/a	In vitro, inhibits viral replication	Garcia et al. (1995)
<i>Allium sativum</i>	Aljoene, Allicin, Allyl methyl thiosulfinate and Methyl allyl thiosulfinate	Oil-macerates and fresh garlic extract	HeLa/Vero cells	Interfere with virus adsorption and penetration	Weber et al. (1992)
<i>Aloe barbadensis</i>	Anthraquinones: Aloe emodin	Hot glycerin extracts	Vero cells	Partially destroy the viral envelope and inactivate them	Sydiskia et al. (1991)
<i>Aloe vera</i>	n/a	Leaves' crude hot glycerin extract	Vero cells	Inhibits virus attachment, entry and replication	Zandi et al. (2007)
<i>Angellaria arvensis</i>	Triterpene saponin	n/a	n/a	In vitro; inhibits the virus replication	Amoros et al. (1987)
<i>Andrographis paniculata</i>	Diterpenes: Andrographolide, Neoandrographolide and 14-deoxy-11,12-didihydroandrographolide	n/a	n/a	n/a	Wiert et al. (2005)
<i>Bergenia ciliata</i>	n/a	Methanolic extracts	Vero cells	n/a	Rajbhandari et al. (2009)
<i>Bostrychia montagnei</i>	Sulfated polysaccharides	Cold and Hot water extracts	Vero cell	Inhibits the virus replication	Duarte et al. (2001)
<i>Caesalpinia pulcherrima</i>	Quercetin derivatives	Aqueous extracts of fruit, stem, leaf, fruit and seed	Human skin basal cell carcinoma cell line (BCC-1/KMC)	Inhibits the virus replication at early stages of cycle	Chiang et al. (2003)
<i>Canavalia ensiformis</i>	Lectins	n/a	n/a	In vitro, inhibits virus penetration	Marchetti et al. (1996)
<i>Cassia angustifolia</i>	Anthraquinones	Hot glycerin extracts	Vero cells, WI-38 cells	Partially destroy the viral envelope and inactivate them	Sydiskia et al. (1991)
<i>Cedrela tubiflora</i>	Acidic polysaccharides	Leaves fraction extracts	n/a	Inhibits virus replication	Craig et al. (2001)
<i>Cicer arietinum</i>	Phenolic compounds	Seed, fruit skin and aerial parts	Madin-darby Bovine Kidney/Vero cells	n/a	Kan et al. (2009)
<i>Clerodendrum Inerme</i>	n/a	Methanol extracts	n/a	Reduce the viral cytopathic effect	Vimalanathan et al. (2009)
<i>Cocos nucifera</i>	Catechin, epicatechin and B-type procyanidins	Husk fiber' water extract	n/a	n/a	Esquenazi et al. (2002)
<i>Conyza canadensis</i>	n/a	Ethyl acetate, chloroform, butanol and methanol extracts of the aerial parts	Human diploid embryonic lung fibroblasts (MRC-5)	n/a	Edziri et al. (2011)
<i>Dianthus Caryophyllus</i>	n/a	Seed crude extracts	Vero and HepG2 cells	n/a	Barakat et al. (2010)

Table 1: Continued

Plant name	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<i>Echinacea purpurea</i>	alkenes and amides	Root's n-hexane extracts	<i>In vitro</i>	n/a	Binns et al. (2002)
<i>Echinacea purpurea</i>	Polysaccharide, Cichoric acid and others	Stems, leaves and flowers water, ethyl acetate, ethanolic and other fractions	Vero cells	n/a	Vimalanathan et al. (2005)
<i>Ceum japonicum</i>	n/a	Hot water extract	Immunosuppressed mice; it suppressed the activity of reverse transcriptase enzyme	n/a	Kageyama et al. (1996)
<i>Glyptopetalum sclerocarpum</i>	Sesquiterpene: Sclerocarpic acid	Sesquiterpene extraction	n/a	n/a	Sotanaphun et al. (1999)
<i>Hamelia virginiana</i>	Oligomeric to polymeric proanthocyanidins	Ultrafiltered crude hydroalcoholic extract	n/a	n/a	Erdelmeier et al. (1996)
<i>Hancornia speciosa</i>	n/a	Ethanolic extract	n/a	n/a	Brandao et al. (2011)
<i>Feliciastrum aureonitens</i>	Galangin	Shoots extracts	n/a	n/a	Meyer et al. (1997)
<i>Himatanthus phagedaeonica</i>	n/a	Ethanolic extract	n/a	n/a	Brandao et al. (2011)
<i>Hysopos officinalis</i>	n/a	Leaves methanolic extracts	Mice and Vero E6 cells; reduce the virus cell number via different mechanism than commercially used drug, acyclovir	n/a	Behbahani (2009)
<i>Macula cochinchinensis</i>	Morin	Ethyl acetate and methanol extracts	<i>In vitro</i>	n/a	Bunyapraphatsara et al. (2000)
<i>Melia azedarach</i>	Meliacarpin	Ethyl acetate extract of leaves	Vero cells	Inhibits the virus replication	Alche et al. (2003)
<i>Melissa officinalis</i>	n/a	Volatile oils	HEP-2 cells	Inhibits virus replication	Allahverdiyev et al. (2004)
<i>Monardica charantia</i>	n/a	n/a	Vero cells	n/a	Praseno and Rintiswati (1997)
<i>Myrica rubra</i>	Tanin (prodelphinidin B-2,3,3'-di-O-gallate)	Bark	Vero cell with cell	Inhibits the viral attachment	Cheng et al. (2003)
<i>Ocimum americanum</i>	n/a	Dichloromethane extract	Green monkey kidney cells	n/a	Yncharoen et al. (2011)
<i>Ocimum basilicum</i>	n/a	Dichloromethane extract	Green monkey kidney cells	n/a	Yncharoen et al. (2011)

Table 1: Continued

Plant name	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<i>Oximum sanctum</i>	n/a	Dichloromethane and methanol extract	Green monkey kidney cells	n/a	Yncharoen et al. (2011)
<i>Ouratea castaneofolia</i>	n/a	Ethanol extract	n/a	n/a	Brando et al. (2011)
<i>Ouratea semiserrata</i>	n/a	Ethanol extract	n/a	n/a	Brando et al. (2011)
<i>Ouratea speciosiflora</i>	n/a	Ethanol extract	n/a	n/a	Brando et al. (2011)
<i>Pandanus amaryllifolius</i>	lectin (pandanin)	Saline extract of the leaves	n/a	n/a	Ooi et al. (2004)
<i>Phyllanthus orbicularis</i>	n/a	Aqueous extract of leaves and stems	n/a	Inhibition of viral entry	Del Barrio and Parra (2000)
<i>Phyllanthus urinaria</i>	Exocercarain	Acetone extract	n/a	n/a	Cheng et al. (2011)
<i>Phyllanthus urinaria</i>	Hippomanin A	Acetone extract	n/a	n/a	Yang et al. (2007)
<i>Podophyllum peltatum</i>	Podophyllotoxin	Aqueous extract	n/a	n/a	Bedows and Hattfield (1982)
<i>Polygonum spectabile</i>	n/a	Ethanol extract	n/a	n/a	Brando et al. (2011)
<i>Pterocaulon sphacelatum</i>	Lactone (acanthoaustralide-1-O-acetate) and flavonoids (quercetin and chrysosplenol D)	Hydroethanol extract	n/a	n/a	Rocha Martins et al. (2011)
<i>Punica granatum</i>	Tannin	Pericarp	n/a	n/a	Zhang et al. (1995)
<i>Quillaja saponaria</i>	Triterpenoid saponins	Aqueous extracts	n/a	n/a	Rener et al. (2007)
<i>Rhamnus frangula</i>	Anthraquinones	Glycerin extracts	n/a	n/a	Sydiskia et al. (1991)
<i>Rhamnus purshianus</i>	Anthraquinones	Glycerin extracts	n/a	n/a	Sydiskia et al. (1991)
<i>Rhus officinale</i>	Anthraquinones	Glycerin extracts	n/a	n/a	Sydiskia et al. (1991)
<i>Rhinacanthus nasutus</i>	Naphthoquinones [rhinacanthin-C and rhinacanthin-D]	n/a	n/a	n/a	Sendl et al. (1996)
<i>Rhus aromatica</i>	n/a	Aqueous extract of root/stem bark	Cell culture	Prevent viral penetration in host cell	Reichling et al. (2009)
<i>Rhus javanica</i>	Moronic acid	Herbal extract	Mice	n/a	Kurokawa et al. (1999)
<i>Sapindaria officinalis</i>	n/a	Lyophilized infusion	n/a	n/a	Serkedjieva et al. (2006)
<i>Sclerotium glucanicum</i>	Scleroglucan	n/a	n/a	Inhibits virus binding	Marchetti et al. (1996)
<i>Spirulina plantensis</i>	n/a	Phosphate buffer and water extracts	n/a	n/a	Shalaby et al. (2010)
<i>Syzygium aromatic</i>	Engenin	Bud	n/a	n/a	Takechi and Tanaka (1981)

n/a = not available

Table 2: Detail description of studies focusing on antiviral plant activities against HIV

Plant name	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<i>Agastache rugosa</i>	Rosmarinic acid	Root's aqueous methanolic extract	<i>In vitro</i>	Inhibits virus integrase enzyme	Kim <i>et al.</i> (1999)
<i>Calophyllum brasiliense</i>	Apetalic acid, Calanolides B and C	Hexane extract of the leaves	n/a	Inhibitory effect on reverse transcriptase	Cesar <i>et al.</i> (2011)
<i>Calophyllum cerasiferum</i>	Inophylum, Calanolide A and Coumarins	n/a	n/a	Inhibitory effect on reverse transcriptase	Saeed and Hussain (2006)
<i>Calophyllum inophylum</i>	Inophylum B and P	Leaves	n/a	n/a	Laure <i>et al.</i> (2008)
<i>Calophyllum teysmannii</i>	Costatolide	Latex	n/a	n/a	Fuller <i>et al.</i> (1994)
<i>Castanospernum austrole</i>	Alkaloid: Castanospermine	Seeds	n/a	n/a	Roja and Heble (1995)
<i>Corydalis yanhusuo</i>	Gossypol and alkaloids	n/a	n/a	Inhibitory effect on reverse transcriptase	Wang and Ng (2001)
<i>Daucus maritimus</i>	n/a	Seeds	n/a	Inhibitory effect on reverse transcriptase	Miladi <i>et al.</i> (2011)
<i>Gelonium multiflorum</i>	Protein (GAP31)	Protein	T-lymphocytes	Inhibits the viral DNA integration to the host DNA and viral replication	Lee-Huang <i>et al.</i> (1995)
<i>Geum japonicum</i>	n/a	Hot water extract	Immunosuppressed mice	Inhibitory effect on reverse transcriptase	Kageyama <i>et al.</i> (1996)
<i>Kadsura heteroclita</i>	Triterpenoid and lignans (dibenzocyclooctadiene-type)	Stems	n/a	n/a	Pu <i>et al.</i> (2008)
<i>Monotes africanus</i>	Flavonoids	Organic extract	n/a	Inhibitory effect on reverse transcriptase	Meragelman <i>et al.</i> (2001) and Routtrakul <i>et al.</i> (2007)
<i>Panax ginseng</i>	Protein panaxagin	n/a	n/a	Inhibitory effect on reverse transcriptase	Ng and Wang (2001)
<i>Phaseolus lunatus</i>	Trypsin inhibitors	n/a	n/a	Inhibitory effect on reverse transcriptase	Wang and Ng (2001)
<i>Phaseolus vulgaris</i>	Lectin	n/a	n/a	Inhibitory effect on reverse transcriptase	Ye <i>et al.</i> (2001) and Fang <i>et al.</i> (2010)
<i>Prunella vulgaris</i>	n/a	Aqueous extracts	n/a	Interference of early, post-virion binding events	Oh <i>et al.</i> (2011)
<i>Quillaja saponaria</i>	Triterpenoid saponins	Aqueous extracts	n/a	n/a	Ronen <i>et al.</i> (2007)
<i>Rhizophora apiculata</i>	Polysaccharide extracted from the leaf	n/a	Cell culture systems	Blocked the binding of HIV-1 virions	Premarnathan <i>et al.</i> (1999b)
<i>Rhizophora mucronata</i>	Polysaccharide bark	Alkaline extract	n/a	Inhibited the viral binding to the cell	Premarnathan <i>et al.</i> (1999a)
<i>Rhus succedanea</i>	Flavonoids	n/a	n/a	Inhibitory effect on reverse transcriptase	Lin <i>et al.</i> (1997)

Table 2: Continued

Plant name	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<i>Ricinus communis</i>	Lectin	n/a	n/a	Inhibitory effect on reverse transcriptase and the N-glycosidases	Wang and Ng (2001)
<i>Shepherdia argentea</i>	Tannins (shephagennins A and B)	Leaf extract	n/a	Inhibitory effect on reverse transcriptase	Yoshida <i>et al.</i> (1996)
<i>Terminalia chebula</i>	n/a	Methanol	Virus-infected baby hamster kidney	n/a	Lee <i>et al.</i> (2011)
<i>Trichosanthes kirilowii</i>	Trichosanthes anti-HIV protein	Protein	n/a	n/a	Lee-Huang <i>et al.</i> (1991)
<i>Vigna unguiculata</i>	Ungulin	Seed protein	n/a	Inhibiting effect on reverse transcriptase and the glycohydrolases $\alpha$ - and $\beta$ -glucosidases	Ye <i>et al.</i> (2000) and Ye and Ng (2001)

n/a: Not available

of *Castanospermum austral* against this virus. Wang and Ng (2001) reported that gossypol and alkaloids of *Corydalis yanhusuo* might be responsible for its inhibitory activity on HIV virus. Ethanolic extract of *Monotes africanus*, which is a rich source of various flavonoids has been reported for its antiviral activity against HIV (Meragelman *et al.*, 2001; Reutrakul *et al.*, 2007). Most of the plants mainly affect the Reverse Transcriptase (RT) activity or penetration of virus particles into host cell. Calanolides which exhibits potential RT inhibitory activity have been found in the hexane extract of *Calophyllum brasiliense* leaves (Cesar *et al.*, 2011). The hexane extract of this plant was found to have inhibitory effect on RT enzyme of HIV. Lectins derived from *Phaseolus vulgaris* are reported for their potential inhibitory effect on HIV RT activity (Ye *et al.*, 2001; Fang *et al.*, 2010). Another seed protein isolated from *Vigna unguiculata* possesses the same effect on RT activity in addition to inhibitory effect on glycohydrolases  $\alpha$ - and  $\beta$ -glucosidases (Ye *et al.*, 2000; Ye and Ng, 2001). Oh *et al.* (2011) reported that aqueous extract of *Prunella vulgaris* interferes with the viron post binding events. Extract of *Rhizophora mucronata* and *Rhizophora apiculata* (a rich source of polysaccharides) blocks the viral binding to the cell surface (Premanathan *et al.*, 1999a; Premanathan *et al.*, 1999b). In another study Lee-Huang *et al.* (1995) studied the effect of a protein isolated from *Gelonium multiflorum* and found that it reduces the virus replication by inhibiting the integration of viral DNA into host genome. *Panax ginseng*, which is also effective against cardiovascular diseases (Jun *et al.*, 2007) has been reported to be effective against HIV by inhibiting the RT activity (Ng and Wang, 2001). *Ricinus communis* an important medicinal plant (Onwuliri and Anekwe, 2001) showed inhibitory effect on RT and N-glycohydrolases of HIV (Wang and Ng, 2001). Another plant *Terminalia chebula*, which has been proven to have many beneficial medicinal properties (Gupta *et al.*, 2008a, b; Shinde *et al.*, 2009; Anam *et al.*, 2009) posses significant anti-HIV activity (Lee *et al.*, 2011). Its methanolic extract showed antiviral activity against HIV when tested on virus infected baby hamster kidney cells.

**Plants with antiviral activity against influenza virus:** It is a single stranded RNA virus with negative sense linear fragmented genome enclosed in a capsid, it belongs to family Orthomyxoviridae and infects both mammals and birds. It has seasonal epidemiology and mainly spread through air when the environment is dry and cold (Lowen *et al.*, 2007). The hospitalization and death rate associated with influenza vary with the age and type of virus, as influenza virus A-caused infection rate is higher than type-B (Thompson *et al.*, 2004). Influenza virus, especially of avian origin is highly virulent disease causing agent in humans and birds; it has long history, put huge burden on human health since 1580 (Farooq *et al.*, 2006; Lazzari and Stohr, 2004). It is responsible for deaths of millions of people; its pandemic nature is due to its variable strains which develop via the reassortment of genetic information. Thus the development of vaccine against it is difficult both in humans and birds.

Sixteen plants were identified which showed antiviral activity against influenza virus. Anthocyanin and polyphenols were among the commonly found active phytochemicals in these plants (Table 3). *Camellia sinensis* is an important herbal plant having significant antioxidant, photochemoprotective, hyperglycemic, antibacterial and many other beneficial heath benefits (Adiloglu and Adiloglu, 2006; Bakar *et al.*, 2006; Shokrzadeh *et al.*, 2006; Hassani *et al.*, 2008; Mohamed and Metwally, 2009; Amutha *et al.*, 2010; Chakraborty and Chakraborti, 2010; Kaur and Saraf, 2011; Obaid *et al.*, 2011). Catechin derivatives obtained from the tea of this plant showed significant inhibitory effect on influenza strains (Song *et al.*, 2005). Ehrhardt *et al.* (2007)

Table 3: Detail description of studies focusing on antiviral plant activities against influenza virus

Plant name	Active compounds	Extract type	Model organism	Mode of action	Reference
<i>Agrimonia pilosa</i>	n/a	n/a	MDCK cells, embryonated chicken eggs	Inhibits RNA synthesis and react with viral membrane	Shin <i>et al.</i> (2010)
<i>Aloe barbadensis</i>	Anthraquinones; Aloe emodin	Hot glycerin extracts	Vero cells	Partial destruction of viral envelope	Sydiska <i>et al.</i> (1991)
<i>Aronia melanocarpa</i>	Anthocyanins	Fruit juice	n/a	n/a	Valcheva-Kuzmanova and Belcheva (2006)
<i>Bergenia ciliata</i>	n/a	Methanolic extracts	MDCK (Madin-Darby Canine Kidney) cells	n/a	Rajbhandari <i>et al.</i> (2009)
<i>Camellia sinensis</i>	Catechin derivatives	Tea	MDCK cell	Inhibits virus replication and hemagglutination	Song <i>et al.</i> (2005)
<i>Cistus incanus</i>	Polyphenol; CYSTUS052	n/a	A549 or MDCK cells; Human patients	Inhibits the viral cell entry by modulating the viral surface structures	Elhrhardt <i>et al.</i> (2007) and Kalus <i>et al.</i> (2009)
<i>Clinanthus siamensis</i>	n/a	Leaf's ethanolic extracts	MDCK cells and mouse	Produce antibodies against virus	Wirotesangthong <i>et al.</i> (2009)
<i>Commelina communis</i>	Alkaloids	n/a	Mice and Madin-Darby canine kidney cells	Inhibits the virus growth and reduce viral titres in lungs	Bing <i>et al.</i> (2009)
<i>Echinacea purpurea</i>	n/a	Commercial extract; Echinaforce®	H-1 (subclone of HeLa cells) cells	Resists virus entry by inhibiting receptor binding and replication	Vimalanathan <i>et al.</i> (2005) and Pleschka <i>et al.</i> (2009)
<i>Ceratium sanguineum</i>	Polyphenols	Methanolic extracts	Murine model	Effect the expression of virus proteins on cell surface	Serkedjieva (1996) and Sokmen <i>et al.</i> (2005)
<i>Narcissus tazetta</i>	Protein (Narcissus tazetta lectin)	Bulbs	n/a	n/a	Ooi <i>et al.</i> (2010)
<i>Pandanus amaryllifolius</i>	Lectin (pandanin)	Saline extract of the leave	n/a	n/a	Ooi <i>et al.</i> (2004)
<i>Rhinacanthus nasutus</i>	Lignans [Rhinacanthin E and Rhinacanthin F]	Aerial parts	n/a	n/a	Kernan <i>et al.</i> (1997)
<i>Saponaria officinalis</i>	n/a	Lyophilized infusion	n/a	n/a	Serkedjieva <i>et al.</i> (2006)
<i>Scutellaria baicalensis</i>	Isoscutellarein-8-methylether	Roots	n/a	Inhibit viral replication	Nagai <i>et al.</i> (1995)
<i>Toddalia asiatica</i>	n/a	Extract	n/a	n/a	Luet <i>et al.</i> (2005)

n/a = Not available

found that *Cistus incanus* possess the antiviral activity by modulating the viral surface in order to inhibit its entry into Madin-Darby Canine Kidney (MDCK) cells. Inhibitory effect against influenza virus has also been reported in human patients (Kalus *et al.*, 2009). Another herb *Echinacea purpurea* also restrict the viral entry by inhibiting the binding of viral receptors to the host cells surface (Pleschka *et al.*, 2009). *Geranium sanguineum*, a rich source of polyphenols showed antiviral activity by effecting the expression of viral proteins on cell surface (Serkedjieva, 1996; Sokmen *et al.*, 2005). A lectin isolated from bulbs of *Narcissus tazetta* showed significant antiviral activity against various strains of influenza virus (Ooi *et al.*, 2010). In another study Ooi *et al.*, (2004) reported the anti-influenza activity of a lectin isolated from saline extract of *Pandanus amaryllifolius* leaves. *Scutellaria baicalensis* an important medicinal plant (Yeh *et al.*, 2010) possess a compound (isoscutellarein-8-methylether) which is thought to be responsible for it antiviral activity against influenza virus (Nagai *et al.*, 1995).

**Plants with antiviral activity against Hepatitis C Virus (HCV):** Hepatitis C is an enveloped virus with +VE single stranded linear RNA, which belongs to the family Flaviviridae. It cause mild-chronic liver disease infecting more than three million people each year, which results in almost 350 000 deaths (WHO, 2011a). It has worldwide spread with 4.8% infection rate in Pakistan. It is a global problem nearly affecting 130 million people; moreover, alone it is responsible for about 27% of cirrhosis and almost 25% of world's hepatocellular carcinoma (Alter, 2007). Its high spread is attributed to poor moral and health conditions, use of drugs, alcohol and contaminated syringes (Roshandel *et al.*, 2007; Zakizad *et al.*, 2009). It is detectable through serum proteins and blood analysis; its chemotherapeutic treatment is difficult but can be treated through herbal products (Ansari *et al.*, 2011; Joseph and Raj, 2011; Pervez, 2000a; Moundipa *et al.*, 2007; Tabassum *et al.*, 2000).

In current study six plants have been identified with proved antiviral activity against HCV. Mainly plant showed inhibitory effect on HCV protease. Hussein *et al.* (2000) reported the inhibitory effect of *Trachyspermum ammi* and *Embelia schimper* methanol extract on HCV protease. *Solanum nigrum* which also exhibits hepatoprotective activity (Subash *et al.*, 2011) has also been reported to have inhibitory effect on HCV (Javed *et al.*, 2011). Cannabis a commonly used drug (Richardson, 2010) is a phytochemical of *Cannabis sativa* a medicinal plant with many beneficial health effects (Arshad and Khan, 2000; Qureshi *et al.*, 2001; Tehranipour and Ebrahimpour, 2009). *Acacia nilotica* has remained the focus of many studies for its multipurpose applications (Shirazi *et al.*, 2001; Banerjee *et al.*, 2004; Ghosh *et al.*, 2004; Elkhalifa *et al.*, 2005; Emtehani and Tabari, 2007) its acetonnic and methanolic extracts have sown anti-HCV effect on liver cells (Rehman *et al.*, 2011). Sylvestre *et al.* (2006) reported the anti-HCV activity of cannabis in human patients. Mainly these plants showed their activity against HCV by targeting its protease (Table 4).

**Plants with antiviral activity against respiratory syncytial virus:** It belongs to the family Paramyxoviridae and is single stranded RNA virus, which is enclosed in an envelope. It is a highly prevalent virus throughout the world with large variation in its genome (Trento *et al.*, 2010). It cause bronchiolitis and other respiratory problems and is a major cause of lower respiratory tract infections; it mostly infects the children under age of six months and also responsible for asthma problems (Mohapatra and Boyapalle, 2008). It causes an infection in 3-7% of healthy adults and 4-7% in the non-healthy adults, who already suffers from lung and heart diseases (Falsey *et al.*,

Table 4: Detail description of studies focusing on antiviral plant activities against hepatitis, respiratory syncytial virus and vesicular stomatitis virus

Plant name	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<b>Hepatitis A Virus</b>	n/a	Phosphate buffer and water extracts	n/a	n/a	Shalaby <i>et al.</i> (2010)
<i>Spirulina platensis</i>	n/a	Seed crude extracts	Vero and HepG2 cells	n/a	Barakat <i>et al.</i> (2010)
<i>Dianthus caryophyllus</i>	n/a	Methyl ester dehydrochelanic acid and methyl brevifolin carboxylate	n/a	n/a	Zhong <i>et al.</i> (1998)
<b>Hepatitis B Virus</b>					
<i>Phyllanthus urinaria</i>	Methyl ester dehydrochelanic acid and methyl brevifolin carboxylate	Seed crude extracts	Vero and HepG2 cells	n/a	
<i>Trachyspermum ammi</i>	n/a	Methanol	n/a	Inhibitory HCV proteases	Hussain <i>et al.</i> (2000)
<i>Endelia schimperi</i>	Benzoguione (embelin (I) and 5-O-methylembelin (II))	Methanol	n/a	Inhibitory HCV proteases	Hussain <i>et al.</i> (2000)
<i>Solanum nigrum</i>	n/a	Methanol and chloroform extracts of seeds	n/a	Inhibitory effect on NS3 protease	Javed <i>et al.</i> (2011)
<i>Acacia nilotica</i>	n/a	Acetonic and methanolic extract	Liver cells	n/a	Rehman <i>et al.</i> (2011)
<i>Cannabis sativa</i>	Cannabis	n/a	Human patients	n/a	Sylvestre <i>et al.</i> (2006)
<i>Daucus maritimus</i>	n/a	Seeds	n/a	Inhibits the reverse transcriptase activity	Miladi <i>et al.</i> (2011)
<b>Respiratory Syncytial Virus</b>					
<i>Narcissus tazetta</i>	Protein (Narcissus tazetta lectin)	Bulbs	n/a	n/a	Ooi <i>et al.</i> (2010)
<i>Selaginella sibirica</i>	Amentoflavone	Ethanol extract	n/a	n/a	Ma <i>et al.</i> (2001)
<i>Schefflera heptaphylla</i>	3,4-Di-O-cafeoylquinic acid and 3,5-di-O-cafeoylquinic acid	n/a	n/a	inhibition of virus-cell fusion	Li <i>et al.</i> (2005)
<i>Barleria prionitis</i>	Iridoid: 6-O-trans-p-coumaroyl-8-O-acetylshanzhisiide methyl ester and its cis isomer	n/a	n/a	n/a	Chen <i>et al.</i> (1998)
<i>Morhamia lutea</i>	Luteoside, verbascoside and isoverbascoside	Roots	n/a	n/a	Kerman <i>et al.</i> (1998)
<b>Vesicular Stomatitis Virus</b>					
<i>Allium sativum</i>	Alloene, allixin and allyl methyl thiosulfinate	Oil/macerates and Fresh garlic extract	HeLa/Vero cells	interfere with virus adsorption and penetration	Weber <i>et al.</i> (1992)
<i>Cedrela tubiflora</i>	Andic polysaccharides	Leaves fraction extracts	n/a	Inhibits virus replication	Craig <i>et al.</i> (2001)
<i>Justicia procumbens</i>	Justicidins A-B and diphyllin derivatives	Aerial parts; methanolic extract	Cultured rabbit lung cells (RL-33)	n/a	Asano <i>et al.</i> (1996)
<i>Melia azedarach</i>	Meliacarpin; Tetranortriterpenoid: Ethyl acetate extract of leaves	Vero cells	Inhibits the virus replication; inhibits viral nucleocapsid uncoating and stops the movement of G-protein thus stop exocytosis of virus	Alche <i>et al.</i> (2003), Barquero <i>et al.</i> (2004)	

n/a: Not available

2005). Moreover it cause burden on population equal to that of influenza A virus, as the patients needs more health facilities and care. Ma *et al.* (2001) found that ethanol extract of *Selaginella sinensis* possess significant activity against respiratory syncytial virus. They further elaborated that this activity of plant could be due to the presence of a biflavonoid (amentoflavone) in its plant extract. A lectin isolated from *Narcissus tazetta* also showed antiviral activity against this virus (Ooi *et al.*, 2010). Li *et al.* (2005) reported the *Schefflera heptaphylla* antiviral activity against respiratory syncytial virus and concluded that this activity was due to the inhibition of fusion of viral particles to the host cell.

**Plant with broad spectrum antiviral activity:** Based on the data presented in Table 1-5 plant with broad spectrum antiviral activity (plants reported for their antiviral activity against two or more than two viruses) were identified. Total 24 plants were found that were resistant to two or more than two viruses (Fig. 2). Two plant species showed antiviral activity against four viruses.




Fig. 2: Plant with broad spectrum antiviral activity

Table 5: Detail description of studies addressing rarely studied viruses

Plant name	Virus	Active compounds	Plant part/Extract type	Model organism	Mode of action	Reference
<i>Aegle marmelos</i>	Human coxsackieviruses	Marmelide	Leaf, root, stem and fruit extracts	Vero cells	Inhibits viral replication at various steps e.g. adsorption penetration etc.	Badam <i>et al.</i> (2002)
<i>Allium sativum</i>	Parainfluenza virus-3 and Human rhinovirus-2	Ajoene, allixin and allyl methyl thiosulfinate	Oil-macerates and fresh garlic extract	HeLa / Vero cells	Interfere with virus adsorption and penetration	Weber <i>et al.</i> (1992)
<i>Anagallis arvensis</i>	Poiovirus	Triterpene saponin	n/a	n/a	Inhibits the virus replication	Amoros <i>et al.</i> (1987)
<i>Azadirachta indica</i>	Dengue virus	n/a	Leaves' aqueous extract	<i>Ades albopictus</i> ' larval cells and Mice	Inhibits virus replication	Parida <i>et al.</i> (2002)
<i>Caesalpinia pulcherrima</i>	Adenoviruses	Quercetin derivatives	Aqueous extracts of fruit, stem, leaf, fruit and seed	Human skin basal cell carcinoma cell line (BCC-1/KMC)	Inhibits the virus replication Chiang <i>et al.</i> (2003) at early stages of cycle	
<i>Canavalia ensiformis</i>	Rubella virus	Lectins	n/a	n/a	Inhibits replication	Marchetti <i>et al.</i> (1996)
<i>Cicer arietinum</i>	Parainfluenza-3 virus	Phenolic compounds	Seed, fruit skin and aerial parts	Madin-Darby bovine kidney/Vero cells	n/a	Kan <i>et al.</i> (2009)
<i>Conyza canadensis</i>	Coxsackie B virus type 3	n/a	Ethyl acetate, chloroform, butanol and methanol extracts of the aerial parts	Human diploid embryonic lung fibroblasts (MRC-5)	n/a	Edziri <i>et al.</i> (2011)
<i>Daucus maritimus</i>	Dengue virus and West Nile virus	n/a	Seeds	n/a	Inhibits the reverse transcriptase	Miladi <i>et al.</i> (2011)
<i>Glycyrrhiza glabra</i>	Japanese encephalitis virus	Glycyrrhizin and licorice	n/a	Porcine stable kidney (PS) and Human cervical carcinoma (HeLa) cell	n/a	Badam (1997)
<i>Helichrysum aureonitens</i>	Coxsackie B virus-1 and Reovirus	Coxsackie B virus-1 and Galangin	Shoots extracts	n/a	n/a	Meyer <i>et al.</i> (1997)
<i>Hibiscus sabdariffa</i>	Measles virus	n/a	Leaves ethanol extract	Hep-2 cells	n/a	Sunday <i>et al.</i> (2010)
<i>Podophyllum peltatum</i>	Measles virus	Podophyllotoxin	Aqueous extract	n/a	n/a	Bedows and Hatfield (1982)
<i>Prunus mume</i>	Human rhinovirus	n/a	Fruit juice	n/a	n/a	Song <i>et al.</i> (2010)
<i>Pterocaulon sphacelatum</i>	Picornavirus	Flavonoid (chrysosplenol C)	Ethanolic extract of aerial parts	n/a	n/a	Semple <i>et al.</i> (1999)
<i>Pterocaulon sphacelatum</i>	Poiovirus	n/a	Hydroethanol extract	n/a	n/a	Recha Martins <i>et al.</i> (2011)
<i>Quillaja saponaria</i>	Reovirus	Triterpenoid saponins	Aqueous extracts	n/a	n/a	Rener <i>et al.</i> (2007)
<i>Sophora flavescentis</i>	Coxsackievirus B3	Sophoridine	n/a	Primarily cultured myocardial cells	n/a	Zhang <i>et al.</i> (2006)
<i>Zingiber officinale</i>	Rhinovirus	Beta-sesquiphellandrene	Dried rhizomes	n/a	n/a	Denyer <i>et al.</i> (1994)

n/a = Not available

*Allium sativum* showed resistance against Herpes virus, Parainfluenza virus-3, Rhinovirus and Vesicular stomatitis virus. *Daucus maritimus* has been reported for its activity against Dengue virus, Hepatitis C virus, HIV and West Nile virus. *Helichrysum aureonitens*, *Pterocaulon sphacelatum* and *Quillaja saponaria* have been reported for their antiviral activity against three viruses each. As shown in figure 2 *Helichrysum aureonitens* is effective against Coxsackievirus, Herpes virus and Reovirus. *Pterocaulon sphacelatum* showed antiviral activity against Herpes virus, Picornavirus and Polio virus. *Quillaja saponaria* has been reported for its activity against Herpes virus, HIV and Reovirus. Almost all plants with broad spectrum activity were effective against herpes viruses. Moreover five plants which were effective against Herpes virus, also showed antiviral activity against Influenza virus. In the same way three plant species showed resistance against both Herpes virus and Vesicular stomatitis virus.

## CONCLUSION

Majority of the studied viruses belongs to the Flaviviridae, Herpesviridae and Picornaviridae family. A large number of plants are available in nature which could act as a source of lead antiviral compounds. Mainly these plants target the enzymes that are involved in replication and integration of virus into host cell. In case of DNA viruses restricted entry of viral particles into host cell or inhibition of viral replication into the host cells were most frequent mode of actions. Destruction of viral envelop was also one of the identified mode of action against DNA viruses. RT plays a significant role in replication of RNA viruses and most of the plants restrict the activity of RT enzyme of virus. In order to design effective drugs against viruses their enzymes that are involved in the key metabolic activities (integration, replication) should be focused. A lot of plant population with possible potential is still uncovered as few plants have been studied in detail in order to identify the active phytochemicals against these viruses. More detailed studies in future will help not only to identify the potential antiviral compounds but also in better understanding of their mode of action for more effective control of these lethal viruses.

## REFERENCES

- Abdelaziz, I. and M. Kandeel, 2011. The protective effects of *Nigella sativa* oil and *Allium sativum* extract on amikacin-induced nephrotoxicity. *Int. J. Pharmacol.*, 7: 697-703.
- Abongo, B.O., M.N.B. Momba, V.K. Malakate and J.N. Mwambakana, 2008. Prevalence of *Escherichia coli* O157:H7 among Diarrhoeic HIV/AIDS patients in the Eastern Cape Province-South Africa. *Pak. J. Biol. Sci.*, 11: 1066-1075.
- Adiloglu, A. and S. Adiloglu, 2006. An investigation on nutritional status of tea (*Camellia sinensis* L.) grown in eastern black sea region of Turkey. *Pak. J. Biol. Sci.*, 9: 365-370.
- Ahasan, M.M., M.M. Billah, M.M. Hasan, K.M.D. Islam and J.A. Shilpi, 2004. Transmission, biochemical manifestation and CD4<sup>+</sup> cell count of HIV: A review. *Pak. J. Biol. Sci.*, 7: 292-300.
- Akomo, E.F.O., C. Zongo, S.D. Karou, L.C. Obame, A. Savadogo, C. Atteke and A.S. Traore, 2009. *In vitro* antiplasmodial and antibacterial activities of *Canthium multiflorum* schum and thonn (Rubiaceae) extracts. *Pak. J. Biol. Sci.*, 12: 919-923.
- Al-Yahya, M.A., 2005. Preliminary phytochemical and pharmacological studies on the rind of pomegranate (*Punica granatum* L.). *Pak. J. Biol. Sci.*, 8: 479-481.
- Alche, L.E., G.A. Ferek, M. Meo, C.E. Coto and M.S. Maier, 2003. An antiviral meliacarpin from Leaves of *Melia azedarach* L. *J. Nat. Res. Part C: Biochem. Biol. Biophys. Virol.*, 58: 215-219.

- Allahverdiyev, A., N. Duran, M. Ozguven and S. Koltas, 2004. Antiviral activity of the volatile oils of *Melissa officinalis* L. against Herpes simplex virus type-2. *Phytomedicine*, 11: 657-661.
- Alqasoumi, S.I., T.A. Al-Howiriny and M.S. Abdel-Kader, 2008. Evaluation of the hepatoprotective effect of *Aloe vera*, *Clematis hirsute*, *Cucumis prophetarum* and bee propolis against experimentally induced liver injury in rats. *Int. J. Pharmacol.*, 4: 213-217.
- Alter, M.J., 2007. Epidemiology of hepatitis C virus infection. *World J Gastroenterol.*, 13: 2436-2441.
- Amabeoku, G.J. and C.G. Kinyua, 2010. Evaluation of the anticonvulsant activity of *Zanthoxylum capense* (Thunb.) Harv. (Rutaceae) in mice. *Int. J. Pharmacol.*, 6: 844-853.
- Amalia, L., E.Y. Sukandar, R.M.A. Roesli and J.I. Sigit, 2008. The effect of ethanol extract of kucai (*Allium schoenoprasum* L.) bulbs on serum nitric oxide level in male wistar rats. *Int. J. Pharmacol.*, 4: 487-491.
- Amoros, M., B. Fauconnier and R.L. Girre, 1987. *In vitro* antiviral activity of a saponin from *Anagallis arvensis*, Primulaceae, against herpes simplex virus and poliovirus. *Antiviral Res.*, 8: 13-25.
- Amutha, M., R. Arunachalam, M. Umamaheswari, A. Usharamalakshmi, S. Ramakrishnan and G. Annadurai, 2010. Medicinal use of *Camellia sinensis* on lactose intolerance. *J. Biol. Sci.*, 10: 112-116.
- Anam, K., R.M. Widharna and D. Kusrini, 2009.  $\alpha$ -Glucosidase inhibitor activity of *Terminalia* species. *Int. J. Pharmacol.*, 5: 277-280.
- Andersson, S., 2010. General polyhedra, virus structure and mutation. *J. Solid*, 225: 309-312.
- Anitha, S. and B.D. Ranjitha Kumari, 2006. *In vitro* flowering in *Rauvolfia tetraphylla* L. *Pak. J. Biol. Sci.*, 9: 422-424.
- Ansari, J.A. and N.N. Inamdar, 2010. The promise of traditional medicines. *Int. J. Pharmacol.*, 6: 808-812.
- Ansari, J.A., S. Ali and M.A. Ansari, 2011. A brief focus on hepatoprotective leads from herbal origin. *Int. J. Pharmacol.*, 7: 212-216.
- Appel, N., T. Schaller, F. Penin and R. Bartenschlager, 2006. From structure to function: New insights into hepatitis C virus RNA replication. *J. Biol. Chem.*, 281: 9833-9836.
- Appidi, J.R., D.S. Grierson and A.J. Afolayan, 2008. Ethnobotanical study of plants used for the treatment of diarrhoea in the Eastern Cape, South Africa. *Pak. J. Biol. Sci.*, 11: 1961-1963.
- Arshad, M. and Q.A. Khan, 2000. Ethnobotanical study of some medicinal plants of Rawal town. *Pak. J. Biol. Sci.*, 3: 1245-1246.
- Asano, J., K. Chiba, M. Tada and T. Yoshii, 1996. Antiviral activity of lignans and their glycosides from *Justicia procumbens*. *Phytochemistry*, 42: 713-717.
- Badam, L., 1997. *In vitro* antiviral activity of indigenous glycyrrhizin, licorice and glycyrrhizic acid (Sigma) on Japanese encephalitis virus. *J. Commun. Dis.*, 29: 91-99.
- Badam, L., S.S. Bedekar, K.B. Sonawane and S.P. Joshi, 2002. *In vitro* antiviral activity of bael (*Aegle marmelos* Corr) upon human coxsackieviruses B1-B6. *J. Commun. Dis.*, 34: 88-99.
- Bakar, M.F.A., A.H. The, A. Rahmat, F. Othman, N. Hashim and S. Fakurazi, 2006. Antiproliferative properties and antioxidant activity of various types of *Strobilanthes crispus* tea. *Int. J. Cancer Res.*, 2: 152-158.
- Banerjee, A., A. Sinhababu, R.K. Kar and S. Mandal, 2004. Micromorphological studies of four fuel wood yielding tropical leguminous plants. *Pak. J. Biol. Sci.*, 7: 100-104.

- Barakat, A.B., S.A. Shoman, N. Dina and O.R. Alfarouk, 2010. Antiviral activity and mode of action of *Dianthus caryophyllus* L. and *Lupinus termes* L. seed extracts against in vitro herpes simplex and hepatitis A viruses infection. *J. Microbiol. Antimicrob.*, 2: 28-29.
- Barquero, A.A., L.E. Alche and C.E. Coto, 2004. Block of vesicular stomatitis virus endocytic and exocytic pathways by 1-cinnamoyl-3, 11-dihydroxymeliacarpin, a tetranortriterpenoid of natural origin. *J. Gen. Virol.*, 85: 483-493.
- Bedows, E. and G.M. Hatfield, 1982. An investigation of the antiviral activity of *Podophyllum peltatum*. *J. Nat. Prod.*, 45: 725-729.
- Behbahani, M., 2009. Anti-viral activity of the methanolic leaf extract of an Iranian medicinal plant *Hyssopus officinalis* against herpes simplex virus. *J. Med. Plants Res.*, 3: 1118-1125.
- Bing, F.H., J. Liu, Z. Li, G.B. Zhang, Y.F. Liao, J. Li and C.Y. Dong, 2009. Anti-influenza-virus activity of total alkaloids from *Commelina communis* L. *Arch. Virol.*, 154: 1837-1840.
- Binns, S.E., J. Hudson, S. Merali and J.T. Arnason, 2002. Antiviral activity of characterized extracts from *Echinacea* spp. (Heliantheae: Asteraceae) against *Herpes simplex* virus (HSV-I). *Planta Med.*, 68: 780-783.
- Biron, K.K., 2006. Antiviral drugs for cytomegalovirus diseases. *Antiviral Res.*, 71: 154-163.
- Brandao, G.C., E.G. Kroon, J.R. dos Santos, J.R. Stehmann, J.A. Lombardi and A.B. de Oliveira, 2011. Antiviral activity of plants occurring in the State of Minas Gerais (Brazil): Part III. *J. Chem. Pharm. Res.*, 3: 223-236.
- Breitbart, M. and F. Rohwer, 2005. Here a virus, there a virus, everywhere the same virus? *Trends Micobiol.*, 13: 278-284.
- Briones, M.S., C.W. Dobard and S.A. Chow, 2010. Role of human immunodeficiency virus type 1 integrase in uncoating of the viral core. *J. Virol.*, 84: 5181-5190.
- Bunyapraphatsara, N., S. Dechsree, C. Yoosook, A. Herunsalee and Y. Panpisutchai, 2000. Anti-herpes simplex virus component isolated from *Maclura cochinchinensis*. *Phytomedicine*, 6: 421-424.
- Burlone, M.E. and A. Budkowska, 2009. Hepatitis C virus cell entry: Role of lipoproteins and cellular receptors. *J. Gen. Virol.*, 90: 1055-1070.
- Cesar, G.Z., M.G. Alfonso, M.M. Marius, E.M. Elizabeth and C.B. Angel *et al.*, 2011. Inhibition of HIV-1 reverse transcriptase, toxicological and chemical profile of *Calophyllum brasiliense* extracts from Chiapas, Mexico. *Fitoterapia*, 82: 1027-1034.
- Chakraborty, D. and S. Chakraborti, 2010. Bioassay-guided isolation and identification of antibacterial and antifungal component from methanolic extract of green tea leaves (*Camellia sinensis*). *Res. J. Phytochem.*, 4: 78-86.
- Chen, J.L., P. Blanc, C.A. Stoddart, M. Bogan and E.J. Rozhon *et al.*, 1998. New iridoids from the medicinal plant *Barleria prionitis* with potent activity against respiratory syncytial virus. *J. Nat. Prod.*, 61: 1295-1297.
- Cheng, H.Y., C.M. Yang, T.C. Lin, L.T. Lin, L.C. Chiang and C.C. Lin, 2011. Excoecarianin, isolated from phyllanthus *Urinaria linnea*, inhibits herpes simplex virus type 2 infection through inactivation of viral particles. *Evidence Based Compl. Alternative Med.*, 10.1093/ecam/nep157
- Cheng, H.Y., T.C. Lin, K. Ishimaru, C.M. Yang, K.C. Wang and C.C. Lin, 2003. *In vitro* antiviral activity of prodelphinidin B-2 3,3-di-O-gallate from *Myrica rubra*. *Planta Med.*, 69: 953-960.
- Chiang, L.C., W. Chiang, M.C. Liu and C.C. Liu, 2003. *In vitro* antiviral activities of *Caesalpinia pulcherrima* and its related flavonoids. *J. Antimicrob. Chemother.*, 50: 194-198.

- Corbett, E.L., C.J. Watt, N. Walker, D. Maher, B.G. Williams, M.C. Raviglione and C. Dye, 2003. The growing burden of tuberculosis: Global trends and interactions with the HIV epidemic. *Arch. Intern. Med.*, 163: 1009-1021.
- Craig, M.I., F. Benencia and F.C. Coulombe, 2001. Antiviral activity of an acidic polysaccharides fraction extracted from *Cedrela tubiflora* leaves Meliaceae plant. *Fitoterapia*, 72: 113-119.
- Czeizel, E. Andrew, Puho, Erzsebe, Acs, Nandor, Banhidy and Ferenc, 2006. A population based case-control study of oral moroxydine, an antiviral agent treatment during pregnancy. *Int. J. Pharmacol.*, 2: 188-192.
- Davison, A.J., R. Eberle, B. Ehlers, G.S. Hayward and D.J. McGeoch *et al.*, 2009. The order *herpesvirales*. *Arch. Virol.*, 154: 171-177.
- Del Barrio, G. and F. Parra, 2000. Evaluation of the antiviral activity of an aqueous extract from *Phyllanthus orbicularis*. *J. Ethnopharmacol.*, 72: 317-322.
- Denyer, C.V., P. Jackson, D.M. Loakes, M.R. Ellis and D.A. Young, 1994. Isolation of antirhinoviral sesquiterpenes from ginger (*Zingiber officinale*). *J. Nat. Prod.*, 57: 658-662.
- Duarte, M.E.R., D.G. Noseda, M.D. Noseda, S. Tulio, C.A. Pujol and E.B. Damonte, 2001. Inhibitory effect of sulfated galactans from the marine alga *Bostrychia montagnei* on herpes simplex virus replication *in vitro*. *Phytomedicine*, 8: 53-58.
- Dupre, J. and M.A. O'Malley, 2009. Varieties of living things: Life at the intersection of lineage and metabolism. *Philos. Theor. Biol.*, 1: 1-25.
- Edziri, H.L., G. Laurent, A. Mahjoub and M. Mastouri, 2011. Antiviral activity of *Conyza canadensis* (L.) Cronquist extracts grown in Tunisia. *Afr. J. Biotechnol.*, 10: 9097-9100.
- Ehrhardt, C., E.R. Hrincius, V. Korte, I. Mazur and K. Droebner *et al.*, 2007. A polyphenol rich plant extract, CYSTUS052, exerts anti influenza virus activity in cell culture without toxic side effects or the tendency to induce viral resistance. *Antiviral Res.*, 76: 38-47.
- Elkhalifa, K.F., S. Ibrahim and A. Hasan, 2005. Variations in tannin's contents of *Acacia nilotica* (L.) Willd. ex Del. in the Sudan. *Pak. J. Biol. Sci.*, 8: 1021-1024.
- Emtehani, M.H. and M. Tabari, 2007. *Acacia nilotica* and *Medicago sativa*, suitable plants for agro-forestry in southern coasts of Iran. *Pak. J. Biol. Sci.*, 10: 1713-1717.
- Erdelmeier, C.A., J. Cinatl, H. Rabenau, H.W. Doerr, A. Biber and E. Koch, 1996. Antiviral and antiphlogistic activities of *Hamamelis virginiana* bark. *Planta Med.*, 62: 241-245.
- Esquenazi, D., M.D. Wigg, M.M. Miranda, H.M. Rodrigues and C.S. Alviano *et al.*, 2002. Antimicrobial and antiviral activities of polyphenolics from *Cocos nucifera* Linn. (Palmae husk fiber extract). *Res. Microbiol.*, 153: 647-652.
- Falsey A.R., P.A. Hennessey, M.A. Formica, C. Cox and E.E. Walsh, 2005. Respiratory syncytial virus infection in elderly and high-risk adults. *N. Engl. J. Med.*, 352: 1749-1759.
- Fang, E.F., P. Lin, J.H. Wong, S.W. Tsao and T.B. Ng, 2010. A lectin with anti-HIV-1 reverse transcriptase, antitumor and nitric oxide inducing activities from seeds of *Phaseolus vulgaris* cv. extralong autumn purple bean. *J. Agric. Food Chem.*, 58: 2221-2229.
- Fard, M.H., S.L. Bodhankar and M. Dikshit, 2008. Cardioprotective activity of fruit of *Lagenaria siceraria* (Molina) standley on doxorubicin induced cardiotoxicity in rats. *Int. J. Pharmacol.*, 4: 466-471.
- Farooq, K., Abdul Hameed, T. Javed, I. Ullah, A.W. Khan and H. Khan, 2006. Comparative immunological response of commercial oil based and liposomal vaccines of avian influenza H7. *Pak. J. Biol. Sci.*, 9: 2402-2410.
- Firth, C., A. Kitchen, B. Shapiro, M.A. Suchard, E.C. Holmes and A. Rambaut, 2010. Using time-structured data to estimate evolutionary rates of double-stranded DNA viruses. *Mol. Biol. Evol.*, 27: 2088-2051.

- Fokunang, C.N., J. Hitchcock, F. Spence, E.A. Tembe-Fokunang, J. Burkhardt, L. Levy and C. George, 2006. An overview of mitochondrial toxicity of nucleoside reverse transcriptase inhibitors associated with HIV therapy. *Int. J. Pharmacol.*, 2: 152-162.
- Fuller, R.W., H.R. Bokesch, K.R. Gustafson, T.C. McKee and J.H. Cardellina *et al.*, 1994. HIV-inhibitory coumarins from latex of the tropical rainforest tree *Calophyllum teysmannii* var. *inophylloide*. *Bioorgan. Med. Chem. Lett.*, 4: 1961-1964.
- Gao, W. and J. Hu, 2007. Formation of hepatitis B virus covalently closed circular DNA: Removal of genome-linked protein. *J. Virol.*, 81: 6164-6174.
- Garcia, G., L. Cavallaro, A. Broussalis, G. Ferraro and V. Martino *et al.*, 1995. Antiviral activity of *Achyrocline flaccida* Wein DC aqueous extract. *Phytother. Res.*, 9: 251-254.
- Geng, Y., P. Dalhaimer, S. Cai, R. Tsai, M. Tewari, T. Minko and D.E. Discher, 2007. Shape effects of filaments versus spherical particles in flow and drug delivery. *Nat. Nanotechnol.*, 2: 249-255.
- Ghosh, S.C., S. Samsuzzaman, R. Sen, M.A.H. Talukder, S.M.A.H.M. Kamal and A.K. Saha, 2004. Effect of tree species and environment on the performance of wheat-T.aman rice and boro rice-t.aman rice cropping pattern. *Pak. J. Biol. Sci.*, 7: 670-673.
- Gorbalenya, A.E., L. Enjuanes, J. Ziebuhr and E.J. Snijder, 2006. Nidovirales: Evolving the largest RNA virus genome. *Virus Res.*, 117: 17-37.
- Gupta, M., B.P. Shaw and A. Mukherjee, 2008a. Evaluation of antipyretic effect of a traditional polyherbal preparation: A double-blind, randomized clinical trial. *Int. J. Pharmacol.*, 4: 190-195.
- Gupta, M., B.P. Shaw and A. Mukherjee, 2008b. Studies on antipyretic-analgesic and ulcerogenic activity of polyherbal preparation in rats and mice. *Int. J. Pharmacol.*, 4: 88-94.
- Hall, H.I., R. Song, P. Rhodes, J. Prejean and Q. An *et al.*, 2008. Estimation of HIV incidence in the United States. *JAMA*, 300: 520-529.
- Harvey, R.A., P.C. Champe, B.D. Fisher and W.A. Strohl, 2006. *Lippincott's Illustrated Reviews: Microbiology*. 2nd Edn., Lippincott Williams and Wilkins, Hagerstown, MD., ISBN: 9780781782159, pp: 354-366.
- Hassan, S.W., F.L. Bilbis, M.J. Ladan, R.A. Umar and S.M. Dangoggo *et al.*, 2006. Evaluation of antifungal activity and phytochemical analysis of leaves, roots and stem barks extracts of *Calotropis procera* (Asclepiadaceae). *Pak. J. Biol. Sci.*, 9: 2624-2629.
- Hassani, A.S., N. Amirmozafari, N. Ordouzadeh, K. Hamdi, R. Nazari and A. Ghaemi, 2008. Volatile components of *Camellia sinensis* inhibit growth and biofilm formation of oral strepto. *Pak. J. Biol. Sci.*, 11: 1336-1341.
- Heneidy, S.Z. and L.M. Bidak, 2004. Potential uses of plant species of the coastal mediterranean region, Egypt. *Pak. J. Biol. Sci.*, 7: 1010-1023.
- Hussein, E.A., A.M. Taj-Eldeen, A.S. Al-Zubairi, A.S. Elhakimi and A.R. Al-Dubaie, 2010. Phytochemical screening, total phenolics and antioxidant and antibacterial activities of callus from *Brassica nigra* L. hypocotyl explants. *Int. J. Pharmacol.*, 6: 464-471.
- Hussein, G., H. Miyashiro, N. Nakamura, M. Hattori, N. Kakiuchi and K. Shimotohno, 2000. Inhibitory effects of sudanese medicinal plant extracts on hepatitis C virus (HCV) protease. *Phytother. Res.*, 14: 510-516.
- Ilboudo, D., D. Karou, W.M.C. Nadembega, A. Savadogo and O.D.S. Pignatelli *et al.*, 2007. Prevalence of human herpes virus-8 and hepatitis B virus among HIV seropositive pregnant women enrolled in the mother-to-child HIV transmission prevention program at saint Camille medical centre in Burkina Faso. *Pak. J. Biol. Sci.*, 10: 2831-2837.

- Ishtiaq, M., W. Hanif, M.A. Khan, M. Ashraf and A.M. Butt, 2007. An ethnomedicinal survey and documentation of important medicinal folklore food phytonims of flora of Samahni valley (Azad Kashmir) Pakistan. Pak. J. Biol. Sci., 10: 2241-2256.
- Jaffary, F., V. Musini, M.A. Nilforoushzadeh and K. Bassett, 2007. Systematic review of imiquimod for the treatment of external genital wart. Int. J. Pharmacol., 3: 1-10.
- Javed, T., U.A. Ashfaq, S. Riaz, S. Rehman and S. Riazuddin, 2011. *In-vitro* antiviral activity of *Solanum nigrum* against hepatitis C virus. Virol. J., Vol. 8, 10.1186/1743-422X-8-26
- Joseph, B. and S.J. Raj, 2011. An overview: Pharmacognostic properties of *Phyllanthus amarus* Linn. Int. J. Pharmacol., 7: 40-45.
- Jun, T., Z. Liancai and W. Bochu, 2007. Effect of ginsenosides on malondialdehyde, nitric oxide and endothelin-1 production in vascular endothelial cells suffering from lipid peroxidation injury. Int. J. Pharmacol., 3: 101-105.
- Kageyama, S., M. Kurokawa, H. Sato, T. Yukawa, H. Ohyama, T. Kurimura, T. Namba and K. Shiraki, 1996. Potent activity of the extract of *Geum japonicum* Thunb. For the prophylaxis of cytomegalovirus infection in AIDS patients. Int. Conf. AIDS, 11: 65-65.
- Kagone, T.S., H. Hien, N. Meda, P.S. Diagbouga and A. Sawadogo *et al.*, 2011. Characterization of HIV-1 genotypes and antiretroviral drug-resistance mutations among patients in Burkina Faso. Pak. J. Biol. Sci., 14: 392-398.
- Kalus, U., A. Grigorov, O. Kadecki, J.P. Jansen, H. Kiesewetter and H. Radtke, 2009. *Cistus incanus* (CYSTUS052) for treating patients with infection of the upper respiratory tract: A prospective, randomised, placebo-controlled clinical study. Antiviral Res., 84: 267-271.
- Kan, A., B. Ozcelik and M. Kartal, 2009. *In vitro* antiviral activities under cytotoxic doses against herpes simplex type-1 and parainfluenza-3 viruses of *Cicer arietinum* L. (chickpea). Afr. J. Pharm. Pharmacol., 3: 627-631.
- Karou, S.D., T. Tchaondo, D.P. Ilboudo and J. Simpore, 2011. Sub-saharan rubiaceae: A review of their traditional uses, phytochemistry and biological activities. Pak. J. Biol. Sci., 14: 149-169.
- Kaur C.D. and S. Saraf, 2011. Photochemoprotective activity of alcoholic extract of *Camellia sinensis*. Int. J. Pharmacol., 7: 400-404.
- Kernan, M.R., A. Amarquaye, J.L. Chen, J. Chan and D.F. Sesin *et al.*, 1998. Antiviral phenylpropanoid glycosides from the medicinal plant *Markhamia lutea*. J. Nat. Prod., 61: 564-570.
- Kernan, M.R., A. Sendl, J.L. Chen, S.D. Jolad and P. Blanc *et al.*, 1997. Two new lignans with activity against influenza virus from the medicinal plant *Rhinacanthus nasutus*. J. Natural Product, 60: 635-637.
- Kim, H.K., H.K. Lee, C.G. Shin and H. Huh, 1999. HIV integrase inhibitory activity of *Agastache rugosa*. Arch. Pharm. Res., 22: 520-523.
- Krishnan, N., S. Ramanathan, S. Sasidharan, V. Murugaiyah and S.M. Mansor, 2010. Antimicrobial activity evaluation of cassia spectabilis leaf extracts. Int. J. Pharmacol., 6: 510-514.
- Kumar, G., G. Sharmila Banu, A.G. Murugesan and M. Rajasekara Pandian, 2007. Preliminary toxicity and phytochemical studies of aqueous bark extract of *Helicteres isora* L. Int. J. Pharmacol., 3: 96-100.
- Kurokawa, M., P. Basnet, M. Ohsugi, T. Hozumi and S. Kadota *et al.*, 1999. Anti-herpes simplex virus activity of moronic acid purified from *Rhus javanica* *in vitro* and *in vivo*. J. Pharmacol. Exp. Ther., 289: 72-78.

- Ky, J.M.K., P. Zerbo, C. Gnoula, J. Simpore, J.B. Nikiema and J. Millogo-Rasolodimby, 2009. Medicinal plants used in traditional medicine in the centre east region of burkina faso. *Pak. J. Biol. Sci.*, 12: 1287-1298.
- Laure, F., P. Raharivelomanana, J.F. Butaud, J.P. Bianchini and E.M. Gaydou, 2008. Screening of anti-HIV-1 inophyllums by HPLC-DAD of *Calophyllum inophyllum* leaf extracts from French Polynesia Islands. *Anal. Chim. Acta*, 624: 147-153.
- Lazzari I.S. and K. Stohr, 2004. Avian influenza and influenza pandemics. *Bull. World Health Organ.*, 82: 242-242.
- Lee, D., K.H. Boo, J.K. Woo, F. Duan and K.H. Lee *et al.*, 2011. Anti-bacterial and anti-viral activities of extracts from *Terminalia chebula* Bark. *J. Korean Soc. Appl. Biol. Chem.*, 54: 295-298.
- Lee-Huang, S., P.L. Huang, H.F. Kung, B.Q. Li and P.L. Huang *et al.*, 1991. TAP 29: An anti-human immunodeficiency virus protein from *Trichosanthes kirilowii* that is nontoxic to intact cells. *PNAS*, 88: 6570-6574.
- Lee-Huang, S., P.L. Huang, P.L. Huang, A.S. Bourinbaiar, H.C. Chens and H.F. Kungii, 1995. Inhibition of the integrase of human immunodeficiency virus (HIV) type 1 by anti-HIV plant proteins MAP30 and GAP31. *Proc. Natl. Acad. Sci. USA*, 92: 8818-8822.
- Li, Y., P.P. But and V.E. Ooi, 2005. Antiviral activity and mode of action of caffeoylquinic acids from *Schefflera heptaphylla* (L.) Frodin. *Antiviral Res.*, 68: 1-9.
- Lin, Y.M., H. Anderson, M.T. Flavin, Y.H. Pai and E. Mata-Greenwood *et al.*, 1997. *In vitro* anti-HIV activity of biflavonoids isolated from *Rhus succedanea* and *Garcinia multiflora*. *J. Nat. Prod.*, 60: 884-888.
- Lowen, A.C., S. Mubareka, J. Steel and P. Palese, 2007. Influenza virus transmission is dependent on relative humidity and temperature. *PLoS Pathog.*, 3: 1470-1476.
- Lu, S.Y., Y.J. Qiao, P.G. Xiao and X.H. Tan, 2005. Identification of antiviral activity of *Toddalia asiatica* (L.) against influenza type A virus. *Zhongguo Zhong Yao Za Zhi*, 30: 998-1001.
- Ma, S.C., P.P. But, V.E. Ooi, Y.H. He, S.H. Lee, S.F. Lee and R.C. Lin, 2001. Antiviral amentoflavone from *Selaginella sinensis*. *Biol Pharm Bull.*, 24: 311-312.
- Mahalakshmi, R., P. Rajesh, N. Ramesh, V. Balasubramanian and V.R. Kannan, 2010. Hepatoprotective activity on *Vitex negundo* Linn. (verbenaceae) by using wistar albino rats in ibuprofen induced model. *Int. J. Pharmacol.*, 6: 658-663.
- Mahzounieh, M., E. Moghtadaei and T. Zahraei Salehi, 2006. Detection of calicivirus genome in calves using Ni/E3 primers in Shahrekord area, Iran. *Pak. J. Biol. Sci.*, 9: 227-230.
- Makambila-Koubemba, M.C., B. Mbatchi, D. Ardid, A. Gelot and C. Henrion *et al.*, 2011. Pharmacological studies of ten medicinal plants used for analgesic purposes in Congo Brassaville. *Int. J. Pharm.*, 7: 608-615.
- Marchetti, M., S. Pisani, V. Pietropaolo, L. Seganti, R. Nicoletti, A. Degener and N. Orsi, 1996. Antiviral effect of a polysaccharide from *Sclerotium glucanicum* towards herpes simplex virus type 1 infection. *Planta Med.*, 62: 303-307.
- Mbaya, A.W. and U.I. Ibrahim, 2011. *In vivo* and *in vitro* activities of medicinal plants on haemic and humoral trypanosomes: A review. *Int. J. Pharmacol.*, 7: 1-11.
- Meragelman, K.M., T.C. McKee and M.R. Boyd, 2001. Anti-HIV prenylated flavonoids from *Monotes africanus*. *J. Nat. Prod.*, 64: 546-548.
- Meyer, J.J.M., A.J. Afolayan, M.B. Taylor and D. Erasmus, 1997. Antiviral activity of galangin isolated from the aerial parts of *Helichrysum aureonitens*. *J. Ethnopharmacol.*, 56: 165-169.

- Miladi, S., N. Abid, C. Debarnot, M. Damak, B. Canard, M. Aouni and B. Selmi, 2011. *In vitro* antiviral activities of extracts derived from *Daucus maritimus* seeds. Natural Product Res., 10.1080/14786419.2010.550263
- Mohamed, A.M. and N.S. Metwally, 2009. Antiaflatoxigenic activities of some plant aqueous extracts against aflatoxin-b1 induced renal and cardiac damage. *J. Pharmacol. Toxicol.*, 4: 1-16.
- Mohapatra, S.S. and S. Boyapalle, 2008. Epidemiologic, experimental and clinical links between respiratory syncytial virus infection and asthma. *Clin. Microbiol. Rev.*, 21: 495-504.
- Momtaz, S. and M. Abdollahi, 2010. An update on pharmacology of *Satureja* species: From antioxidant, antimicrobial, antidiabetes and anti-hyperlipidemic to reproductive stimulation. *Int. J. Pharmacol.*, 6: 346-353.
- Moundipa, P.F., S. Ngouela, G.A. Tchamba, N.F. Njayou, P.D.D. Chuisseau, Z.F. and E. Tsamo, 2007. Antihepatotoxic activity of *Xylopia phloiodora* extracts on some experimental models of liver injury in rats. *Int. J. Pharmacol.*, 3: 74-79.
- Nagai, T., Y. Suzuki, T. Tomimori and H. Yamada, 1995. Antiviral activity of plant flavonoid, 5,7,4'-trihydroxy-8-methoxyflavone, from the roots of *Scutellaria baicalensis* against influenza A (H3N2) and B viruses. *Biol. Pharm. Bull.*, 18: 295-299.
- Ndjonka, D., C. Agyare, K. Luersen, A. Hensel and E. Liebau, 2010. *In vitro* anti-leishmanial activity of traditional medicinal plants from cameroon and Ghana. *Int. J. Pharmacol.*, 6: 863-871.
- Ng, T.B. and H. Wang, 2001. Panaxagin, a new protein from Chinese ginseng possesses anti-fungal, anti-viral, translation-inhibiting and ribonuclease activities. *Life Sci.*, 68: 739-749.
- Obaid, A.Y., O.A. Abu-Zinadah and H.K. Hussein, 2011. The beneficial effects of green tea extract and its main derivatives in repairing skin burns of rabbit. *Int. J. Biol. Chem.*, 5: 103-115.
- Oguntibeju, O.O., W.M.J Van den Heever and F.E. Van Schalkwyk, 2007. Supplementation effect on body weight and BMI of HIV-positive/AIDS patients. *Int. J. Pharmacol.*, 3: 120-122.
- Oh, C., J. Price, M.A. Brindley, M.P. Widrlechner and L. Qu *et al.*, 2011. Inhibition of HIV-1 infection by aqueous extracts of *Prunella vulgaris* L. *Virol. J.*, 8: 188-188.
- Ojha, S.K., M. Nandave, S. Arora, R. Narang, A.K. Dinda and D.S. Arya, 2008. Chronic administration of *Tribulus terrestris* Linn. extract improves cardiac function and attenuates myocardial infarction in rats. *Int. J. Pharmacol.*, 4: 1-10.
- Oldstone, M.B., 1993. Rous-whipple award lecture. viruses and diseases of the twenty-first century. *Am. J. Pathol.*, 143: 1241-1249.
- Onwuliri, V.A. and G.E. Anekwe, 2001. Amino acids and other biochemical components of *Ricinus communis* (Variety Minor), an anti-conceptive seed. *Pak. J. Biol. Sci.*, 4: 866-868.
- Ooi, L.S., W.S. Ho, K.L. Ngai, L. Tian, P.K. Chan, S.S. Sun and V.E. Ooi, 2010. *Narcissus tazetta* lectin shows strong inhibitory effects against respiratory syncytial virus, influenza A (H1N1, H3N2, H5N1) and B viruses. *J. Biosci.*, 35: 95-103.
- Ooi, L.S.M., S.S.M. Sun and V.E.C. Ooi, 2004. Purification and characterization of a new antiviral protein from the leaves of *Pandanus amaryllifolius* (Pandanaceae). *Int. J. Biochem. Cell. Biol.*, 36: 1440-1446.
- Parida, M.M., C. Upadhyay, G. Pandya and A.M. Jana, 2002. Inhibitory potential of neem (*Azadirachta indica* Juss) leaves on Dengue virus type-2 replication. *J. Ethnopharmacol.*, 79: 273-278.
- Patton, G.C., C. Coffey, S.M. Sawyer, R.M. Viner and D.M. Haller *et al.*, 2009. Global patterns of mortality in young people: A systematic analysis of population health data. *The Lancet*, 374: 881-892.

- Pervez, K., 2000a. Effect of combination chemotherapy on hepatitis C virus in hepatic patients. *Pak. J. Biol. Sci.*, 3: 969-970.
- Pervez, K., 2000b. Hepatitis C virus core protein expression for enzyme linked immunosorbant assay. *Pak. J. Biol. Sci.*, 3: 1096-1097.
- Pichereau, C., K. Desseaux, A. Janin, C. Scieux and R.P. de Latour *et al.*, 2011. The complex relationship between human herpesvirus 6 and acute graft-versus-host disease. *Biol. Blood Marrow Transplant.*, 10.1016/j.bbmt.2011.07.018
- Pichlmair, A., O. Schulz, C.P. Tan, T.I. Naslund, P. Liljestrom, F. Weber and C. Reis e Sousa, 2006. RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-phosphates. *Science*, 314: 997-1001.
- Pleschka, S., M. Stein, R. Schoop and J.B. Hudson, 2009. Anti-viral properties and mode of action of standardized Echinacea purpurea extract against highly pathogenic avian Influenza virus (H5N1, H7N7) and swine-origin H1N1 (S-OIV). *Virol. J.*, 6: 197-197.
- Praseno, S.S. and N. Rintiswati, 1997. Antiviral activity of *Momordica charantia*: A preliminary study on in vitro anti-herpes simplex virus. *Periodic Med. Sci.*, 29: 121-123.
- Premanathan, M., K. Kathiresan, N. Yamamoto and H. Nakashima, 1999a. *In vitro* anti-human immunodeficiency virus activity of polysaccharide from *Rhizophora mucronata* Poir. *Biosci. Biotechnol. Biochem.*, 63: 1187-1191.
- Premanathan, M., R.A.H. Izumi, K. Kathiresan, M. Nakano, N. Yamamoto and H. Nakashima, 1999b. Anti-viral properties of a mangrove plant, *Rhizophora apiculata* Blume, against human immunodeficiency virus. *Antiviral Res.*, 15: 113-122.
- Pu, J.X., L.M. Yang, W.L. Xiaoa, R.T. Li and C. Lei *et al.*, 2008. Compounds from *Kadsura heteroclite* and related anti-HIV activity. *Phytochemistry*, 69: 1266-1272.
- Qureshi, S.A., A. Nawaz, S.K. Udani and B. Azmi, 2009. Hypoglycaemic and Hypolipidemic activities of *Rauwolfia serpentina* in Alloxan-induce diabetic rats. *Int. J. Pharmacol.*, 5: 323-326.
- Qureshi, S.J., S. Bano, T. Mohammad and M.A. Khan, 2001. Medicinal potential of poisonous plants of tehsil Kahuta from district Rawalpindi, Pakistan. *Pak. J. Biol. Sci.*, 4: 331-332.
- Rahman, S., M.M. Akbor, A. Howlader and A. Jabbar, 2009. Antimicrobial and cytotoxic activity of the alkaloids of amlaki (*Emblica officinalis*). *Pak. J. Biol. Sci.*, 12: 1152-1155.
- Raja, K.S. Q. Wang and M.G. Finn, 2003. Icosahedral virus particles as polyvalent carbohydrate display platforms. *Chembiochem*, 42: 1348-1351.
- Rajbhandari, M., R. Mente, P.K. Jha, R.P. Chaudhary and S. Bhattarai *et al.*, 2009. Antiviral activity of some plants used in Nepalese traditional medicine. *Evidence-Based Complement. Alternat. Med.*, 63: 517-522.
- Rehman, S., U.A. Ashfaq, S. Riaz, T. Javed and S. Riazuddin, 2011. Antiviral activity of *Acacia nilotica* against Hepatitis C Virus in liver infected cells. *Virol. J.*, 86: 220-220.
- Reichling, J., A. Neuner, M. Sharaf, M. Harkenthal and P. Schnitzler, 2009. Antiviral activity of *Rhus aromatica* (fragrant sumac) extract against two types of herpes simplex viruses in cell culture. *Pharmazie*, 64: 538-541.
- Reutrakul, V., N. Ningnuek, M. Pohmakotr, C. Yoosook and C. Napaswad *et al.*, 2007. Anti HIV-1 flavonoid glycosides from *Ochna integerrima*. *Planta Med.*, 73: 683-688.
- Richardson, T.H., 2010. Cannabis use and mental health: A review of recent epidemiological research. *Int. J. Pharmacol.*, 6: 796-807.

- Rocha Martins, L.R., M.A. Brenzan, C.V. Nakamura, B.P. Dias Filho, T.U. Nakamura, L.E. Ranieri Cortez and DA. Garcia Cortez, 2011. *In vitro* antiviral activity from *Acanthospermum australe* on herpesvirus and poliovirus. *Pharm Biol.*, 49: 26-31.
- Roja, G. and M.R. Heble, 1995. Castanospermine, an HIV inhibitor from tissue cultures of *Castanospermum australe*. *Phytotherapy Res.*, 9: 540-542.
- Roner, M.R., J. Sprayberry, M. Spinks and S. Dhanji, 2007. Antiviral activity obtained from aqueous extracts of the Chilean soapbark tree (*Quillaja saponaria* Molina). *J. Gen. Virol.*, 88: 275-285.
- Roshandel, G., S. Semnani, N. Abdolahi, A.A. Keshtkar and S. Besharat *et al.*, 2007. Prevalence of hepatitis D virus infection in HbsAg positive subjects in Iran. *Pak. J. Biol. Sci.*, 10: 1751-1754.
- Saeed, S.A. and A. Hussain, 2006. Anti-HIV drugs of the herbal type. DAWN Sci-tech World, <http://archives.dawn.com/dawnftp/72.249.57.55/dawnftp/weekly/science/archive/060603/science5.htm>.
- Sampathkumar, P., B. Dheeba, V. Vidhyasagar, T. Arulprakash and R. Vinothkannan, 2008. Potential antimicrobial activity of various extracts of *Bacopa monnieri* (Linn.). *Int. J. Pharmacol.*, 4: 230-232.
- Selisko, B., J.C. Guillemot, K. Alvarez and B. Canard, 2007. Opportunities in the development of anti-dengue drugs. Annual report, Scientific working group report on dengue, TDR.
- Semalty, M., A. Semalty, Geeta P. Joshi and M.S.M. Rawat, 2010. *In vivo* hair growth activity of herbal formulations. *Int. J. Pharmacol.*, 6: 53-57.
- Semple, S.J., S.F. Nobbs, S.M. Pyke, G.D. Reynolds and R.L. Flower, 1999. Antiviral flavonoid from *Pterocaulon sphacelatum*, an australian aboriginal medicine. *J. Ethnopharmacol.*, 68: 283-288.
- Sendl, A., J.L. Chen, S.D. Jolad, C. Stoddart and E. Rozhon *et al.*, 1996. Two new naphthoquinones with antiviral activity from *Rhinacanthus nasutus*. *J. Nat. Prod.*, 59: 808-811.
- Serkedjieva, J., 1996. A polyphenolic extract from *Geranium sanguineum* L. inhibits influenza virus protein expression. *Phytotherapy Res.*, 10: 441-443.
- Serkedjieva, J., N. Manolova, I. Zgorniak-Nowosielska, B. Zawilinska and J. Grzybek, 2006. Antiviral activity of the infusion (SHS-174) from flowers of *Sambucus nigra* L., aerial parts of *Hypericum perforatum* L. and roots of *Saponaria officinalis* L., against influenza and herpes simplex viruses. *Phytother. Res.*, 4: 97-100.
- Shalaby, E.A., S.M. Shanab and V. Singh, 2010. Salt stress enhancement of antioxidant and antiviral efficiency of *Spirulina platensis*. *J. Med. Plants Res.*, 4: 2622-2632.
- Shin, W.J., K.H. Lee, M.H. Park and B.L. Seong, 2010. Broad-spectrum antiviral effect of *Agrimonia pilosa* extract on influenza viruses. *Microbiol. Immunol.*, 54: 11-19.
- Shinde, S.L., S.B. Junne, S.S. Wadje and M.M.V. Baig, 2009. The diversity of antibacterial compounds of terminalia species (Combretaceae). *Pak. J. Biol. Sci.*, 12: 1483-1486.
- Shinwari, M.I. and M.A. Khan, 2000. Vegetation comparison of sacred, reserved and unreserved sites of Rumli village at Margalla Hills National Park, Islamabad. *Pak. J. Biol. Sci.*, 3: 1681-1683.
- Shirazi, M.U., B. Khanzada, S.M. Alam, R. Ansari, S.M. Mujtaba, M. Ali, M. Ali and M.A. Khan, 2001. Seasonal nutrient variations in two *Acacia* species growing under saline environment. *Pak. J. Biol. Sci.*, 4: 514-517.

- Shokrzadeh, M., A.G. Ebadi, S.S. Mirshafiee and M.I. Choudhary, 2006. Effect of the aqueous green leaf extract of green tea (*Camellia sinensis*) on glucose level of rat. Pak. J. Biol. Sci., 9: 2708-2711.
- Sluis-Cremer, N. and G. Tachedjian, 2008. Mechanisms of inhibition of HIV replication by nonnucleoside reverse transcriptase inhibitors. Virus Res., 134: 147-156.
- Sokmen, M., M. Angelova, E. Krumova, S. Pashova, S. Ivancheva, A. Sokmen and A. Serkedjieva, 2005. *In vitro* antioxidant activity of polyphenol extracts with antiviral properties from *Geranium sanguineum* L. Life Sci., 76: 2981-2993.
- Soleimanjahi, H., M. Jamalidoost, F. Fotouhi and Z. Meshkat, 2007. Amplification and cloning of herpes simplex virus type 2 glycoprotein G from an Iranian isolate. Pak. J. Biol. Sci., 10: 955-958.
- Song, J.H., K.J. Im, S.W. Chae and H.J. Choi, 2010. Prunus mume possess anti-human rhinovirus activit. J. Cosmet. Publ. Health, 6: 1-3.
- Song, J.M., K.H. Lee and B.L. Seong, 2005. Antiviral effect of catechins in green tea on influenza virus. Antiviral Res., 68: 66-74.
- Sonibare, M.A., T.O. Lawal and O.O. Ayodeji, 2011. Antimicrobial evaluation of plants commonly used in the management of psychosis opportunistic infections. Int. J. Pharmacol., 7: 492-497.
- Sotanaphun, U., V. Lipipun, R. Suttisri and R. Bavovada, 1999. A new antiviral and antimicrobial sesquiterpene from *Glyptopetalum sclerocarpum*. Planta Med., 65: 257-258.
- Subash, K.R., K.S. Ramesh, B.V. Charan, F. Britto, N.J. Rao and S. Vijaykumar, 2011. Study of hepatoprotective activity of *Solanum nigrum* and *Cichorium intybus*. Int. J. Pharmacol., 7: 504-509.
- Subbarao, K. and T. Joseph, 2007. Scientific barriers to developing vaccines against avian influenza viruses. Nat. Rev. Immunol., 7: 267-278.
- Sukandar, E.Y., H. Permana, I.K. Adnyana, J.I. Sigit, R.A. Ilyas, P. Hasimun and D. Mardiyah, 2010. Clinical study of turmeric (*Curcuma longa* L.) and garlic (*Allium sativum* L.) extracts as antihyperglycemic and antihyperlipidemic agent in type-2 diabetes-dyslipidemia patients. Int. J. Pharmacol., 6: 456-463.
- Sunday, O.A., A.B. Munir, O.O. Akeeb, A.A. Bolanle and S.O. Badaru, 2010. Antiviral effect of *Hibiscus sabdariffa* and *Celosia argentea* on measles virus. Afr. J. Microbiol. Res., 4: 293-296.
- Suttle, C.A., 2005. Viruses in the sea. Nature, 437: 356-361.
- Sydiskia, R.J., D.G. Owen, J.L. Lohr, K.H.A. Rosler and R.N. Blomster, 1991. Inactivation of enveloped viruses by anthraquinones extracted from plants. Antimicrob. Agents Chemother., 35: 2463-2466.
- Syed, G.H., Y. Amako and A. Siddiqui, 2010. Hepatitis C virus hijacks host lipid metabolism. Trends Endocrinol. Metab., 21: 33-40.
- Sylvestre, D.L., B.J. Clements and Y. Malibu, 2006. Cannabis use improves retention and virological outcomes in patients treated for hepatitis C. Eur. J. Gastroenterol. Hepatol., 18: 1057-1063.
- Tabassum, F., R. Khurshid and M.W. Akhtar, 2000. Identification of marker protein, in patient with chronic liver disease. Pak. J. Biol. Sci., 3: 1509-1510.
- Takechi M. and Tanaka Y., 1981. Purification and characterization of antiviral substance from the bud of *Syzygium aromaticum*. Planta Med., 42: 69-74.
- Tehranipour, M. and S. Ebrahimpour, 2009. Evaluating the effect of aquatic extract of *Cannabis sativa* seed on spatial memory consolidation in rats. J. Biol. Sciences, 9: 884-888.

- Thompson, W.W., D.K. Shay, E. Weintraub, L. Brammer, C.B. Bridges, N.J. Cox and K. Fukuda, 2004. Influenza-associated hospitalizations in the United States. *JAMA*, 292: 1333-1340.
- Trento, A., I. Casas, A. Calderon, M.L. Garcia-Garcia, C. Calvo, P. Perez-Brena and J.A. Melero, 2010. Ten years of global evolution of the human respiratory syncytial virus BA genotype with a 60-nucleotide duplication in the G protein gene. *J. Virol.*, 84: 7500-7512.
- Valcheva-Kuzmanova, S.V. and A. Belcheva, 2006. Current knowledge of *Aronia melanocarpa* as a medicinal plant. *Folia Med. (Plovdiv)*, 48: 11-17.
- Vimalanathan, S., L. Kang, V.T. Amiguet, J. Livesey, J.T. Arnason and J. Hudson, 2005. *Echinacea purpurea* aerial parts contain multiple antiviral compounds. *Pharm. Biol.*, 43: 740-745.
- Vimalanathan, S., S. Ignacimuthu and J.B. Hudson, 2009. Medicinal plants of Tamil Nadu (Southern India) are a rich source of antiviral activities. *Pharm. Biol.*, 47: 422-429.
- WHO, 2001. Herpes simplex virus type 2: Programmatic and research priorities in developing countries. Report of a WHO/UNAIDS/LSHTM Workshop | London, UK. <http://www.who.int/hiv/pub/sti/pub9/en/index.html>.
- WHO, 2011a. Hepatitis C. Fact sheet N°164. <http://www.who.int/mediacentre/factsheets/fs164/en/>.
- WHO, 2011b. World Health Statistics. WHO, Geneva, Switzerland.
- Wang, H.X. and T.B. Ng, 2001. Examination of lectins, polysaccharopeptide, polysaccharide, alkaloid, coumarin and trypsin inhibitors for inhibitory activity against human immunodeficiency virus reverse transcriptase and glycohydrolases. *Planta Med.*, 67: 669-672.
- Wapling, J., S. Srivastava, M. Shehu-Xhilaga and G. Tachedjian, 2007. Targeting human immunodeficiency virus type 1 assembly, maturation and budding. *Drug Target Insights*, 2: 159-182.
- Weber, N.D., A.D. Douglas, J.A. North, B.K. Murray, L.D. Lawson and B.G. Hughes, 1992. *In vitro* virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta Med.*, 58: 417-423.
- Wiart, C., K. Kumar, M.Y. Yusof, H. Hamimah and Z.M. Fauzi et al., 2005. Antiviral properties of ent-labdene diterpenes of *Andrographis paniculata* Nees, inhibitors of herpes simplex virus type 1. *Phytother. Res.*, 19: 1069-1070.
- Wirottesangthong, M., T. Nagai, H. Yamada, S. Amnuoypol and C. Mungmee, 2009. Effects of *Clinacanthus siamensis* leaf extract on influenza virus infection. *Microbiol. Immunol.*, 53: 66-74.
- Yang, C.M., H.Y. Cheng, T.C. Lin, L.C. Chiang and C.C. Lin, 2007. Hippomanin a from acetone extract of *Phyllanthus urinaria* inhibited HSV-2 but not HSV-1 infection *in vitro*. *Phytother. Res.*, 21: 1182-1186.
- Ye, X.Y. and T.B. Ng, 2001. Isolation of unguilin, a cyclophilin-like protein with anti-mitogenic, antiviral and antifungal activities, from black-eyed pea. *J. Protein Chem.*, 20: 353-359.
- Ye, X.Y., H.X. Wang and T.B. Ng, 2000. Structurally dissimilar proteins with antiviral and antifungal potency from cowpea (*Vigna unguiculata*) seeds. *Life Sci.*, 67: 3199-3207.
- Ye, X.Y., T.B. Ng, P.W.K. Tsang and J. Wang, 2001. Isolation of a homodimeric lectin with antifungal and antiviral activities from red kidney bean (*Phaseolus vulgaris*) seeds. *J. Protein Chem.*, 20: 367-375.
- Yeh, J.H., H.F. Chiu, J.S. Wang, J.K. Lee and T.C. Chou, 2010. Protective effect of baicalein extracted from *Scutellaria baicalensis* against lipopolysaccharide-induced glomerulonephritis in mice. *Int. J. Pharmacol.*, 6: 81-88.

- Yoshida, T., H. Ito, T. Hatano, M. Kurata and T. Nakanishi *et al.*, 1996. New hydrolyzable tannins, shephagenins A and B, from *Shepherdia argentea* as HIV-1 reverse transcriptase inhibitors. *Chem. Pharm. Bull. (Tokyo)*, 44: 1436-1439.
- Yucharoen, R., S. Anuchapreeda and Y. Tragoolpua, 2011. Anti-herpes simplex virus activity of extracts from the culinary herbs *Ocimum sanctum* L., *Ocimum basilicum* L. and *Ocimum americanum* L. *Afr. J. Biotechnol.*, 10: 860-866.
- Zakizad, M., F. Salmeh, T. Yaghoobi, M. Yaghoubian and M.B. Nesami *et al.*, 2009. Seroprevalence of hepatitis C infection and associated risk factors among addicted prisoners in Sari-Iran. *Pak. J. Biol. Sci.*, 12: 1012-1018.
- Zambon, M.C., 2001. The pathogenesis of influenza in humans. *Rev. Med. Virol.*, 11: 227-241.
- Zandi, K., M. Abbaszadeh, K. Sartavi and Z. Rastian, 2007. Antiviral activity of *Aloe vera* against herpes simplex virus type 2: An *in vitro* study. *Afr. J. Biotechnol.*, 6: 1770-1773.
- Zhang, J., B. Zhan, X. Yao, Y. Gao and J. Shong, 1995. Antiviral activity of tannin from the pericarp of *Punica granatum* L. against genital herpes virus *in vitro*. *Zhongguo Zhong Yao Za Zhi*, 20: 556-558.
- Zhang, Y., H. Zhu, G. Ye, C. Huang and Y. Yang *et al.*, 2006. Antiviral effects of sophoridine against coxsackievirus B3 and its pharmacokinetics in rats. *Life Sci.*, 78: 1998-2005.
- Zhong, Y., C. Zuo, F. Li, X. Ding and Q. Yao *et al.*, 1998. Chemical constituents of *Phyllanthus urinaria* L. and its antiviral activity against hepatitis B virus. *Zhongguo Zhong Yao Za Zhi*, 23: 363-364.
- Ziyaeyan, M., A. Japoni, M.H. Roostaei, S. Salehi and H. Soleimanjahi, 2007. A serological survey of herpes simplex virus type 1 and 2 Immunity in pregnant women at labor stage in Tehran, Iran. *Pak. J. Biol. Sci.*, 10: 148-151.