



Asian Journal of **Biochemistry**

ISSN 1815-9923



Academic
Journals Inc.

www.academicjournals.com

Enzymatic and Oven-drying Method of Processing Rubber Seeds for Animal Feed and the Evaluation of the Toxicity of Such Feed in Rats

P.N. Okafor and N.O. Anyanwu
Department of Biochemistry, Michael Okpara University of Agriculture,
Umudike, P.M.B 7267 Umuahia Abia State, Nigeria

Abstract: The hydrolytic activity of endogenous β -glucosidase of *Hevea brasiliensis* towards the cyanogenic glycosides of rubber seeds as a means of processing was investigated followed by the study of acute toxicity of the enzymatically and oven dried processed feeds from these seeds towards rats. Crushing/ grating of the rubber seeds and allowing interaction of the endogenous β -glucosidase with the cyanogenic glycosides content for a period of 60 min resulted in about 90-95% hydrolysis of the cyanogenic glycoside by the β -glucosidase. Oven drying of the crushed seeds after 60 min at a temperature of 60°C for 180 min gave rise to about 81% reduction in the total cyanogens content. Ingestion of feed compounded from the oven dried seed containing 9.25 mg CN⁻¹ kg⁻¹ did not produce any sign of acute toxic effects in rats after 72 h. Elevation in blood glucose and Thiocyanate was observed but the activities of aspartate and alanine aminotransferases and alkaline phosphatase did not show any significant change compared to those of control. The same was true of the total protein and serum albumin levels of the animals. The results are discussed from toxicological and feed value points of view.

Key words: Enzymatic, oven drying, processing, *H. brasiliensis*, toxicity, rats

Introduction

The importance of increased livestock production in developing countries like Nigeria cannot be over emphasized. The human need for foods with high biological values such as meat and eggs and the possibility of covering them in this part of the world has become so necessary with increasing population and disease conditions. In spite of the great demand for protein of high biological value for human consumption, pigs and poultry production in the tropics (developing countries) is scanty and inefficient as a rule (Preston, 1984).

The number of cattle, sheep, goats and chicken found in developing countries in 1991 far out numbered those in developed countries but the quantity of meat and milk produced in developed countries was far higher than that produced in developing countries (FAO, 1992). The underlying reason for this situation is the system of management and feeding strategies in the developing countries (Iyayi and Tawe, 1994).

The use of unconventional feeds has been advanced as one of the ways to improve animal production in the tropics (Iyayi and Tawe, 1994). Of particular interest among these unconventional feeds such as cassava peels, sorghum spent grains, wheat offals and others is the rubber seed meal.

Corresponding Author: Dr. P.N. Okafor, Department of Biochemistry, Michael Okpara University of Agriculture, Umudike, P.M.B 7267 Umuahia Abia State, Nigeria

That rubber seeds could be used for feeding animals including monogastric species have been reported (Ong and Yeong, 1977, Nwokolo, 1987; Ly *et al.*, 2001). That rubber seed cake has nutritive values and is being used in cattle and poultry feeds has also been reported (Thomas *et al.*, 1998). However, the occurrence of cyanogenic glycosides, linamarin and lotaustralin in this plant (*H. brasiliensis*) appears to limit its utilization for animal nutrition. This is because enzymatic hydrolysis of the glycosides by endogenous β -glucosidase, linamarase, occur when the plant tissue is damaged either mechanically or otherwise, thereby liberating HCN which is a deadly poison. Despite this constraint, some millers for livestock feed in Nigeria have started including rubber seed and other cyanogenic plant materials in their various feeds prepared for domestic animals (Okafor and Nwabuko, 2003).

A wide variety of different methods of processing the rubber seeds to reduce their content of cyanogenic glycosides and hence their toxicity are known. These methods comprise of different combinations of drying, soaking, boiling and fermentation of whole seeds; all of these processes reduce the total cyanide content of the seeds. The most effective treatments for reducing the cyanide content was reported to be 20 h of soaking in water combined with 1 h of cooking (Selle *et al.*, 1983). However, a simpler and more effective treatment could be achieved using enzymatic hydrolysis. In view of this, we thought worthwhile to evaluate the use of endogenous β -glucosidase together with over drying in the detoxification of cyanogenic glycosides as a means of processing these seeds for animal feed. The possible acute effect of such processed feed toward animals will also be investigated.

Materials and Methods

This study was carried out in the Department of Biochemistry Laboratory, Federal University of Agriculture, Umudike, Nigeria. The *Hevea brasiliensis* seeds used in this experiment were collected from the Rubber Plantation of Michael Okpara University of Agriculture Umudike, Nigeria.

Treatment of Seeds

Fresh seeds were used because the seeds deteriorate very rapidly after falling on the ground, due to moisture content. Shelled seeds were macerated/grated and the endogenous β -glucosidase was allowed to interact with the cyanogenic glycosides for 60 min. The cyanogenic glycosides content of the seeds was determined immediately after maceration and at an interval of 10 min for 60 min using the method of Egan *et al.* (1998). This was followed by oven drying the mash at 60°C for 180 min. The total cyanogen content of the mash was determined at the end of 180 min. Cyanide release from the rubber seed by exogenous β -glucosidase was also carried out. This was to monitor the extent of hydrolysis of the cyanogenic glycoside content of the seeds by the endogenous enzyme. The seeds were first extracted in orthophosphoric acid (0.1 M) solution and then incubated with linamarase from cassava with activity of 3 units/mL. The cyanide released was then determined according to the method of Egan *et al.* (1998).

Animal Experiment

The *Hevea brasiliensis* seeds processed as described above were used to formulate feed for rats. The composition of the feed was 76% rubber seed, 10% protein supplement in the form of defatted soy bean, 4% vitamin mixture, 4% oil 4% banana flavour and 2% salt mixture.

Animal Treatment

Sixteen albino rat weighing 120 g on the average were used for the feeding experiment. The rats were allowed to acclimatize to the laboratory environment for about 7 days. This was followed by starving them for 24 h to ensure that most of the cyanogens in their blood and urine if any, were eliminated. The animals were grouped into 4 with group 1 as the control. Groups 2,3 and 4 were fed rubber seed meal for 24, 48 and 72 h, respectively. The animals were sacrificed after each time interval and their blood and urine collected for cyanide, thiocyanate and liver enzyme determinations.

Assay for Cyanide and Thiocyanate

The cyanide and thiocyanate of blood and urine were measured using the methods of Essier *et al.* (1993) and Sorbo (1953), respectively.

Enzyme Assay

The assay for some plasma enzymes as indicator of changes in liver, kidney and muscle cell damage resulting from cyanide poisoning was carried out. Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were as recommended by Reitman and Frankel. (1957) and alkaline phosphatase as described by Klein *et al.* (1960). Glucose was also determined using glucose oxidase method (Passey *et al.*, 1977), total protein by Biuret method as described by Layne (1957) and serum albumin by the method of Doumas *et al.* (1971).

Results and Discussion

From the data reported in Table 1, the results show that the endogenous β -glucosidase of the seeds has high hydrolytic activity toward the cyanogenic glucosides. About 90-95% of the total cyanogenic glycosides of seeds were hydrolyzed within 60 min as measured by the release of HCN. The cyanide release by the endogenous β -glucosidase showed higher degree of hydrolytic activity when compared to that of cassava linamarase under the assay conditions.

The implication of the high hydrolytic activity of the seed β -glucosidase is that it could be used as a means of detoxification during processing of rubber seeds for animal feed or industry. This could be achieved by crushing/grating the seeds and allowing the endogenous enzyme to hydrolyze the cyanogenic glycosides content for a period of 60 min or more. This will result in the breakdown of most of the cyanogenic glycosides to give hydrocyanic acid and other non-glucosidic cyanide that could be easily eliminated from the preparation.

Oven drying of the mashed seeds at 60°C for a period of 30-180 min resulted in about 81% total cyanogen removal (Table 2). The high reduction in cyanide levels of the mashed seeds resulted from the release of endogenous β -glucosidase enzyme, which hydrolyzed the cyanogenic glycoside content to give volatile hydrocyanic acid. *H. brasiliensis* linamarase has been reported to be active even at 80°C (Vitharange *et al.*, 1991). Thus, mashing the seeds of this plant, allowing the interaction of its cyanogenic glycosides content with endogenous β -glucosidase, followed by oven drying resulted in dried mash with very low level of residual cyanide.

The benefits of processing these seeds into animal feed cannot be overemphasized. Rubber Seed Meal (RSM) could be used successfully for broilers depending on the source of animal protein (Rajaguru and Wettinmuniy, 1975). The seeds have been reported to be a good source of energy, calcium, valine, isoleucine, phenylalanine and tyrosine (Selle *et al.*, 1983). Ravindran (1995) has demonstrated that it is possible to formulate layer diets using rubber seed meal and achieve acceptable production and lower food cost under small farm conditions in tropical countries. In spite

Table 1: Total cyanide release from *Hevea brasiliensis* seed by endogenous and exogenous β -glucosidases

| Incubation time (min) | Cyanide production (mg 100 ⁻¹ g) | |
|-----------------------|---|--------------------------------|
| | Endogenous β -glucosidase | Exogenous β -glucosidase |
| 10 | 68.90 | 9.12 |
| 20 | 158.80 | 15.05 |
| 30 | 215.03 | 18.71 |
| 40 | 270.86 | 22.18 |
| 50 | 275.62 | 47.52 |
| 60 | 353.42 | 63.52 |

Total cyanide content of fresh seed was determined to be 391.6 mg 100⁻¹g.

Table 2: Percentage cyanide removal during oven drying of marshed rubber seeds at 60°C.

| Time (min) | Cyanide removal (%) |
|------------|---------------------|
| 30 | 36.89 |
| 60 | 45.86 |
| 90 | 50.32 |
| 120 | 62.18 |
| 150 | 77.38 |
| 180 | 81.27 |

Table 3: Biochemical effects on rats fed with rubber seed meal for 72 h.

| Blood parameter | T ₂₄ | T ₄₈ | T ₇₂ | C72 |
|--|-----------------|-----------------|-----------------|-------|
| Total protein (mg dL ⁻¹) | 5.23 | 5.48 | 5.39 | 5.7 |
| Serum Albumin (g dL ⁻¹) | 3.9 | 3.8 | 3.9 | 3.8 |
| Glucose (mg dL ⁻¹) | 56.00 | 60.00 | 67.00 | 48.00 |
| ALT (iu L ⁻¹) | 21.00 | 24.00 | 21.00 | 22.00 |
| AST (iu L ⁻¹) | 67.00 | 65.00 | 62.00 | 66.00 |
| Alkaline phosphatase (iu L ⁻¹) | 30.00 | 33.00 | 30.00 | 30.00 |
| Serum thiocyanate | 9.12 | 11.16 | 12.13 | 7.30 |

of these and other reports of successful use of the seeds in animal feeding, arguments still persist in scientific circles as to the safety of rubber seed meal for sustainable livestock production. Reduction in growth, response particularly in broilers and growers less than 3 months of age when rubber seed meal was supplemented for up to 30% (Rajaguru and Wettimuny, 1975), lowered egg production, thickness, hatchability of incubated eggs and weight of chicks hatched out (Rajaguru, 1973) are among the report against the use of rubber seeds in animal production especially as constituent in breeders rations. These effects may be attributed to a amino acid imbalance (Rajaguru and Wettimuny, 1975) of RSM that lowers the biological value of protein and the presence of an artificial factor which has not been identified yet (Rajaguru, 1973).

It is worthy to note that deleterious factors such as cyanogens when present in animal feed could result in growth depression and hamper hatchability of incubated eggs (Okafor and Maduagwu, 1999). In addition, they result in depletion of a amino acids of the body when ingested. This is because sulphur-containing a amino acids are needed for cyanide detoxification resulting in a condition akin to a amino acid imbalance and hence reduced protein synthesis. Thus, proper processing, of these seeds to reduce the total cyanogens content will result in improved performance when used as animal feed.

Ingestion of feed compounded from the oven dried mashed seeds containing 9.25 mg CN⁻¹ kg⁻¹ by rats resulted in non-significant changes in the blood total protein, albumin, aspartate aminotransferase (AST), alanine aminotransferase (ALT) and alkaline phosphatase activities; (Table 3) indicating that such feed may not have acute toxic effects on animals. Therefore the use of rubber seed meal as animal feed should be encouraged and the use of its endogenous enzyme for processing during feed formulation should be imbibed.

References

- Doumas, B.T., W.A. Watson and H.G. Briggs, 1971. Albumin standards and the measurement of serum albumin with Bromocresol Green. *Clinical Chimica Acta*, 31: 87.
- Egan, S.V., H.H. Yeoh and J.H. Bradbury, 1998. Simple picrate paper kit for determination of cyanogenic potential of cassava flour. *J. Sci. Food Agric.*, 75: 258-262.
- Esser A.J.A., M. Bosvold, R.M. Van Der Grift III and A.G.A. Voragen, 1993. Studies on the quantification of specific cyanogens in cassava products and introduction of a new chromogen. *J. Sci. Food Agric.*, 63: 287- 296.
- FAO, 1992. Quarterly Bulletin of Statistics 5(3).
- Iyayi, E.A and O.O. Tewe, 1994. Cassava feeding in a small holder livestock units *Acta Horticulturae*, 375: 261-269.
- Klein, B., P.A. Read and L.A. Babson, 1960. Rapid method for quantitative determination of alkaline phosphatase. *Clin. Chem.*, 6: 269-275.
- Layne, E., 1957. Spectrophotometric and turbidometric methods for measuring proteins: In: Colowick, S.P. and Kaplan, N.O. (Eds). *Methods in Enzymology*. Vol. III. New York, Academic Press.
- Ly, J., Ty. Chhay and C. Phiny, 2001. Evaluation of nutrients of rubber seed meal in Mong Cai pigs. *Livestock Research for Rural Development*, 13: 2
- Nwokolo, E., 1987. Biochemical and nutritional qualities of rubber seed meal. *Trop. Agri., (Trinidad)*. 64: 170-171.
- Okafor, P.N. and C.U. Nwabuko, 2003. Cyanide, nitrate and nitrite contents of livestock feeds in Umuahia, Nigeria. *Biokemistri*, 14: 57-63.
- Okafor, P.N. and E.N. Maduagwu, 1999. Toxicity evaluation of novel cyanide compounds of cassava tissue towards chicken embryos. *W.A. J. Biol. Applied Chem.*, 44: 24-27.
- Ong, H.K. and S.W. Yeong, 1977. Prospect of rubber seed meal for feeding swine and poultry. In proceedings of the symposium of Foodstuff for livestock in South East Asia. C.K Lumpur pp: 337-344.
- Passey, R.B., M.I. Gillum, J.B. Fuller, R.M. Urry and M.L. Giles, 1977. Evaluation and comparison of glucose methods and reference method recommended in proposed product class standard. *Clin. Chem.*, 23: 131.
- Preston, T.R., 1984. Estrategia parallel desarrollo de Sistemas alimenticios para el tropico. *Taller Intl. Fic-Filipinas* pp: 247-264.
- Rajaguru, A.S.A., 1973. Effects of rubber seed meal on the performance of matured chicken. *RRISL Bulletin*, 8: 39-45.
- Rajaguru, A.S.B. and S.G. Wettimuny, 1975. Rubber seed meal as a protein supplement in poultry feeding. *RRISL Bulletin*, Vol. 9.
- Ravindran, V., 1995. Evaluation of layer diet formulated from non-conventional feeding stuffs. *Br. Poul. Sci.*, 36: 165-170.
- Reitman, S. and S. Frankel, 1957. Determination of aspartate and alanine aminotransferase activities in blood serum and tissues. *Am. J. Clin. Pathol.*, 28: 56.
- Selle, C.M., E. Gonzalez de Mejia, L.G. Elias and R. Bressani, 1983. *Archivos Latino Americanos De Nutrition*, 33: 884-901.
- Sorbo, B.H., 1953. On the substrate specificity of rhodanese. *Acta Chem. Scandinavica*, 7: 1137-1145.

- Thomas, V., V.C. Mercykutty and C.K. Saraswarthyama, 1998. Rubber seeds: Its biological and industrial applications- A review. *The Planta*, 74: 437-443.
- Vitharauge, M.G., J.M.P. Nirmla and A.M. Abeyoskara, 1991. Some studies on controlling the action of lipase linamarase during rubber seed kernel processing. *J. Natl. Sci. Coun. Sri-Lanka*, 19: 143-150.