



American Journal of
**Drug Discovery
and Development**

ISSN 2150-427X



Academic
Journals Inc.

www.academicjournals.com

Biosurfactants in Pharmaceutical Industry: A Mini-Review

Eshrat Gharaei-Fathabad

Department of Pharmacognosy and Biotechnology, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

ABSTRACT

Biosurfactants can be served as green alternatives in a variety of applications including bioremediation, pharmaceuticals, agricultural disease control and cosmetics. Biosurfactant mixtures produced by microbes and they are genus- and sometimes species-specific. Because of their short fatty acid tails and polar head groups, biosurfactants are highly sticky and both hydrophilic and hydrophobic. In pharmaceuticals, biosurfactants can be used for gene delivery and recovery of intracellular products as well as they can be served as antimicrobial substances and emulsifying agents.

Key words: Biosurfactant, gene delivery, pharmaceuticals, bioemulsifier, antimicrobial activity

INTRODUCTION

A surfactant is an amphiphilic agent with both lipophilic and hydrophilic structural moieties in its molecule. Surfactants are widely used for industrial, agricultural, food, cosmetic and pharmaceutical applications. Most of these compounds are chemically synthesized and potentially cause environmentally and toxically problems (Schramm *et al.*, 2003; Makkar and Rockne, 2003). However, it is only in the past few decades that surface active molecules of microbial origin, referred to as biosurfactants, have gained considerable interest (Desai and Banat, 1997; Healy *et al.*, 1996). Biosurfactants are surface-active metabolites produced by microorganisms when grown on water miscible or oily substrates: they either remain adherent to microbial cell surfaces or are secreted in the culture broth (Abouseouda *et al.*, 2008; Correa-Bicca *et al.*, 1999; Cunha *et al.*, 2004; Das and Mukherjee, 2007). They possess the characteristic property of reducing the surface and interfacial tensions using the same mechanisms as chemical surfactants (Singh *et al.*, 2007). Microbial surfactants constitute a diverse group of surface-active molecules and are known to occur in a variety of chemical structures, such as glycolipids, lipopeptides and lipoproteins, fatty acids, neutral lipids, phospholipids and polymeric and particulate structures (Desai and Banat, 1997). The features that make them commercially promising alternatives to chemically synthesized surfactants are their lower toxicity, higher biodegradability and hence, greater environmental compatibility, better foaming properties (useful in mineral processing) and stable activity at extremes of pH, salinity and temperature (Amiriyani *et al.*, 2004; Batista *et al.*, 2006; Bento *et al.*, 2005; Bhattacharyya *et al.*, 2003; Bodour *et al.*, 2003; Cameotra and Makkar, 2004; Chen *et al.*, 2007; Christofi and Ivshina, 2002; Cohen and Exerowa, 2007). Unlike chemical surfactants, which are mostly derived from petroleum feedstock, these molecules can be produced by microbial fermentation processes (Fusconi *et al.*, 2010; Silva *et al.*, 2009) using cheaper agrobased substrates and waste materials (Fox and Bala, 2000; Maneerat, 2005; Nitschke *et al.*, 2004; Panilaitis *et al.*, 2007; Rashedi *et al.*, 2005; Rivardo *et al.*, 2009; Rivera *et al.*, 2007; Rodrigues *et al.*, 2006a).

Although, most biosurfactants are considered to be secondary metabolites, some may play essential roles for the survival of biosurfactant-producing microorganisms through facilitating nutrient transport or microbe-host interactions or by acting as biocide agents (Dehghan-Noudeh *et al.*, 2005; Déziel *et al.*, 1996; Fernandes *et al.*, 2007; Rodrigues *et al.*, 2006a). Biosurfactant roles include increasing the surface area and bioavailability of hydrophobic water-insoluble substrates, heavy metal binding (Kavamura and Esposito, 2010; Hoffiman *et al.*, 2010), bacterial pathogenesis and biofilm formation (Fiechter, 1992; Gautam and Tiagi, 2006; Healy *et al.*, 1996; Simoes *et al.*, 2010).

In various industrial processes, they are potentially useful surface-active agents for emulsion (Dhakephalkar *et al.*, 2010; Amaral *et al.*, 2009; Huang *et al.*, 2010). Polymerization, wetting, foaming, phase dispersion, emulsification and de-emulsification (Kosaric, 1992; Martinez-Checa *et al.*, 2007). Biosurfactants have also been found to possess several properties of therapeutic and biomedical importance (Gudiana *et al.*, 2010). They have antibacterial, antifungal (Joshi *et al.*, 2008) and antiviral properties; they inhibit fibrin clot formation and they have anti-adhesive action against several pathogenic microorganisms (Mulligan, 2005; Rodrigues *et al.*, 2004, 2006a; Singh and Cameotra, 2004).

Recently, biosurfactants received much attention in nanobiotechnology criteria (Koopmans and Aggeli, 2010; Palanisamy, 2008; Rodriguez *et al.*, 2010; Cevc and Vierl, 2010; Solanki and Murthy, 2010; Reddy *et al.*, 2009; Palanisamy and Riachur, 2009).

Surface-active compounds produced by microorganisms are of two main types, those that reduce surface tension at the air-water interface (biosurfactants) and those that reduce the interfacial tension between immiscible liquids, or at the solid-liquid interface (bioemulsifiers). Biosurfactants usually exhibit emulsifying capacity but bioemulsifiers do not necessarily reduce surface tension (Kakugawa *et al.*, 2002; Konishi *et al.*, 2007; Langer *et al.*, 2006; Freitas *et al.*, 2009).

Here, we discuss the role and applications of biosurfactants focusing on medicinal pharmaceutical perspectives.

BIOSURFACTANT CLASSIFICATION

Unlike chemically synthesized surfactants, which are classified according to the nature of their polar grouping, biosurfactants are categorized mainly by their chemical composition and their microbial origin. According to the studies of Desai and Banat (1997) and Gautam and Tiagi (2006), biosurfactants, based on the structure of their hydrophilic part, are mainly classified into 5 categories:

- Glycolipids
- Lipopeptides
- Fatty acids
- Polymer type
- Particulate biosurfactants (Desai and Banat, 1997; Gautam and Tiagi, 2006).

Glycolipids: Most known biosurfactants are glycolipids. They are carbohydrates in combination with long-chain aliphatic acids or hydroxyaliphatic acids. Among the glycolipids, the best known are rhamnolipids, trehalolipids and sophorolipids (Morita *et al.*, 2006; Sullivan, 1998; Thanomsab *et al.*, 2007).

Lipopeptides and lipoproteins: A large number of cyclic lipopeptides including decapeptide antibiotics (gramicidins) and lipopeptide antibiotics (polymyxins) possess remarkable surface-active properties.

Fatty acids, phospholipids and neutral lipids: Several bacteria and yeasts produce large quantities of fatty acid and phospholipid surfactants during growth on n-alkanes.

Polymeric biosurfactants: The best-studied polymeric biosurfactants are emulsan, liposan, mannoprotein and other polysaccharide-protein complexes.

Particulate biosurfactants: Extracellular membrane vesicles partition hydrocarbons to form a microemulsion which plays an important role in alkane uptake by microbial cells (Monteiro *et al.*, 2007; Mukherjee *et al.*, 2006; Ortiz *et al.*, 2006).

Thus the majority of biosurfactants include low-molecular-weight glycolipids (GLs), lipopeptides (LPS), flavolipids (FLs), phospholipids and high-molecular-weight polymers such as lipoproteins, lipopolysaccharide-protein complexes and polysaccharide-protein-fatty acid complexes. Biosurfactants have a great deal of structural diversity. The common lipophilic moiety of a biosurfactant molecule is the hydrocarbon chain of a fatty acid, whereas the hydrophilic part is formed by ester or alcohol groups of neutral lipids, by the carboxylate group of fatty acids or amino acids (or peptides), organic acid in the case of flavolipids, or, in the case of glycolipids, by the carbohydrate (Rodrigues *et al.*, 2006b; Ruiz-Garc *et al.*, 2005; Santa Anna *et al.*, 2002; Singh *et al.*, 2007).

Potential applications of biosurfactants in industries: Surfactants offer extraordinary benefits to many industries. They are involved in infinite number of different industrial processes and physicochemical phenomenon; increasing mobility, increasing solubility, lubrication, removing soil or scouring (Pei *et al.*, 2009; Lai *et al.*, 2009), wetting, rewetting, softening, retarding dyeing rate, fixing dyes, making emulsions, stabilizing dispersions, coagulating suspended solids, making foams (Hirata *et al.*, 2009), preventing foam formation and defoaming (Kosaric, 1992; Zang and Miller, 1992; Zouboulis *et al.*, 2003). The most significant application of biosurfactants was studied in bioremediation for example in removing heavy metals from soils (Asci *et al.*, 2010; Wang and Mulligan, 2009a, b; Gusiatin *et al.*, 2009; Mulligan, 2009; Frazetti *et al.*, 2009; Nayak *et al.*, 2009).

In addition of all these benefits, biosurfactants have a large number of bioactivities: inhibit bacterial growth (Flagas and Makris, 2009; Sabate *et al.*, 2009), toxic effects, immune stimulant, tumor growth inhibition, antibiotic, cell lysis (haemolysis) (Dehghan-Noudeh *et al.*, 2005), plant pathogenicity (Joshi *et al.*, 2008), effects on migration of human neutrophils, respiratory action (anti- asthma activity), food digestion (Nitschke and Costa, 2007), inhibition of cell wall synthesis, fungicidal properties (Joshi *et al.*, 2008) or enzyme stimulation, bio regulatory effects, etc. Hence they play part in many processes in nature (Rodrigues *et al.*, 2006b).

These molecules have an unlimited number of uses that involves every industry and every aspect of life: oil industry, pharmaceuticals, testing quality of condoms, hygiene and cosmetics, cement, beer and beverages, textiles, paint, detergents and cleaning (Rai and Mukherjee, 2010) and food processing (Arauz *et al.*, 2009). However, the applications depend on applied properties and the mechanism of action.

Potential applications of biosurfactants in pharmaceuticals: Kakugawa *et al.* (2002), as well as Mukherjee *et al.* (2006) demonstrated that the biosurfactants could have a wide range of applications in pharmaceutical fields.

Gene delivery: The establishment of an efficient and safe method for introducing exogenous nucleotides into mammalian cells is critical for basic sciences and clinical applications such as gene therapy. Among various methods for gene transfection, (Zhang *et al.*, 2010; Fujita *et al.*, 2009; Liu *et al.*, 2010) lipofection using cationic liposomes is considered to be a promising way to deliver foreign gene to the target cells without side-effects. Although, several kinds of cationic liposomes for lipofection have been developed, further studies are still required to develop a non-viral vector which has comparable efficiency to viral vectors (Inoh *et al.*, 2001, 2004; Kitamoto *et al.*, 2002; Maitani *et al.*, 2007; Okayama *et al.*, 1997).

Kitamoto *et al.* (2002) demonstrated that in comparison with commercially available cationic liposomes, liposomes based on biosurfactants show increasing efficiency of gene transfection. Ueno *et al.* (2007) have been developing some new techniques and methodologies for the liposome-based gene transfection. They introduced biosurfactants in this field. They examined MEL-A-containing liposomes for gene transfection.

Immunological adjuvants: Bacterial lipopeptides constitute potent non-toxic and non-pyrogenic immunological adjuvants when mixed with conventional antigens. A marked enhancement of the humoral immune response was obtained with the low molecular mass antigens iturin AL, herbicolin A and microcystin (MLR) coupled to poly-L-lysine (MLR-PLL) in rabbits and in chickens (Rodrigues *et al.*, 2006b).

Inhibition the adhesion of pathogenic organisms to solid surfaces: Biosurfactants have been found to inhibit the adhesion of pathogenic organisms to solid surfaces or to infection sites (Das *et al.*, 2009); thus, prior adhesion of biosurfactants to solid surfaces might constitute a new and effective means of combating colonization by pathogenic microorganisms (Rivardo *et al.*, 2009). Pre-coating vinyl urethral catheters by running the surfactin solution through them before inoculation with media resulted in a decrease in the amount of biofilm formed by *Salmonella typhimurium*, *Salmonella enterica*, *E. coli* and *Proteus mirabilis* (Rodrigues *et al.*, 2004). Furthermore, Rodrigues *et al.* (2004) demonstrated that biosurfactants greatly reduced microbial numbers on prostheses and also induced a decrease in the airflow resistance that occurs on voice prostheses after biofilm formation.

Recovery of intracellular products: Surfactants have also been used to permeabilise or lyse cells after fermentation as part of the protocol for recovery of intracellular products. Reverse micelle solutions were used for selective permeabilization of *Escherichia coli* to facilitate extraction of penicillin acylase. This process can be achieved by using biosurfactants (Singh *et al.*, 2007).

It is well known that highest efficiencies in terms of overall release of intracellular proteins from microbial cells are achieved through aggressive mechanical cell disintegration methods. However, in addition to releasing intracellular proteins, these methods solubilize most of the protein components associated with cell walls, organelles and membranes. More selective permeabilization, achieved by using reagents which render the cell envelope more porous are beneficial for selective release of target proteins where the objective is to obtain an extracted product with a high specific

activity or where further protein purification is required. Biosurfactants can be the reagents of choice for membrane permeabilization (Desai and Banat, 1997).

Thus, in the recovery of purified intracellular proteins use of selected biosurfactants to permeabilise cells with selective protein release represents a promising purification option. In selecting biosurfactants for these applications the primary consideration is the efficiency and selectivity of the surfactant in permeabilizing cells with the selective release of the desired product. It is also important to insure the chosen biosurfactant has no negative impact on the stability or activity of the product since biosurfactants may bind to proteins and other bioactive molecules (Desai and Banat, 1997; Singh *et al.*, 2007).

Antimicrobial activity: The diverse structures of biosurfactants confer them to display versatile performance (Ajesh and Sreejith, 2009; Zhao *et al.*, 2010). By its structure, biosurfactant is supposed to exert its toxicity on the cell membrane permeability as a detergent like effect. One useful property of many biosurfactants is their antimicrobial activity (Rahman and Ano, 2009). Several biosurfactants have strong antibacterial, antifungal and antiviral activity. Other medically relevant uses of biosurfactants include their role as anti-adhesive agents to pathogens, making them useful for treating many diseases and as therapeutic and probiotic agents. The MEL, a glycolipid biosurfactant, exerted the growth inhibition and differentiation-inducing activities against human leukemia cell lines by directly affect intracellular signal transduction through phosphate cascade system (Tabatabaee *et al.*, 2005; Tahzibi *et al.*, 2004; Techaoei *et al.*, 2007; Thaniyavarn *et al.*, 2006).

Das *et al.* (2009) have reported a biosurfactant produced by marine *B. circulans* that had a potent antimicrobial activity against Gram-positive and Gram-negative pathogenic and semi-pathogenic microbial strains including MDR strains.

Fernandes *et al.* (2007) investigated the antimicrobial activity of biosurfactants from *Bacillus subtilis* R14 against 29 bacterial strains. Their results demonstrated that lipopeptides have a broad spectrum of action, including antimicrobial activity against microorganisms with multidrug-resistant profiles (Fernandes *et al.*, 2007).

Rodrigues *et al.* (2006c) mentioned in their review about biosurfactants that MELs produced by *Candida antarctica*, rhamnolipids produced by *P. aeruginosa* and lipopeptides produced by *B. subtilis*31 and *B. licheniformis* have been shown to have antimicrobial activities (Rodrigues *et al.*, 2006d).

Biosurfactants for cosmetics: Many biosurfactant properties such as emulsification and de-emulsification, foaming, water binding capacity, spreading and wetting properties effect on viscosity and on product consistency, can efficiently be utilized by the above industry.

Surfactants as emulsifiers, foaming agents, solubilizers, wetting agents, cleansers, antimicrobial agents, mediators of enzyme action in various dosage forms like creams, lotions, liquids, pastes, powders, sticks, gels, films, sprays could be used and may be replaced by biosurfactants (Tugrul and Cansunar, 2005; Tuleva *et al.*, 2002; Ueno *et al.*, 2007; Urum and Pekdemir, 2004; Villeneuve, 2007; Youssef *et al.*, 2007).

Cosmetic products using surfactants including; insect repellents, antacids, bath products, acne pads, antidandruff products, contact lens solution, hair colours and care products, deodorants, nail care, body massage accessories, lipsticks, lip makers, eye shades, mascaras, soap, tooth pastes and polishes, denture cleansers, adhesives, antiperspirants, lubricated condoms, baby products, foot

care, mousses, antiseptics, shampoos, conditioners, shampoos, conditioners, shave and depilatory products, moisturizers, health and beauty products (Schramm *et al.*, 2003). All of these applications could be replaced by using biosurfactants.

CONCLUSION

Chemically synthesized surface-active compounds are widely used in the pharmaceutical, cosmetic, petroleum and food industries. However, with the advantages of biodegradability and production on renewable-resource substrates, biosurfactants may eventually replace their chemically synthesized counterparts. So far, the use of biosurfactants has been limited to a few specialized applications because biosurfactants have been economically uncompetitive. There is a need to gain a greater understanding of the physiology, genetics and biochemistry of biosurfactant-producing strains and to improve process technology to reduce production costs (Youssef *et al.*, 2004, 2007; Zang and Miller, 1992; Zouboulis *et al.*, 2003).

REFERENCES

- Abouseouda, M., R. Maachib, A. Amranec, S. Bouderguaa and A. Nabia, 2008. Evaluation of different carbon and nitrogen sources in production of biosurfactant by *Pseudomonas fluorescens*. *Desalination*, 223: 143-151.
- Ajesh, K. and K. Sreejith, 2009. Peptide antibiotics: An alternative and effective antimicrobial strategy to circumvent fungal infections. *Peptide*, 30: 999-1006.
- Amaral, P., M. Colao, M.A. Coelho, G. Fontes and M. Nele, 2009. Characterization of a bioemulsifier produced from glycerol and glucose by *Yarrowia lipolytica*. *New Biotechnol.*, 25: s138-s138.
- Amiryan, A., M.M. Assadi, V.A. Saggadian and A. Noohi, 2004. Bioemulsan production by Iranian oil reservoirs microorganisms. *Iran. J. Environ. Health Sci. Eng.*, 1: 28-35.
- Arauz, L.J., A.F. Jozala, P.G. Mazzola and T.C.V. Penna, 2009. Nisin biotechnological production and application: A review. *Trends Food Sci. Technol.*, 20: 146-154.
- Asci, Y., M. Nurbas and Y.S. Acikel, 2010. Investigation of sorption/desorption equilibria of heavy metals ions on/from quartz using rhamnolipid biosurfactant. *J. Environ. Manage.*, 91: 724-731.
- Batista, S.B., A.H. Mounter, F.R. Amorim and M.R. Totola, 2006. Isolation and characterization of biosurfactant/bioemulsifier-producing bacteria from petroleum contaminated sites. *Bioresour. Technol.*, 97: 868-875.
- Bento, M.F., D.F.A. Camargo, B.C. Okeke and Jr. W.T. Frankenberger, 2005. Diversity of biosurfactant producing microorganisms isolated from soils contaminated with diesel oil. *Microbiol. Res.*, 160: 249-255.
- Bhattacharyya, S., M. Ghosh and D.K. Bhattacharyya, 2003. *Pseudomonas* strains as source of microbial surface active molecules. *J. Oleo Sci.*, 52: 221-224.
- Bodour, A.A., K.P. Drees and R.M. Maier, 2003. Distribution of biosurfactant-producing bacteria in undisturbed and contaminated arid Southwestern soils. *Applied Environ. Microbiol.*, 69: 3280-3287.
- Cameotra, S.S. and R.S. Makkar, 2004. Recent applications of biosurfactants as biological and immunological molecules. *Curr. Opin. Microbiol.*, 7: 262-266.
- Cevc, G. and U. Vierl, 2010. Nanotechnology and transdermal rout: A state of the art review and critical appraisal. *J. Control Release*, 141: 277-299.
- Chen, C., S.C. Baker and R. Darton, 2007. The application of a high throughput analysis method for the screening of potential biosurfactants from natural sources. *J. Microbiol. Method*, 70: 503-510.

- Christofi, N. and I.B. Ivshina, 2002. Microbial surfactants and their use in field studies of soil remediation. *J. Applied Microbiol.*, 93: 915-929.
- Cohen, R. and D. Exerowa, 2007. Surface forces and properties of foam films from rhamnolipid biosurfactants. *Adv. Colloid Interfac.*, 134: 24-34.
- Correa-Bicca, F., C. Fleckl and M.A.Z. Ayub, 1999. Production of biosurfactant by hydrocarbon degrading *Rhodococcus ruber* and *Rhodococcus erythropolis*. *Rev. Microbiol.*, 30: 1-5.
- Cunha, C.D., M. Rosário, A.S. Rosado and S.G.F. Leite, 2004. *Serratia* sp. SVGG16: A promising biosurfactant producer isolated from tropical soil during growth with ethanol-blended gasoline. *Process. Biochem.*, 39: 2277-2282.
- Das, K. and A.K. Mukherjee, 2007. Comparison of lipopeptide biosurfactants production by *Bacillus subtilis* strains in submerged and solid state fermentation systems using a cheap carbon source: Some industrial applications of biosurfactants. *Process Biochem.*, 42: 1191-1199.
- Das, P., A.K. Mukherjee and R. Sen, 2009. Antiadhesive action of a marine microbial surfactant. *Colloids Surf B: Biointerfaces*, 71: 183-186.
- Dehghan-Noudeh, G., M. Housaindokht and B.S.F. Bazzaz, 2005. Isolation, characterization and investigation of surface and hemolytic activities of a lipopeptide biosurfactant produced by *Bacillus subtilis* ATCC 6633. *J. Microbiol.*, 43: 272-276.
- Desai, J.D. and I.M. Banat, 1997. Microbial production of surfactants and their commercial potential. *Microbiol. Mol. Biol. Rev.*, 61: 47-64.
- Dhakephalkar, P.K, A.G. Banpurkar and B.A. Chopade, 2010. Biosurfactants, bioemulsifiers and exopolysaccharides from marine microorganisms. *Biotechnol. Adv.*, 10.1016/j.biotechadv.2010.02.006
- Déziel, E., A. Paquette, R. Villemur, F. Lepine and J.G. Bisailon, 1996. Biosurfactant production by a soil *Pseudomonas* strain growing on polycyclic aromatic hydrocarbons. *Applied Environ. Microbiol.*, 62: 1908-1912.
- Fernandes, P.A.V., I.R. Arruda, A.F.B Santos, A.A. Araújo, A.M.S. Maior and E.A. Ximenes, 2007. Antimicrobial activity of surfactants produced by *Bacillus subtilis* R14 against multidrug-resistant bacteria. *Braz J. Microbiol.*, 38: 704-709.
- Fiechter, A., 1992. Integrated systems for biosurfactant synthesis. *Pure Applied Chem.*, 64: 1739-1743.
- Flagas, M.E. and Makris, 2009. Probiotic bacteria and biosurfactants for nosocomial infection control: A hypothesis. *J. Hospital Infection*, 71: 301-306.
- Fox, S.L. and G.A. Bala, 2000. Production of surfactant from *Bacillus subtilis* ATCC 21332 using potato substrates. *Bioresour. Technol.*, 75: 235-240.
- Frazetti, A., P. Caredda, C. Ruggeri, P.L. Colla, E. Tamburini, M. Papacchinis and G. Bestetti, 2009. Potential applications of surface active compounds by *Gordonia* sp. Strain BS29 in soil remediation technologies. *Chemosphere*, 75: 801-807.
- Freitas, F., V.D. Alves, M. Carvalheira, N. Costa, R. Oliveira and M.A.M. Reis, 2009. Emulsifying behaviour and rheological properties of extracellular polysaccharide produced by *Pseudomonas oleovorans* grown on glycerol by product. *Carbo Polymer*, 78: 549-556.
- Fujita, T., M. Furuhashi, Y. Hattori, H. Kawakami, K. Toma and Y. Maitani, 2009. Calcium enhanced delivery of tetraarginine-PEG-liquid coated DNA/Protamine complexes. *Int. J. Pharm.*, 368: 186-192.
- Fusconi, R., R.M.N., Assuncao, R.M. Guimaraes, G.R. Fiho and A.E.H. Machado, 2010. Exopolysaccharide produced by *Gordoniapolsoprenivorans* CCT 7137 in GYM commercial medium: FT-IR study and emulsifying activity. *Carbo Polym.*, 79: 403-408.

- Gautam, K.K. and V.K. Tiagi, 2006. Microbial surfactants: A review. *J. Oleo Sci.*, 55: 155-166.
- Gudiana, E.J., J.A. Teixeira and L.R. Rodrigues, 2010. Isolation and functional characterization of a biosurfactant produced by *Lactobacillus paracasei*. *Colloids Surfaces B: Biointerfaces*, 16: 298-304.
- Gusiatin, Z., E. Klimiuk, T. Pokoj and D. Kulikowaska, 2009. Biosurfactant using soil remediation highly contaminated with heavy metals. *New Biotechnol.*, 25: s287-s287.
- Healy, M.G., C.M. Devine and R. Murphy, 1996. Microbial production of biosurfactants. *Resour. Conserv. Recy.*, 18: 41-57.
- Hirata, Y.S., M. Ryu, Y. Oda, K. Igarashi, A. Nagatsuka, T. Furata and M. Sugiura, 2009. Novel characteristics of sophorolipids, yeast glycolipid biosurfactants, as biodegradable low foaming surfactants. *J. Biosci. Bioeng.*, 108: 142-146.
- Hoffman, D.R., P.P. Anderson, C.M. Schobert, M.B. Gault, W.J. Blanford and T.R. Sandrin, 2010. toxicity of *Cadmium cobalt* and copper during naphthalene biodegradation. *Bioresour. Technol.*, 101: 2672-2677.
- Huang, X.F., W. Guan, J. Liu, L.J. Lu, J. Cheng and X.Q. Zhou, 2010. Characterization and polygenetic analysis of bioemulsifier-producing bacteria. *Bioresour. Technol.*, 101: 317-323.
- Inoh, Y., D. Kitamoto, N. Hirashima and M. Nakanishi, 2001. Biosurfactants of MEL: A increase gene transfection mediated by cationic liposomes. *Biochem. Biophysic Res. Commun.*, 289: 57-61.
- Inoh, Y., D. Kitamoto, N. Hirashima and M. Nakanishi, 2004. Biosurfactant MEL: A dramatically increases gene transfection via membrane fusion. *J. Control Release*, 94: 423-431.
- Joshi, S., C. Bharucha, A.J. Desai, 2008. Production of biosurfactant and antifungal compound by fermented food isolate *Bacillus subtilis* 20B. *Bioresour. Technol.*, 99: 4603-4608.
- Kakugawa, K., M. Tamai, K. Imamura, K. Miyamoto and S. Miyoshi *et al.*, 2002. Isolation of yeast *Kurtzmanomyces* sp. i-11, novel producer of mannosylerythritol lipid. *Biosci. Biotechnol. Biochem.*, 66: 188-191.
- Kavamura, V.N. and E. Esposito, 2010. Biotechnological strategies applied to the decontamination of soils polluted with heavy metals. *Biotechnol. Adv.*, 28: 61-69.
- Kitamoto, D., Isoda H. and T.N. Hara, 2002. Functions and potential applications of glycolipid biosurfactants-from energy-saving materials to gene delivery carriers. *J. Bio. Sci. Bio. Eng.*, 94: 187-201.
- Konishi, M., T. Morita, T. Fukuoka, T. Imura, K. Kakugawa and D. Kitamoto, 2007. Production of different types of mannosylerythritol lipids as biosurfactants by the newly isolated yeast strains belonging to the genus *Pseudozyma*. *Applied Microbiol. Biotechnol.*, 75: 521-531.
- Koopmans, R.J. and A. Aggeli, 2010. Nanobiotechnology-quo vadis?. *Curr. Opin. Microbiol.*,
- Kosaric, N., 1992. Biosurfactants in industry. *Pure Applied Chem.*, 64: 1731-1737.
- Lai, C.C., Y.C. Huang, Y.H. Wei and J.S. Chang, 2009. Biosurfactant-enhanced removal of total petroleum hydrocarbons from contaminated soil. *J. Hazard Mater.*, 170: 495-505.
- Langer, O., O. Palme, V. Wray, H. Tokuda and S. Lang, 2006. Production and modification of bioactive biosurfactants. *Process Biochem.*, 41: 2138-2145.
- Liu, J., A. Zou and B. Mu, 2010. Surfactin effect on the physicochemical property of PC liposome. *Colloids Surfaces A: Physicochem. Eng. Aspects*, 10.1016/j.colsurfa.2010.03.021
- Maitani, Y., S. Igarashi, M. Sato and Y. Hattori, 2007. Cationic liposome (DC-Chol/DOPE=1: 2) and a modified ethanol injection method to prepare liposomes, increased gene expression. *Int. J. Pharma.*, 342: 33-39.

- Makkar, R.S. and K.J. Rockne, 2003. Comparison of synthetic surfactants and biosurfactants in enhancing biodegradation of polycyclic aromatic hydrocarbons. *Environ. Toxicol. Chem.*, 22: 2280-2292.
- Maneerat, S., 2005. Production of biosurfactants using substrates from renewable-resources. *Songklanakarin J. Sci. Technol.*, 27: 675-683.
- Martinez-Checa, F., F.L. Toledo, K.E. Mabrouki, E. Quesada and C. Calvo, 2007. Characteristics of bioemulsifier V2-7 synthesized in culture media added of hydrocarbons: Chemical composition, emulsifying activity and rheological properties. *Bioresour. Technol.*, 98: 3130-3135.
- Monteiro, S.A., G.L. Sasaki, L.M. Souza, J.M. Meira and J.M. Araujo *et al.*, 2007. Molecular and structural characterization of the biosurfactant produced by *Pseudomonas aeruginosa* DAUPE 614. *Chem. Phys. Lipids*, 147: 1-13.
- Morita, T., M. Konishi, T. Fukuoka, T. Imura and D. Kitamoto, 2006. Physiological differences in the formation of the Glycolipid biosurfactants, Mannosylerythritol lipids, between *Pseudozyma antarctica* and *Pseudozyma aphidis*. *Applied Microbiol. Biotechnol.*, 74: 307-315.
- Mukherjee, S., P. Das and R. Sen, 2006. Towards commercial production of microbial surfactants. *Trends Biotechnol.*, 24: 509-515.
- Mulligan, C.N., 2005. Environmental applications for biosurfactants. *Environ. Pollut.*, 133: 183-198.
- Mulligan, C.N., 2009. Recent advances in the environmental applications of biosurfactants. *Curr. Opin. Colloid Interface Sci.*, 14: 372-378.
- Nayak, A.S., M.H. Vijaykumar and T.B. Karegoudar, 2009. Characterization of biosurfactant produced by *Pseudoxanthomonas* sp. PNK-04 and its application in bioremediation. *Int. Biodeter Biodegrad*, 63: 73-79.
- Nitschke, M. and S.G. Costa, 2007. Biosurfactants in food industry. *Trends Food Sci. Technol.*, 18: 252-259.
- Nitschke, M., C. Ferraz and G.M. Pastore, 2004. Selection of microorganisms for biosurfactant production using agroindustrial wastes. *Braz J. Microbiol.*, 35: 81-85.
- Okayama, R., M. Noji and M. Nakanishi, 1997. Cationic cholesterol with a hydroxyethylamino head group promotes significantly liposome-mediated gene transfection. *FEBS Lett.*, 408: 232-234.
- Ortiz, A., J.A. Teruel, M.J. Espuny, A. Marques, A. Manresa and F.J. Aranda, 2006. Effects of dirhamnolipid on the structural properties of phosphatidylcholine membranes. *Int. J. Pharmaceutics*, 325: 99-107.
- Palanisamy, P. and A.M. Raichur, 2009. Synthesis of spherical NiO nanoparticles through a novel biosurfactant mediated emulsion technique. *Mater. Sci. Eng. C*, 29: 199-204.
- Palanisamy, P., 2008. Biosurfactant mediated synthesis of NiO nanorods. *Mater. Lett.*, 62: 743-746.
- Panilaitis, B., G.R. Castro, D. Solaiman and D.L. Kaplan, 2007. Biosynthesis of emulsan biopolymers from agro-based feedstocks. *J. Applied Microbiol.*, 102: 531-537.
- Pei, X., X. Zhan and Z. Zhou, 2009. Effect of biosurfactant on the sorption of phenanthrene onto original and H₂O₂-treated soils. *J. Environ. Sci.*, 21: 1378-1385.
- Rahman, M.S. and T. Ano, 2009. Production characteristics of lipopeptide antibiotics in biofilm fermentation of *Bacillus subtilis*. *J. Environ. Sci.*, 21: s36-s39.
- Rai, S.K. and A.K. Mukherjee, 2010. Statistical optimization of producing, purification and industrial application of a laundry detergent and organic solvent-stable subtilisin-like serine protease (Alzwiiprase) from *Bacillus subtilis* DM-04. *Biochem. Eng. J.*, 48: 173-180.

- Rashedi, H., E. Jamshidi, M.M. Assadi and B. Bonakdarpour, 2005. Isolation and production of biosurfactant from *Pseudomonas aeruginosa* isolated from Iranian Southern wells oil. *Int. J. Environ. Sci. Tech.*, 2: 121-127.
- Reddy, A.S., C.Y. Chen, S.C. Bakers, C.C. Chen and J.S. Jean *et al.*, 2009. Synthesis of silver nanoparticles using surfactin: A Biosurfactant as stabilizing agent. *Mater. Lett.*, 63: 1227-1230.
- Rivardo, F., R.J. Turner, G. Allegrone, H. Ceri and M.G. Martinotti, 2009. Anti-adhesion activity of two biosurfactants produced by *Bacillus* spp. prevents biofilm formation of human bacterial pathogens. *Applied Microbiol. Biotechnol.*, 86: 541-553.
- Rivera, O.M.P., A.B. Moldes, A.M. Torrado and J.M. Dominguez, 2007. Lactic acid and biosurfactants production from hydrolyzed distilled grape marc. *Process Biochem.*, 42: 1010-1020.
- Rodrigues, L., H.C.V. Mei, J. Teixeira and R. Oliveira, 2004. Influence of biosurfactants from probiotic bacteria on formation of biofilms on voice prostheses. *Applied Environ. Microbiol.*, 70: 4408-4410.
- Rodrigues, L., A. Moldes, J. Teixeira and R. Oliveira, 2006a. Kinetic study of fermentative biosurfactant production by *Lactobacillus strains*. *Biochem. Eng. J.*, 28: 109-116.
- Rodrigues, L., I.M. Banat, J. Teixeira and R. Oliveira, 2006b. Biosurfactants: Potential applications in medicine. *J. Antimicrobial Chemotherapy*, 57: 609-618.
- Rodrigues, L., J.A. Teixeira and R. Oliveira, 2006c. Low-cost fermentative medium for biosurfactant production by probiotic bacteria. *Biochem. Eng. J.*, 32: 135-142.
- Rodrigues, L.R., J.A. Teixeira, H.C.V. Meib and R. Oliveira, 2006d. Isolation and partial characterization of a biosurfactant produced by *Streptococcus thermophilus* A. *Coll. Surf B: Biointerfaces*, 53: 105-112.
- Rodriguez, A.P., D. Delgado, M.A. Solinis, J.L. Pedraz, E. Echevarria, J.M. Rodriguez and A.R. Gascon, 2010. Solid lipid nanoparticles as potential tools for gene therapy: *In vivo* protein expression after intravenous administration. *Inter. J. Pharm.*, 385: 157-162.
- Ruiz-Garc, C., E. Quesada, E. Martinez-Checa, I. Llamas, M.C. Urdaci and V. Be Jar, 2005. *Bacillus axarquiensis* sp. nov. and *Bacillus malacitensis* sp. nov., isolated from river-mouth sediments in Southern Spain. *Int. J. Syst. Evol. Microbiol.*, 55: 1279-1285.
- Sabate, D.C., L. Carrillo and M.C. Audisio, 2009. Inhibition of paenibacillus larvae and ascospaera apis by *Bacillus subtilis* isolated from honey bee gut and honey samples. *Res. Microbiol.*, 160: 193-199.
- Santa Annal, L.M., G.V. Sebastian, E.P. Menezes, T.L.M. Alves, A.S. Santos, Jr. N. Pereira and D.M.G. Freire,, 2002. Production of biosurfactants from *Pseudomonas aeruginosa* PA1 isolated in oil environments. *Braz J. Chem. Eng.*, 19: 159-166.
- Schramm, L.L., E.N. Stasiuk and D.G. Marangoni, 2003. Surfactants and their applications. *Ann. Rep. Program Chem. Sec.*, 99: 3-48.
- Silva, G.P., M. Mack and J. Contiero, 2009. Glycerol: A promising and abundant carbon source for industrial microbiology. *Biotechnol. Adv.*, 27: 30-39.
- Simoes, M., L.C. Simoes and M.J. Vieira, 2010. A review of current and emergent biofilm control strategies. *LWT- Food Sci. Technol.*, 43: 573-583.
- Singh, A., J.D. Hamme and O.P. Ward, 2007. Surfactants in microbiology and biotechnology. Part 2. Application aspects. *Biotechnol. Adv.*, 25: 99-121.

- Singh, P. and S.S. Cameotra, 2004. Potential applications of microbial surfactants in biomedical sciences. *Trends Biotechnol.*, 22: 142-146.
- Solanki, J.N. and Z.V.P. Murthy, 2010. Highly monodisperse and sub-nano silver particles synthesis via microemulsion technique. *Colloids Surf. A: Physicochem. Eng. Aspects*, 359: 31-38.
- Sullivan, E.R., 1998. Molecular genetics of biosurfactant production. *Curr. Opin. Biotechnol.*, 9: 263-269.
- Tabatabaee, A., M.A. Mazaheri, A.A. Noohi and V.A. Sajadian, 2005. Isolation of biosurfactant producing bacteria from oil reservoirs. *Iran. J. Environ. Health Sci. Eng.*, 2: 6-12.
- Tahzibi, A., F. Kamal and M. Mazaheri Assadi, 2004. Improved production of rhamnolipids by a *Pseudomonas aeruginosa* mutant. *Iran. Biomed. J.*, 8: 25-31.
- Techaoei, S., P. Leelapornpisid, D. Santiarwarn and S. Lumyong, 2007. Preliminary screening of biosurfactant producing microorganisms isolated from hot spring and garages in Noetheren Thailand. *KMITL Sci. Tech. J.*, 7: 38-43.
- Thaniyavarn, J., A. Chongchin, N. Wanitsuksombut and S. Thaniyavarn, 2006. Biosurfactant production by *Pseudomonas aeruginosa* A41 using palm oil as carbon source. *J. Gen. Applied Microbiol.*, 52: 215-222.
- Thanomsab, B., W. Pumeechockchai, A. Limtrakul, P. Arunrattiyakorn, W. Petchleelaha, T. Nitoda and H. Kanzaki, 2007. Chemical structures and biological activities of rhamnolipids produced by *Pseudomonas aeruginosa* B189 isolated from milk factory waste. *Bioresour. Technol.*, 98: 1149-1153.
- Tugrul, T. and E. Cansunar, 2005. Detecting surfactant-producing microorganisms by the drop-collapse test. *World J. Microbiol. Biotechnol.*, 21: 851-853.
- Tuleva, B.K., G.R. Ivanov and N.E. Christova, 2002. Biosurfactant production by a new *Pseudomonas putida* strain. *Z Naturforsch [C]*, 57: 356-360.
- Ueno, Y., N. Hirashima, Y. Inoh, T. Furuno and M. Nakanishi, 2007. Characterization of biosurfactant-containing liposomes and their efficiency for gene transfection. *Biol. Pharm. Bull.*, 30: 169-172.
- Urum, K. and T. Pekdemir, 2004. Evaluation of biosurfactants for crude oil contaminated soil washing. *Chemosphere*, 57: 1139-1150.
- Villeneuve, P., 2007. Lipases in lipophilization reactions. *Biotechnol. Adv.*, 25: 515-536.
- Wang, S. and C.N. Mulligan, 2009a. Rhamnolipid biosurfactant-enhanced soil flushing for the removal of arsenic and heavy metals from mine tailing. *Proc. Biochem.*, 44: 296-301.
- Wang, S. and C.N. Mulligan, 2009b. Arsenic mobilization from mine tailings in the presence of a biosurfactant. *Applied Geology*, 24: 928-935.
- Youssef, N.H., K.E. Duncan, D.D. Nagle, K.H. Savage, R.M. Knapp and M.J. McInerney, 2004. Comparison of methods to detect biosurfactant production by diverse microorganisms. *J. Microbiol. Methods*, 56: 339-347.
- Youssef, N., D.R. Simpson, K.E. Duncan, M.G. McInerney, M. Folmsbee, M. Fincher and R.M. Knapp, 2007. *In situ* biosurfactant production by bacillus strains injected into a limestone petroleum reservoir. *Applied Environ. Microbiol.*, 73: 1239-1247.
- Zang, Y. and R.M. Miller, 1992. Enhanced octadecane dispersion and biodegradation by a *Pseudomonas rhamnolipid* surfactant (Biosurfactant). *Applied Environ. Microbiol.*, 58: 3276-3282.

- Zhang, Y., H. Li, J. Sun, J. Gao, W. Liu, B. Li, Y. Guo and J. Chen, 2010. Dc-chol/Dope cationic liposomes: A comparative study of the influence factors on plasmid pDNA and Si RNA gene delivery. *Int. J. Pharm.*, 390: 198-207.
- Zhao, Z., Q. Wang, K. Wang, K. Brian, C. Liu and Y. Cou, 2010. Study of the antifungal activity of *Bacillus vallismortis* ZZ 185 in vitro and identification of its antifungal components. *Bioresour. Technol.*, 101: 292-297.
- Zouboulis, A.I., K.A. Matis, N.K. Lazaridis and P.N. Golyshin, 2003. The use of biosurfactants in flotation: Application for the removal of metal ions. *Mineral Eng.*, 16: 1231-1236.