



American Journal of  
**Plant Physiology**

ISSN 1557-4539



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)

### **Effect of Atmospheric Nitrogen Dioxide on Mulukhiya (*Corchorus olitorius*) Growth and Flowering**

<sup>1</sup>S.E.H. Adam, <sup>1</sup>B.M.A. Abdel-Banat, <sup>1,2</sup>A. Sakamoto, <sup>1,2</sup>M. Takahashi and <sup>2</sup>H. Morikawa

<sup>1</sup>Department of Mathematical and Life Science, Graduate School of Science,  
Hiroshima University, Kagamiyama 1-3-1, Higashi-Hiroshima 739-8526, Japan

<sup>2</sup>Core Research for Evolutional Science and Technology,  
Japan Science and Technology Corporation, Kawaguchi 332-0012, Japan

---

**Abstract:** This study focuses on evaluation of the effect of NO<sub>2</sub> at ambient concentrations on vegetative growth and reproduction of mulukhiya (*Corchorus olitorius*) plants. Treatment of mulukhiya plants with 50 ppb NO<sub>2</sub> in a controlled chamber, promoted vegetative growth and flowering as well. Treated plants produced more biomass and flowered early, whereas the control plants produced less biomass and flowered late. This result indicates that ambient NO<sub>2</sub> boost plants' growth parameters and enhance development. NO<sub>2</sub> may act directly as an external signaling molecule to stimulate flowering and/or indirectly to promote leaf growth and hence affect the flowering signals that moves from the leaf through the phloem to the shoot apical meristem (SAM) leading to the overall performance of plant development and reproduction.

**Key words:** Mulukhiya, nitrogen dioxide, growth, flowering, fumigation

---

#### **INTRODUCTION**

Flowering in plants is controlled internally and externally by developmental and environmental signals. Genetic studies of the timing of flowering in *Arabidopsis thaliana* have revealed four major controlling pathways. The photoperiod and vernalization pathways integrate environmental signals into the floral decision, whereas the autonomous and gibberellin (GA) pathways act independently of environmental cues (Simpson and Dean, 2002). Recently, Tamaki *et al.* (2007) and Corbesier *et al.* (2007) proved that florigen, the signal that tells a plant to flower, is a protein that moves from leaves through the phloem to induce flowering in the shoot apical meristem (SAM). Exogenously supplied NO<sub>2</sub> plays important roles in plant growth and metabolism (Morikawa *et al.*, 2004; Takahashi *et al.*, 2005, 2006; Adam *et al.*, 2008).

In contrast to the general view that NO<sub>2</sub> acts as a destructive gas, at ambient levels it has been found to stimulate vegetative growth of plants such as *Nicotiana plumbaginifolia* and horticultural plants including lettuce, sunflower, cucumber and pumpkin by acting as a signaling molecule rather than as a significant nutrient source (Takahashi *et al.*, 2005; Adam *et al.*, 2008). The effect of atmospheric NO<sub>2</sub> on decontamination of cadmium by kenaf (*Hibiscus cannabinus*) has been investigated (Takahashi *et al.*, 2008). Growing of kenaf seedlings in air supplemented with NO<sub>2</sub> increased the cadmium content per stem by more than 30% than that in plants grown in NO<sub>2</sub> free air (Takahashi *et al.*, 2008).

Studies on the effects of NO<sub>2</sub> on plants have mainly concentrated on measuring changes in vegetative growth parameters and much less are known about alteration and phenology of reproductive organs. Regulation of flowering by NO<sub>2</sub> has not previously been reported.

---

**Corresponding Author:** Suaad E.H. Adam, Department of Mathematical and Life Science, Graduate School of Science, Hiroshima University, Kagamiyama 1-3-1, Higashi-Hiroshima 739-8526, Japan

In this study, the relationship between the pollutant  $\text{NO}_2$  and the floral initiation in mulukhiya (*Corchorus olitorius*) has been investigated. Mulukhiya, [also called moroheiya in Japanese], is a nutritious vegetable known throughout the world, especially in the Middle East and Africa. While investigating the effects of  $\text{NO}_2$  on this vegetable species, we found that fumigation with 50 ppb  $\text{NO}_2$  stimulates its vegetative growth as well as flowering initiation. Present findings suggest that  $\text{NO}_2$  may act on mulukhiya by modulating the flowering regulator(s) directly as an external signal or indirectly by modulating the endogenous flowering stimulus.

## MATERIALS AND METHODS

This study was carried out at the Department of Mathematical and Life Sciences, Graduate School of Science, Hiroshima University, Japan from March 2005 to March 2007.

Mulukhiya seeds were sown in plastic pots containing perlite and vermiculite (v/v) and then divided into two groups. Each group was placed in a growth chamber, one supplied with 50 ppb  $\text{NO}_2$  and the other  $\text{NO}_2$ -free as described by Adam *et al.* (2008) and Takahashi *et al.* (2005). Double chambers were used (Fig. 1). Briefly, two glass-walled  $\text{NO}_2$ -fumigation chambers (1.5 · 1 · 0.7 m in width, height and depth; model HM1500; Nippon Medical and Chemical Instruments Co., Osaka, Japan) were placed in a confined glasshouse (6.9 · 2.4 · 3.0 m in width, height and depth; model BTH-P1-TH; Nippon Medical and Chemical Instruments Co.). The air entering the glasshouse (at a rate of 4 m<sup>3</sup> min<sup>-1</sup>) from outside was scrubbed of  $\text{NO}$ ,  $\text{NO}_2$  and  $\text{O}_3$  by using a scrubber with an activated charcoal and  $\text{KMnO}_4$  system (Purelite, Nippon Puretec Co., Tokyo, Japan) as described by Wildt *et al.* (1997) so that the  $\text{NO}$ ,  $\text{NO}_2$  and  $\text{O}_3$  concentrations in the glasshouse were kept at <5 nL L<sup>-1</sup>. Temperature and relative humidity in the glasshouse were controlled at 22±0.3°C and 70±4%, respectively, while  $\text{CO}_2$  concentration was kept at 340±80 µL L<sup>-1</sup> by a climate control system (CTH-G 100; Nippon Medical and Chemical Instruments Co.). Air entering the + $\text{NO}_2$  chamber (at a rate of 1 L min<sup>-1</sup>) was supplemented with  $\text{NO}_2$ . The final concentration of  $\text{NO}_2$  in the chamber was kept at 50±10 nL L<sup>-1</sup>. Air entering the - $\text{NO}_2$  chamber at the same rate as that for + $\text{NO}_2$  chamber




Fig. 1: Sketch describes the fumigation chambers used in this research

was not supplemented with  $\text{NO}_2$  but was kept at  $<5 \text{ nL L}^{-1} \text{NO}_2$ . Both chambers were equipped with independent climate control systems (CTH-C 70; Nippon Medical and Chemical Instruments Co.) to maintain temperature,  $\text{CO}_2$  concentration and relative humidity at the same values as those for the glasshouse.

Plants were grown for 6 weeks, irrigated with half-strength MS medium (Murashige and Skoog, 1962). The numbers of flowers and flower buds were counted at the time of harvest. Shoot biomass and root biomass were measured after the samples were freeze-dried. Total leaf area and stem length of each sample were measured.

For further analysis of flowering time and seed formation, experiments under the same conditions were performed for a longer time (about 9 weeks) to allow complete flowering and seed setting. Statistical significance of the data was evaluated using student's t-test.

## RESULTS AND DISCUSSION

Treatment with  $\text{NO}_2$  for a prolonged period (6 weeks) at a concentration of 50 ppb activated flowering and seed formation in mulukhiya (Fig. 2). Moreover, the treatment stimulated all measured growth parameters (Fig. 3). Shoot and root biomass were increased in treated plants compared to the controls and similar results were obtained with total leaf area and stem length. However, there was no difference in the number of leaves.




Fig. 2:  $\text{NO}_2$  enhances mulukhiya growth and development. (A) A photograph of plants fumigated treated with 50 ppb  $\text{NO}_2$  (right) for 6 weeks and the controls ( $\text{NO}_x < 5 \text{ ppb}$ ) (left). (B) 9 weeks-old pods of the fumigated treated (right) and the control plants (left). (C) The relative maturity of the seeds of the treated plants (top) and the controls (bottom). Bar, 1 cm




Fig. 3: Effects of NO<sub>2</sub> treatment on the growth and flowering of mulukhiya plants. Shown are leaf number, total leaf area, shoot biomass, root biomass, flower bud number and flower number from plants treated with 50 ppb NO<sub>2</sub> for 6 weeks under natural lighting and irrigation with a medium containing 10 mM KNO<sub>3</sub>. Values for treated plants (grey columns) are normalized relative to a value of 100 for the control plants (black columns). Control plants had leaf number 9.40±1.64, total leaf area 30.58±7.80 (cm<sup>2</sup>), shoot biomass 0.05±0.02 (g), root biomass 0.007±0.003 (g), flower bud number 2.10±0.74 and flower number 0.01±0.32 (n = 8, except for the total leaf area n = 3)

During the 6 weeks, treated plants started to flower earlier than the control plants. At the time of harvest, 7 out of 8 plants had at least one open flower. The mean value of the flowers number was 0.8 for the treated plants and 0.1 for the control plants and all plants had a larger number of flower buds (4.25) compared to the controls (2.1) with no open flowers (Fig. 2, 3).

Shoot and root biomass showed a significant increase upon treatment: 0.1 and 0.01 g for shoot and root, respectively, while the controls showed 0.05 and 0.007 g for shoot and root, respectively (Fig. 3), which is in a general agreement with the previous reports (Morikawa *et al.*, 2004; Takahashi *et al.*, 2005; Adam *et al.*, 2008) that showed NO<sub>2</sub> stimulates plant growth.

The number of leaves was not affected by NO<sub>2</sub>. The number of leaves produced on the primary shoot before the first flower is initiated is a standard indicator for flowering time. Plants with more leaves flower later than those with fewer leaves (He *et al.*, 2004). However, the total leaf area increased upon treatment with NO<sub>2</sub>. The growth conditions of the plants generally and leaves in particular might have a direct effect on the flowering decision, in addition to the other factors that control flowering time. Hence, while the direct flowering signal moves from leaves to SAM, the effect of NO<sub>2</sub> on flowering time might be by indirectly affecting leaf growth and the floral stimulus that is produced in the leaves, although Corbesier and Coupland (2006) stated that early flowering is associated with small leaves. Another possibility is that NO<sub>2</sub> might directly affect the floral decision center to speed up this process, or it may interact with or be integrated with flowering pathways to regulate the floral stimulus. In contrast, Nitric Oxide (NO) suppresses flowering of Arabidopsis plants by affecting the expression of regulatory genes in flowering pathways (He *et al.*, 2004).

In conclusion, our results indicated that 50 ppb NO<sub>2</sub> promotes not only the vegetative growth but also accelerates flowering and reproduction in mulukhiya. The importance of this observation lies on the fact that keeping all generally considered harmful gases at their naturally balanced states remediates the environment devastation threats. However, the mechanism by which NO<sub>2</sub> modulates flowering time remains to be investigated. Further study will uncover the mechanism behind these results and support the hypothesis that NO<sub>2</sub> is capable of significant regulation of the plant life cycle.

#### ACKNOWLEDGMENTS

This study was supported in part by a Grant-in-Aid for Scientific Research (No. 16208033) from the Japan Society for the Promotion of Science.

#### REFERENCES

- Adam, S.E.H., J. Shigeto, A. Sakamoto, M. Takahashi and H. Morikawa, 2008. *Atmospheric nitrogen dioxide at ambient levels stimulates growth and development of horticultural plants*. *Botany*, 86: 213-217.
- Corbesier, L. and G. Coupland, 2005. Photoperiodic flowering of *Arabidopsis*: Integrating genetic and physiological approaches to characterization of the floral stimulus. *Plant Cell Environ.*, 28: 54-66.
- Corbesier, L. and G. Coupland, 2006. The quest for florigen: A review of recent progress. *J. Exp. Bot.*, 57: 3395-3403.
- Corbesier, L., C. Vincent, S. Jang, F. Fornara and Q. Fan *et al.*, 2007. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science*, 316: 1030-1033.
- He, Y., R.H. Tang, Y. Hao, R.D. Stevens and C.W. Cook *et al.*, 2004. Nitric oxide represses the *Arabidopsis* floral transition. *Science*, 305: 1968-1971.
- Knot, T.J.E., 1934. Effect of a localized photoperiod on spinach. *Proc. Am. Soc. Hortic. Sci.*, 31: 152-154.
- Morikawa, H., M. Takahashi, A. Sakamoto, T. Matsubara and G. Arimura *et al.*, 2004. Formation of unidentified nitrogen in plants: An implication for a novel nitrogen metabolism. *Planta*, 219: 14-22.
- Murashige, T. and F. Skoog, 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol.*, 15: 473-497.
- Simpson, G.G. and C. Dean, 2002. *Arabidopsis*, the rosetta stone of flowering time?. *Science*, 296: 285-289.
- Takahashi, M., M. Nakagawa, A. Sakamoto, C. Ohsumi, T. Matsubara and H. Morikawa, 2005. Atmospheric nitrogen dioxide gas is a plant vitalization signal to increase plant size and the contents of cell constituents. *New Phytol.*, 168: 149-154.
- Takahashi, M., S.E.H. Adam, T. Furuhashi, H. Tsukaya and T. Matsubara *et al.*, 2006. Atmospheric nitrogen oxides are a plant vitalization signal. COST 859 WG2 and WG3 2nd Scientific Workshop, 31 August-2 September, Saint-Etienne, France, pp: 105-105.
- Takahashi, M., S.E.H. Adam, D. Konaka and H. Morikawa, 2008. Nitrogen dioxide at an ambient level improves the capability of kenaf (*Hibiscus cannabinus*) to decontaminate cadmium. *Int. J. Phytoremediat.*, 10: 73-76.
- Tamaki, S., S. Matsuo, L.H. Wong, S. Yokoi and K. Shimamoto, 2007. Hd3a protein is a mobile flowering signal in rice. *Science*, 316: 1033-1036.
- Wildt, J., D. Kley, A. Rockel, P. Rockel and H.J. Segsneider, 1997. Emission of NO from several higher plant species. *J. Geophys. Res.*, 102: 5919-5927.