



American Journal of
Plant Physiology

ISSN 1557-4539



Academic
Journals Inc.

www.academicjournals.com

Photosynthetic Traits and Activities of Antioxidant Enzymes in Blackgram (*Vigna mungo* L. Hepper) Under Cadmium Stress*

Sarvajeet Singh, Nafees A. Khan, Rahat Nazar and Naser A. Anjum
Plant Physiology and Biochemistry Division, Department of Botany,
Aligarh Muslim University, Aligarh-202 002, India

Abstract: Cadmium (Cd) is a non-essential heavy metal that does not have any metabolic use and can be harmful even at low concentrations. Blackgram (*Vigna mungo* L. Hepper cv. T9) plants were grown in pots containing a mixture of soil and compost treated with 0, 25, 50 and 100 mg Cd kg⁻¹ soil as CdCl₂ for 30 days. The changes in total Chlorophyll (Chl), Chl a/b, net photosynthetic rate (P_n), stomatal conductance (g_s), Water Use Efficiency (WUE) and Carbonic Anhydrase (CA) activity were noted. The activities of antioxidative enzymes in root and leaf were also assayed together with the content of Thiobarbituric Acid Reactive Substances (TBARS) and hydrogen peroxide (H₂O₂). The concentration of Cd in root and leaf increased with the increasing Cd concentrations. Greatest decrease in photosynthetic traits was observed with 100 mg Cd kg⁻¹ soil. The activity of Superoxide Dismutase (SOD) increased in leaf but decreased in root, whereas the activity of catalase (CAT) decreased in both root and leaf. By contrast to CAT, the activity of ascorbate peroxidase (APX) increased in root and leaf. However, GR activity increased in root and decreased in leaf. The results suggest that the antioxidative enzymes showed differential pattern in root and leaf and the decrease in photosynthesis with 100 mg Cd kg⁻¹ soil was associated with the accumulation of TBARS and H₂O₂ content and reduction in Chl content, stomatal conductance and CA activity.

Key words: Antioxidant enzymes, cadmium, carbonic anhydrase, oxidative stress, photosynthesis

INTRODUCTION

Cadmium (Cd) is a common metal pollutant introduced into the environment through industrial activities, sewage sludge application and commercial phosphorus fertilizers and subsequently become a part of the food chain (Wagner, 1993). It is easily taken up by plants and causes toxicity even at low concentrations (Sanita di Toppi and Gabbriellini, 1999). It reduces photosynthesis through inhibiting photosynthetic pigments synthesis (Somasekaraiah *et al.*, 1992; Drazkiewicz *et al.*, 2003; Mobin and Khan, 2007) and the enzymes involved in CO₂ fixation (Di Filippis and Ziegler, 1993; Seregin and Ivanov, 2001) and also reduces plant growth (Arduini *et al.*, 2004; Wojcik and Tukiendorf, 2005; Khan *et al.*, 2006). Plants activate antioxidative enzymes system to reduce the adverse effects of Cd stress (Dixit *et al.*, 2001; Shah *et al.*, 2001; Cho and Seo, 2005; Mobin and Khan, 2007), the response of which depends on plant species and the tissue analyzed (Gallego *et al.*, 1999; Vitoria *et al.*, 2001; Ferreira *et al.*, 2002; Fornazier *et al.*, 2002; Cardoso *et al.*, 2002). It is assumed that differential activities of antioxidative enzymes in root and leaf may reduce the adverse effects of Cd stress and protect photosynthetic machinery from oxidative stress. The purpose of the present work was to evaluate the oxidative stress, response of antioxidative enzymes system in root and leaf and photosynthetic potential of blackgram (*Vigna mungo*) subjected to cadmium stress.

Corresponding Author: Nafees A. Khan, Plant Physiology and Biochemistry Division, Department of Botany,
Aligarh Muslim University, Aligarh-202 002, India Tel: +91-571-2702016

*Originally Published in American Journal of Plant Physiology, 2008

MATERIALS AND METHODS

Plant Material and Growth Conditions

An experiment was conducted in the naturally illuminated greenhouse of the Department of Botany, Aligarh Muslim University, Aligarh, India. A mixture of soil and compost (3:1) with neutral in reaction (pH 7.1) was used for the study. The chemical properties of the soil were: organic carbon, 0.38%; CEC, 2.78 meq 100 g⁻¹ soil; nitrogen, 88.4 mg kg⁻¹ soil; phosphorus, 8.4 mg kg⁻¹ soil; potassium, 110.6 mg kg⁻¹ soil and cadmium, 0.31 mg kg⁻¹ soil. Soil was mixed with appropriate amount of CdCl₂ to achieve 0, 25, 50 and 100 mg Cd kg⁻¹ soil. Seeds of blackgram (*Vigna mungo* L. Hepper cv. T9) were sown in 23 cm diameter clay pots containing 4 kg soil on June 15, 2006. After germination, two plants per pot were maintained and watered with deionized water as and when required. Each Cd treatment as well as control was replicated three times. After 30 days of sowing, photosynthetic traits in leaves, activities of antioxidant enzymes and contents of Thiobarbituric Acid Reactive Substances (TBARS) and H₂O₂ were determined in root and leaf.

Determination of Cadmium

Cadmium content was determined in dried root and leaf samples digested in concentrated HNO₃-HClO₄ (3:1, v/v) and cadmium concentration was determined by atomic absorption spectrophotometer (GBG, 932 plus, Australia).

Measurement of Photosynthetic Traits

Leaf chlorophyll content was determined by its extraction in 90% acetone and the absorbance was read spectrophotometrically (Lichtenthaler, 1987).

The activity of Carbonic Anhydrase (CA) was determined in leaves used for photosynthetic measurement by the method of Dwivedi and Randhava (1974). Net photosynthetic rate (P_N), stomatal conductance (gs) and intercellular CO₂ concentration (Ci) were measured on fully expanded uppermost leaves on two plants per treatment using Li6200 portable photosynthesis system (LiCOR, Nebraska, USA) on a sunny day. During the measurements the air relative humidity, temperature and ambient CO₂ concentration were 68±5%, 24±2°C and 350±15 µmol mol⁻¹, respectively. Water Use Efficiency (WUE) was calculated by dividing the values of P_N with gs as described by Dudley (1996).

Determination of TBARS and H₂O₂

The content of TBARS in the root/leaf was determined as described by Dhindsa *et al.* (1981). The TBARS content was calculated using the extinction coefficient (155 mM⁻¹ cm⁻¹). The content of H₂O₂ was measured in root/leaf by the method described by Jena and Choudhuri (1981). The H₂O₂ content was calculated using the extinction coefficient (0.28 µmol⁻¹ cm⁻¹).

Enzyme Extraction and Assay

Root/leaf samples were homogenized with an extraction buffer containing 100 mM potassium phosphate buffer (pH 7.0), 0.5% Triton X-100 and 1% polyvinylpyrrolidone (PVP) using chilled mortar and pestle. The homogenate was centrifuged at 15000 x g for 20 min at 4°C. The supernatant obtained after centrifugation was used for the enzyme assays. For Ascorbate Peroxidase (APX), extraction buffer was supplemented with 2 mM ascorbate.

The activity of Superoxide Dismutase (SOD) was assayed by monitoring the inhibition of photochemical reduction of Nitroblue Tetrazolium (NBT) according to Dhindsa *et al.* (1981). One unit of SOD activity was defined as the amount of enzyme required to cause 50% inhibition of the reaction of NBT.

The Activity of Catalase (CAT) was measured by the method of Aebi (1984) and was determined by monitoring the disappearance of H₂O₂ using the extinction coefficient 0.036 mM⁻¹ cm⁻¹. One unit of enzyme was defined as the amount of enzyme necessary to decompose 1 μmol of H₂O₂ min⁻¹ at 25°C.

The activity of APX was determined according to Nakano and Asada (1981). APX activity was calculated by using extinction coefficient 2.8 mM⁻¹ cm⁻¹. One unit of enzyme is the amount necessary to decompose 1 μmol of substrate min⁻¹ at 25°C.

The activity of Glutathione Reductase (GR) was determined as described by Foyer and Halliwell (1976) by monitoring the glutathione dependent oxidation of NADPH. The activity of GR was calculated by using extinction coefficient 6.2 mM⁻¹ cm⁻¹. One unit of enzyme is the amount necessary to decompose 1 μmol of NADPH min⁻¹ at 25°C.

The protein content in the samples was determined using Bovine Serum Albumin (BSA, Sigma) as standard (Bradford, 1976).

Statistical Analysis

The results are presented as means±standard error. Data were subjected to ANOVA test (SPSS ver. 11 Inc., Chicago, USA) and means were compared by using Duncan's Multiple Range Test, taking p<0.05 as significant.

RESULTS

Cadmium Accumulation

The accumulation of Cd in root and leaf increased with increasing Cd concentration in the soil (Fig. 1). However, for each Cd treatment its concentration was greater in roots than leaves. Cd concentration in the roots and leaves increased by 2.3 and 4.9 fold, respectively, with 100 mg Cd kg⁻¹ soil compared to 25 mg Cd kg⁻¹ soil.

Photosynthetic Traits

Increasing concentration of Cd significantly decreased the photosynthetic traits, P_N, g_s, C_i, Chl and CA activity. Greatest significant reduction was observed with 100 mg Cd kg⁻¹ soil. In plants treated with 100 mg Cd kg⁻¹ soil, P_N and g_s were lowered by 55 and 50%, respectively, in comparison to control. The C_i and CA activity remained unaffected with 25 mg Cd kg⁻¹ soil, but significantly reduced with 50 and 100 mg Cd kg⁻¹ soil. In comparison to control, maximum significant reduction in C_i and CA activity of 19 and 49% with 100 mg Cd kg⁻¹ soil was observed. It was also noticed that 100 mg Cd kg⁻¹ soil significantly reduced Chl content. Maximum reduction in Chl content of 43% was noted with 100 mg Cd kg⁻¹ soil in comparison to control. The ratio of Chl a to Chl b increased with the increasing Cd concentration (Table 1).

Table 1: Changes in chlorophyll content (Chl), Chl a/b and net photosynthetic rate (P_N), stomatal conductance (g_s), Water Use Efficiency (WUE), intercellular CO₂ Concentration (C_i) and Carbonic Anhydrase (CA) activity of blackgram (*Vigna mungo* L. Hepper) exposed to Cadmium (Cd) after 30 days of sowing. Values are means of three replications±SE. Data followed by different letters within a row are significantly different at p<0.05

Parameters	Cadmium (Cd) concentrations (mg kg ⁻¹ soil)			
	0	25	50	100
Total Chl (mg g ⁻¹ FW)	1.730±0.02 ^a	1.680±0.09 ^a	1.240±0.05 ^b	0.980±0.01 ^c
Chl a/b	1.500±0.01 ^b	1.580±0.01 ^b	1.810±0.03 ^a	1.800±0.02 ^a
P _N (μmol CO ₂ m ⁻² sec ⁻¹)	12.070±0.30 ^a	10.310±0.45 ^b	7.760±0.27 ^c	5.470±0.248 ^d
g _s (mol m ⁻² sec ⁻¹)	0.249±0.009 ^a	0.218±0.006 ^b	0.175±0.008 ^c	0.125±0.011 ^d
WUE (μmol mol)	48.470±0.66 ^b	47.290±0.36 ^b	44.340±0.49 ^a	43.760±0.44 ^a
C _i (μmol mol ⁻¹)	290±4.61 ^a	282±4.61 ^a	263±3.51 ^b	234±3.05 ^c
CA activity (μmol CO ₂ g ⁻¹ FW)	1.620±0.047 ^a	1.400±0.125 ^a	1.170±0.018 ^b	0.820±0.005 ^c




Fig. 1: Changes in cadmium accumulation in root and leaf of blackgram (*Vigna mungo* L. Hepper cv. T9) exposed to Cadmium (Cd) after 30 days of sowing. Values are means of three replications \pm SE. Data followed by different letter(s) in a graph line are significantly different at $p < 0.05$

Contents of TBARS and H₂O₂

The contents of TBARS and H₂O₂ in root and leaf were measured to observe the involvement of oxidative stress (Fig. 2). Roots showed higher contents of TBARS and H₂O₂ than leaves. In roots and leaves, TBARS increased maximally by 57 and 73% with 50 mg Cd kg⁻¹ soil in comparison to control. The effect of 50 and 100 mg Cd kg⁻¹ soil was not significantly different. Leaves and roots exhibited an increase of 113 and 166% in H₂O₂ content with 100 mg Cd kg⁻¹ soil in comparison to control (Fig. 1).

Antioxidant Enzyme Activities

Antioxidative enzymes responded differentially in roots and leaves to Cd treatments. Root SOD activity decreased significantly with 100 mg Cd kg⁻¹ soil but remained statistically non-significant with 25 and 50 mg Cd kg⁻¹ soil when compared with control. Maximum significant reduction in root SOD activity of 10% was observed with 100 mg Cd kg⁻¹ soil in comparison to control. In leaves, SOD activity was increased with increasing Cd concentration. Leaf SOD activity was significantly increased by 23% with 100 mg Cd kg⁻¹ soil in comparison to control, whereas, the effect of 25 and 50 mg Cd kg⁻¹ soil remained non-significant (Fig. 3).

In comparison to control, 25 mg Cd kg⁻¹ soil did not showed any change in root CAT activity but significant decrease in its activity was observed with 50 and 100 mg Cd kg⁻¹ soil. Significant reduction in root CAT activity of 46% was observed with 100 mg Cd kg⁻¹ soil. CAT activity in the leaves remained unchanged with 25 and 50 mg Cd kg⁻¹ soil, whereas, 100 mg Cd kg⁻¹ soil caused significant reduction of 25% in its activity when compared with control.

The activity of APX in roots and leaves increased with increasing Cd concentrations. Roots showed greater increase in APX activity than the leaves. In roots, 25 and 50 mg Cd kg⁻¹ soil significantly increased the APX activity but it remained same as in 50 mg Cd kg⁻¹ soil and 100 mg Cd kg⁻¹ soil. Maximum significant increase of 271% in root APX activity was observed with 50 mg Cd kg⁻¹ soil compared to control. APX activity in leaves was increased significantly with all Cd concentrations compared to control. Maximum increase in leaf APX activity of 113% was observed with 50 mg Cd kg⁻¹ soil in comparison to control.

Cd treatments increased the GR activity in roots but decreased in leaves with 100 mg Cd kg⁻¹ soil. In roots, it showed greatest significant increase of 55% with 50 mg Cd kg⁻¹ soil, however, its effect was similar to 50 mg Cd kg⁻¹ soil (Fig. 3).




Fig. 2: Changes in (a) Thiobarbituric Acid Reactive Substances (TBARS) and (b) H₂O₂ content in root and leaf of blackgram (*Vigna mungo* L. Hepper cv. T9) exposed to Cadmium (Cd) after 30 days of sowing. Values are means of three replications±SE. Data followed by different letter(s) in a graph line are significantly different at p<0.05




Fig. 3: Changes in (a) Superoxide Dismutase (SOD), (b) Catalase (CAT), (c) Ascorbate Peroxidase (APX) and (d) Glutathione Reductase (GR) activity in the root and leaf of blackgram (*Vigna mungo* L. Hepper cv. T9) exposed to Cadmium (Cd) after 30 days of sowing. Values are means of three replications±SE. Data followed by different letter(s) in a graph line are significantly different at p<0.05

DISCUSSION

The contents of TBARS are considered as an index of lipid peroxidation and Cd phytotoxicity (Pandolfini *et al.*, 1992; De Vos *et al.*, 1993; Lozano-Rodriguez *et al.*, 1997). An increased level of H₂O₂ content in root and leaf due to Cd stress caused elevated generation of reactive oxygen species and

lipid peroxidation. The activities of antioxidative enzymes showed a differential pattern in root and leaf and cooperated synergistically to protect photosynthetic machinery and maintain photosynthesis. Moreover, Cd accumulation differed in root and leaf and was translocated less to leaf. Superoxide dismutase constitutes the primary step of cellular defense. It dismutates $O_2^{\cdot-}$ to H_2O_2 and O_2 . Further, the accumulation of H_2O_2 is restricted through the action of catalase or by ascorbate-glutathione cycle, where ascorbate peroxidase reduces it to H_2O . Finally glutathione reductase catalyzes the NADPH-dependent reduction of oxidized glutathione to the reduced glutathione (Noctor *et al.*, 2002). With the increasing Cd concentration the activity of SOD increased in leaf but decreased in root, whereas, the activity of CAT decreased in both root and leaf. Previous reports have also shown variable changes in the SOD activity in plants exposed to different metals including Cd (Chongpraditrum *et al.*, 1992; Somashekaraiah *et al.*, 1992; Luna *et al.*, 1994; Gallego *et al.*, 1996; Okamoto *et al.*, 2001; Schickler and Caspi, 1999). The Cd-induced decline in CAT activity has also been reported by Somashekaraiah *et al.* (1992) and Gallego *et al.* (1996). The increase in APX activity in root and leaf indicates efficient conversion of H_2O_2 to H_2O . The activity of GR was also activated in root under Cd exposure, indicating operation of ascorbate-glutathione cycle at high rate to detoxify the ROS formed in the roots and to keep the glutathione in reduced form (Cobbett, 2000).

A greater reduction in Chl content and the decrease in stomatal conductance and CA activity due to Cd cumulatively contributed to the decrease in net photosynthetic rate. The decrease in stomatal conductance due to Cd and a parallel decrease in intercellular CO_2 concentration suggest the involvement of stomatal limitations to photosynthesis. Cadmium stress also produced disturbances in water balance and thus reduction in water use efficiency was observed with Cd treatments. This might be due to the inhibition of absorption and translocation of water, as previously observed by Barcelo and Poschenrieder (1990). The decrease in photosynthesis due to Cd has also been attributed to the increase in mesophyll resistance (Lamoreaux and Chaney, 1978) and decrease in the activity of ribulose 1,5 bisphosphate carboxylase by binding the SH group of the enzyme (Stiborova *et al.*, 1986; Vassilev *et al.*, 2003). Further, the observed higher decrease in Chl b than Chl a and the increase in Chl a to Chl b ratio may be linked to the reduction in Light Harvesting Chlorophyll Proteins (LHCPs) (Loggini *et al.*, 1999) and decrease in photosynthesis due to Cd. The reduction in LHCPs content is an adaptive defence mechanism of chloroplast, which allows them to reduce the adverse condition (Asada *et al.*, 1998). The decrease in Chl content has been also shown due to the inhibition of protochlorophyllide reductase and synthesis of 5-aminolevulinic acid (Stobart *et al.*, 1985). The decrease in photosynthesis due to Cd toxicity has been reported in the literature (Sawhney *et al.*, 1990; Sheoran *et al.*, 1990; Khan *et al.*, 2006; Mobin and Khan, 2007).

It is concluded that, blackgram (*Vigna mungo*) exhibited oxidative stress in root and leaf and plants maintained a highly integrated differential antioxidative enzymes system in root and leaf to protect photosynthetic apparatus and maintain photosynthesis against oxidative damage.

REFERENCES

- Aebi, H., 1984. Catalase *in vitro*. *Methods Enzymol.*, 105: 121-126.
- Arduini, I., A. Masoni, M. Mariotti and L. Ercoli, 2004. Low cadmium application increase miscanthus growth and cadmium translocation. *Environ. Exp. Bot.*, 52: 89-100.
- Asada, K., T. Endo, J. Mano and C. Miyake, 1998. Molecular Mechanism for Relaxation of and Protection from Light Stress. In: *Stress Responses of Photosynthetic Organisms*, Saton, K. and N. Murata (Eds.). Amsterdam: Elsevier, pp: 37-52.
- Barcelo, J. and Ch. Poschenrieder, 1990. Plant water relations as affected by heavy metal stress: A review. *J. Plant Nutr.*, 13: 1-37.

- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248-254.
- Cardoso, P.F., S.M.G. Molin, G.J.G. Pereira, A.P. Vitoria and R.A. Azevedo, 2002. Response of rice inbred lines to cadmium exposure. *J. Plant Nutr.*, 25: 927-944.
- Chongpraditrum, P., S. Mori and M. Chino, 1992. Excess copper induces a cytosolic Cu, Zn-superoxide dismutase in soybean roots. *Plant Cell Physiol.*, 33: 239-244.
- Cho, U.H. and N.H. Seo, 2005. Oxidative stress in *Arabidopsis thaliana* exposed to cadmium is due to hydrogen peroxide accumulation. *Plant Sci.*, 168: 113-120.
- Cobbett, C.S., 2000. Phytochelatins and their roles in heavy metal detoxification. *Plant Physiol.*, 123: 825-832.
- De Vos, C.H.R., W.M. Ten Boukum, R. Vooijs, H. Schat and L.J. De Kok, 1993. Effect of copper on fatty acid composition and peroxidation of lipids in the roots of copper tolerant and sensitive *Silene cucubalus*. *Plant Physiol. Biochem.*, 31: 151-158.
- Dhindsa, R.S., P. Plumb-Dhindsa and T.A. Thorpe, 1981. Leaf senescence: Correlated with increased levels of membrane permeability and lipid peroxidation and decreased levels of superoxide desmutase and catalase. *J. Exp. Bot.*, 32: 93-101.
- Di Filippis, L.F. and H. Ziegler, 1993. Effect of sub lethal concentration of zinc, cadmium and mercury on the photosynthetic carbon reduction cycle of *Euglena*. *J. Plant Physiol.*, 142: 167-172.
- Dixit, V., V. Pandey and R. Shyam, 2001. Differential antioxidant responses to cadmium in roots and leaves of pea (*Pisum sativum* L., cv Azad). *J. Exp. Bot.*, 52: 1101-1109.
- Drazkiewicz, M., A. Tukendorf and T. Baszynski, 2003. Age-dependent response of maize leaf segments to cadmium treatment: Effect on chlorophyll fluorescence and phytochelatin accumulation. *J. Plant Physiol.*, 160: 247-254.
- Dudley, S.A., 1996. Differing selection on plant physiological traits in response to environmental water availability: A test of adaptive hypothesis. *Evolution.*, 50: 92-102.
- Dwivedi, R.S. and N.S. Randhava, 1974. Evaluation of a rapid test for the hidden hunger of zinc in plants. *Plant Soil.*, 40: 445-451.
- Ferreira, R.R., R.F. Fornazier, A.P. Vitoria, S.M.G. Molina, P.J. Lea and R.A. Azevedo, 2002. Changes in antioxidant enzyme activities in soybean under cadmium stress. *J. Plant Nutr.*, 25: 327-342.
- Fornazier, R.F., R.R. Ferreira, A.P. Vitoria, S.M.G. Molina, P.J. Lea and R.A. Azevedo, 2002. Effect of cadmium on antioxidant enzyme activities in sugarcane. *Biol. Plant.*, 41: 91-97.
- Foyer, C.H. and B. Halliwell, 1976. The presence of glutathione and glutathione reductase in chloroplasts: A proposed role in ascorbic acid metabolism. *Planta*, 133: 21-25.
- Gallego, S.M., M.P. Benavides and M.L. Tomaro, 1996. Effect of heavy metal ion excess on sunflower leaves: Evidence for involvement of oxidative stress. *Plant Sci.*, 121: 151-159.
- Gallego, S.M., M.P. Benavides and M.L. Tomaro, 1999. Effect of cadmium ions on antioxidant defense system in sunflower cotyledons. *Biol. Plant.*, 42: 49-55.
- Jena, S. and M.A. Choudhuri, 1981. Glycolate metabolism of three submerged aquatic angiosperms during aging. *Aquat. Bot.*, 12: 345-354.
- Khan, N.A., I. Ahmad, S. Singh and R. Nazar, 2006. Variation in growth, photosynthesis and yield of five wheat cultivars exposed to cadmium stress. *World J. Agric. Sci.*, 2: 223-226.
- Lamoreaux, R.J. and W.R. Chaney, 1978. The effect of cadmium on net photosynthesis, transpiration and dark respiration of excised silver maple leaves. *Physiol. Plant*, 43: 231-236.
- Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods Enzymol.*, 148: 350-382.
- Loggini, B., A. Scartazza, E. Brugnoli and F. Navari-Izzo, 1999. Antioxidant defense system, pigment composition and photosynthetic efficiency in two wheat cultivars subjected to drought. *Plant Physiol.*, 119: 1091-1099.

- Lozano-Rodríguez, E., L.E. Hernández, P. Bonay and R.O. Carpena-Ruiz, 1997. Distribution of cadmium in shoot and root tissues of maize and pea plants: Physiological disturbances. *J. Exp. Bot.*, 306: 123-128.
- Luna, C.M., C.A. González and V.S. Trippi, 1994. Oxidative damage caused by an excess of Cu in oat leaves. *Plant Cell Physiol.*, 35: 11-15.
- Mobin, M. and N.A. Khan, 2007. Photosynthetic activity, pigment composition and antioxidative response of two mustard (*Brassica juncea*) cultivars differing in photosynthetic capacity subjected to cadmium stress. *J. Plant Physiol.*, 164: 601-610.
- Nakano, Y. and K. Asada, 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.*, 22: 867-880.
- Noctor, G., L. Gomez, H. Vanacker and C.H. Foyer, 2002. Interactions between biosynthesis, compartmentation and transport in the control of glutathione homeostasis and signaling. *J. Exp. Bot.*, 53: 1283-1303.
- Okamoto, O.K., E. Pinto, L.R. Latorie, E.J.H. Becharie and P. Colepicola, 2001. Antioxidative modulation in response to metal-induced oxidative stress in algal chloroplasts. *Arch. Environ. Contam. Toxicol.*, 40: 18-24.
- Pandolfini, T., R. Gabbriellini and C. Comparini, 1992. Nickel toxicity and peroxidase activity in seedlings of *Triticum aestivum* L. *Plant Cell Environ.*, 15: 719-725.
- Sanita di Toppi, L. and R. Gabbriellini, 1999. Response to cadmium in higher plants. *Environ. Exp. Bot.*, 41: 105-130.
- Sawhney, V., I.S. Sheoran and R. Singh, 1990. Nitrogen fixation, photosynthesis and enzymes of ammonia assimilation and ureide biosynthesis in nodules of mungbean (*Vigna radiata*) grown in presence of cadmium. *Indian J. Exp. Biol.*, 28: 883-886.
- Schickler, H. and H. Caspi, 1999. Response of antioxidative enzymes to nickel and cadmium stress in hyperaccumulator plants of the genus *Alyssum*. *Physiol. Plant*, 105: 39-44.
- Seregin, I.V. and V.B. Ivanov, 2001. Physiological aspects of cadmium and lead toxic effects on higher plants. *Russ. J. Plant Physiol.*, 48: 606-630.
- Shah, K., R.G. Kumar, S. Verma and R.S. Dubey, 2001. Effect of Cadmium on lipid peroxidation, superoxide anion generation and activities of antioxidant enzymes in growing rice seedlings. *Plant Sci.*, 161: 1135-1144.
- Sheoran, I.S., N. Agarwal and R. Singh, 1990. Effect of cadmium and nickel on *in vivo* carbon dioxide exchange rate of pigeon pea (*Cajanus cajan* L.). *Plant Soil*, 129: 243-249.
- Somashekaraiah, B.V., K. Padmaja and A.R.K. Prasad, 1992. Phytotoxicity of cadmium ions on germinating seedlings of mung bean (*Phaseolus vulgaris*): Involvement of lipid peroxides in chlorophyll degradation. *Physiol. Plant*, 85: 85-89.
- Stiborova, M., M. Doubravona, A. Brezinova and A. Friedrich, 1986. Effect of heavy metal ions on growth and biochemical characteristics of photosynthesis of barley (*Hordeum vulgare* L.). *Photosynthetica*, 20: 418-425.
- Stobart, A.K., W.T. Griffiths, A. Ameen-Bukhari and R.P. Sherwood, 1985. The effect of Cd²⁺ on the biosynthesis of chlorophyll in leaves of bwiley. *Physiol. Plant*, 63: 291-293.
- Vassilev, A., F. Lidon, P.S. Campos, J.C. Ramalho, M.G. Barreiro and I. Yordanov, 2003. Cu-induced changes in chloroplast lipids and photosystem 2 activity in barley plants. *Bulg. J. Plant Physiol.*, 29: 33-43.
- Vitoria, A.P., P.J. Lea and R.A. Azevedo, 2001. Antioxidant enzymes responses to cadmium in radish tissues. *Phytochemistry*, 57: 701-710.
- Wagner G.J., 1993. Accumulation of cadmium in crop plants and its consequences to human health. *Adv. Agron.*, 51: 173-212.
- Wojcik, M. and A. Tukiendorf, 2005. Cadmium uptake, localization and detoxification in *Zea mays*. *Biol. Plant*, 49: 237-245.