




International Journal of **Virology**

ISSN 1816-4900



Academic
Journals Inc.

www.academicjournals.com

Comparative Epidemiology of Infection with Human Cytomegalovirus in Cairo and South London

¹Salwa El-S. Abdel Hamid, ¹Kouka S.E. Abdel-Wahab, ¹Laila H. Saleh, ³Graham E. Davis, ⁴Ian Hastie and ¹James C. Booth

¹Virology Laboratory, Department of Microbiology, Al Azhar University, Faculty of Medicine (for Girls), Cairo, Egypt

²Department of Medical Microbiology, ³Department of Child Health, ⁴Department of Geriatric Medicine, St George's Hospital Medical School, London

Corresponding Author: Salwa El-S. Abdel Hamid, Virology Laboratory, Department of Microbiology, Al Azhar University, Faculty of Medicine (for Girls), Cairo, Egypt

ABSTRACT

Human cytomegalovirus (HCMV) infection induces HCMV IgM antibodies in acute and HCMV IgG antibodies in convalescent phases while HCMV IgG antibodies last for years. Testing for HCMV IgG antibodies as an age-related pattern of infection was studied in Cairo and South London showed that HCMV infection was acquired earlier in life in Cairo than in London. At large 95.6% of 720 sera from the general population in Cairo were seropositive by the age of 5 years compared with 33.3% of 313 sera from British subjects in London. HCMV IgM antibodies were detected only in a small proportion of elderly individuals in both cities. Testing of urine specimens by virus culture showed that congenital infection with HCMV was more common in Cairo (1.28%), compared with (0.0%) in London. Urine samples from 8.6% of Egyptian children aged 2 weeks to 5 years were positive for HCMV by culture, compared with 3.6% aged 6-10 years and 5.8% aged 11-15 years. In children from London, HCMV was isolated from the urines of 11% aged 2 weeks to 5 years, 5.2% of those aged 6-10 years but from none aged 11-15 years.

Key words: Epidemiology, cytomegalovirus, infection, Cairo

INTRODUCTION

HCMV infection occurs world-wide and, once it is acquired, persists in the host probably for life either in states of latency or low level replication, with sporadic episodes of reactivation. Reactivation is detected with greater frequency in immunocompromised subjects, especially kidney and bone marrow transplant (Griscelli *et al.*, 2001; Kamar *et al.*, 2008) recipients and infection with Human Immunodeficiency Virus (HIV) (Kim *et al.*, 2006). That reactivation occurs in immunocompetent individuals is witnessed by the birth of significant numbers of congenitally-infected babies, mostly in the developing world, whilst almost everyone acquires HCMV and develops a detectable immune response within the first year of life. Molecular analysis has shown that some individuals harbor more than one strain of HCMV which suggests that exogenous reinfection, despite a specific immune response, is a feature of the natural history of HCMV (Yamamoto *et al.*, 2010; Novak *et al.*, 2008).

Although considerable information exists on the prevalence of HCMV infection, with age, in different populations, as determined by serology, somewhat less is known about the prevalence of

active infection in the general population, as shown by virus excretion, especially in developing countries. In the latter congenital HCMV affects up to 2% of births compared to 0.5% in developed countries; however, an incidence of 1.4% has been reported in one location in West Africa (Boppana *et al.*, 2001; Schopfer *et al.*, 1978).

Like congenital infection, perinatal and postnatal infection results in excretion of HCMV in infants' urine, persisting for several years until such time as the immune response matures. Cell mediated immune lymphoproliferative responses to HCMV are weaker in HCMV antibody-positive adults from Arabic and West African countries than adults native to Britain (Omisakin *et al.*, 1996; Bello and Whittle, 1991; Booth *et al.*, 1987). Whether, this is reflected in a higher prevalence of virus excretion is not known. In order to provide a direct comparison of the epidemiology of HCMV infection in developed and developing societies we have undertaken a series of investigations, in a single laboratory, on specimen materials collected in Cairo, Egypt and in a cosmopolitan population in South London, England.

MATERIALS AND METHODS

Subjects and specimens: Serum specimens were collected from 720 Egyptians: three hundred and ninety eight apparently healthy subjects whose age ranged from 16 years to 75 years; ninety sera from neonates plus children (age range from few days old up to 15 years) and one hundred and sixty six sera were obtained from pregnant females in labor. Mostly the sera were obtained from the mothers of the babies whose urines were examined for evidence of congenital CMV infection. In addition sixty six sera were obtained from hospitalized patients suffering from bilharzial complications whose ages were between 15 and 73 years old. One hundred and thirteen sera from British neonates and children and two hundred sera from British adults were kindly given by Virus Laboratory at St. George's Hospital Medical School, London.

- Urine specimens from Egyptian and British newborn babies
- Urine specimens from Egyptian and British older children and adolescents

Tests for antibody to HCMV: Sera were tested for CMV IgG by an in-house indirect Enzyme-Immuno Assay (EIA) (Booth *et al.*, 1982) and for CMV IgM by both in-house indirect EIA (Kangro *et al.*, 1984) and commercial capture EIA (CMV-IgM-ELA-Medac, Hamburg).

Detection of HCMV in urine samples: Urine specimens were inoculated into duplicate tube cultures of primary Human Embryonic Lung (HEL) fibroblasts which were examined microscopically every other day for evidence of cytopathology, for five weeks before being regarded as negative, with refeeding with fresh medium every 3-4 days. Confirmation of cytopathology as being due to CMV was by immunofluorescent staining with CMV Monoclonal Anti Body (MAB) to a CMV early protein (HCMV-3; Cogent Diagnostics Ltd., Edinburgh).

Urine specimens were also tested for CMV by a rapid culture method in monolayers of HEL fibroblasts growing in 24-well culture plates (Steel *et al.*, 1988) which were washed thoroughly with fresh medium after allowing the inoculums to adsorb for 2 h at 37°C then refed and returned to the incubator for two days. The culture medium was discarded and the cell sheets were fixed in methanol and then were stained for 1 h at 37°C with HCMV-3 monoclonal as above. After thorough washing in phosphate-buffered saline (pbs), the cell sheets were reacted 1 h at 37°C with rabbit anti-mouse F(ab)₂; (Tago, Burlingame, Ca., USA) labeled with Alkaline Phosphatase (AP)

followed by incubation with a solution of AP substrate Fast Red TR (Sigma). CMV-infected cells were identified by their prominent, red cytoplasmic staining when the culture plates were examined under an inverted microscope at 100x magnification.

RESULTS

Seroepidemiology: Of the serum samples that were tested for HCMV IgG, overall 97.1% of those from Egypt and 50.1% from Britain were positive. Among the Egyptians, 92% were already positive for HCMV IgG by 5 years of age and 100% by age 50 years (Table 1) among the British, 29-37% were positive in the age groups less than 15 years, rising to 71% in those over 45 years (Table 2).

Tests for CMV IgM were positive in the serum from a British subject aged 76 years and in three Egyptians over the age of 40, namely a 43 year-old woman, a 52 year-old woman and a 71 year-old man. All were in good health. Sera from 66 bilharzial patients positive for HCMV IgG were all negative on testing for HCMV IgM. Interestingly the mean level of CMV IgG was higher in bilharzial patients than for healthy Egyptians of the same age (Fig. 1).

Intrauterine CMV infection: Urine specimens from 178 Egyptian babies less than 2 weeks old yielded HCMV isolates in 2 cases (1.3%) (Table 3); 176 urines from babies less than 2 weeks old in South London were negative (0%) (Table 4). Both CMV and adenovirus were isolated from a urine specimen belonging to an Egyptian 6-day old healthy baby boy.

Perinatal and postnatal infection: Urines from 8.6% of Cairene children aged 2 weeks to 5 years were positive for HCMV by culture, compared with 3.6% aged 6-10 years and 5.8% aged

Table 1: Detection of CMV-IgG and IgM in sera from Egyptians of different age groups

Age range	No. of sera tested	No. of CMV-IgG positive sera (%)	No. of CMV-IgM positive sera (%)
1 day-6 months	8	8 (11)	-
7 months-1 year	7	5 (71.4)	-
13 months-5 years	46	44 (95.6)	-
6 years-10 years	13	12 (92.3)	-
11 years-15 years	16	16 (100)	-
16 years-20 years	82	79 (96.3)	-
21 years-30 years	123	119 (96.7)	-
31 years-40 years	78	77 (98.7)	-
41 years-50 years	42	41 (97.6)	1 (2.3)
Over 50 years	73	73 (100)	2 (2.7)
Total	488	474 (97.1)	3 (0.6)

Table 2: Detection of CMV-IgG and IgM in sera from Londoners (UK) of different age groups

Age range	No. of sera tested	No. of CMV-IgG positive sera (%)	No. of CMV-IgM positive sera (%)
1 day-6 months	32	13 (40.6)	0
7 months-1 year	13	6 (46.1)	0
13 months-5 years	33	11(33.3)	0
6 years-15 years	35	10(28.6)	0
16 years-24 years	41	21 (51.2)	0
25 years-35 years	125	72 (57.6)	0
Over 35 years	34	24(70.6)	1(2.9)
Total	313	157 (50.1)	1 (0.3)




Fig. 1: Frequency distribution for CMV-IgG antibodies in Egyptian healthy adults and bilharzial patients; detected by ELISA test and reported as optical density values

Table 3: CMV isolation from Egyptian children and neonates urines

Age range	No. of urines inoculated into HEL culture	No. of CMV positives(%)
Less than 2 weeks	178	2 (1.28)
2 weeks-5 years	23	2 (8.6)
6 years-10 years	55	2 (3.6)
11 years-15 years	17	1 (5.8)

* HEL: Human embryonic lung fibroblast

Table 4: CMV isolation from British children and neonates urines

Age range	No. of urines inoculated into HEL culture	No. of positives (%)
Less than 2 weeks	176	None (0)
2 weeks-5 years	451	50 (11)
6 years-10 years	211	11 (5.2)
11 years-15 years	21	0 (0)

* HEL: Human embryonic lung fibroblast

11-15 years (Table 3). In children from South London, HCMV was isolated from the urines of 11% aged 2 weeks to 5 years, from 5.2% of those aged 6-10 years but from none aged 11-15 years (Table 4).

DISCUSSION

The high prevalence of HCMV IgG from an early age in the population of Cairo is consistent with previous findings for other Middle Eastern countries (Fig. 2) (Amer *et al.*, 1989; Peckham, 1989; Daboubi and Al-Zaben, 2000; Abou-El-Yazed *et al.*, 2008). Likewise, the pattern for South London confirms previous reports for this and other similar locations (Stern and Elek, 1965). An unexpected finding was the detection of HCMV IgM particularly in older adults in Cairo. In London one HCMV- IgM case detected may have been the result of a primary infection in someone who had escaped contact exposure with HCMV in infancy, the more




Fig. 2: Comparison of prevalence of CMV-IgG antibodies detected by Elisa test among the studies Egyptian and British subjects

so because 29.4% of subjects over 35 years old were HCMV IgG-negative. However, this was unlikely to have been the explanation in the subjects from Cairo especially two of these were mothers who had children. It may be that they had experienced recent household contact reinfection or had reactivated their latent HCMV. In bilharzial patients, the failure to detect HCMV IgM, suggests that antiviral Cell Mediated Immunity (CMI) is not depressed to an extent as to favour reactivation of latent HCMV. Others have also reported detecting HCMV IgM in the elderly (Weymouth *et al.*, 1990); its presence was attributed to virus reactivation due to declining antiviral immunocompetence with increasing age (Saltzman and Peterson, 1987).

The incidence of congenital CMV infection in Egypt (1.28%) compares well with the data that are available in reports from Egypt (5.71%) (Morgan *et al.*, 2003); low socioeconomic population in London (1.4%) (Stagno *et al.*, 1986) and in Chilean and low-income Birmingham groups (1.7 and 1.9%), respectively (Stagno *et al.*, 1982). In none of these studies has the value approached the figure of (14%) reported for the Gambia by Bello and Whittle (1991). Studies in Europe and the USA have given values of 0.2-0.5% for socially advantaged communities which are not far removed from our present findings of less than 0.6% in South London (Pass, 2005). Clearly there is still much to learn about the importance of congenital HCMV infection in societies of the third world.

Congenital and perinatal HCMV infections are followed by prolonged urinary excretion of the virus which continues for many years in association with depressed specific CMI to CMV (Noyola *et al.*, 2000; Hanshaw, 1971) and such immunity is gained, generally at about the age of five years (Pass *et al.*, 1983). In the Egyptian population, however, despite infection having been acquired in early infancy, virus excretion continued at least up to the age of fifteen years which was the oldest age group in the present study. Previously we have shown that young adults in the Middle East and West Africa who are positive for HCMV IgG give lower readings in lymphocyte proliferation tests with antigen for HCMV in comparison with Caucasians in Britain who are also seropositive to HCMV (Omisakin *et al.*, 1996; Booth *et al.*, 1987). The interpretation of these

findings was that HCMV infection in early infancy, as in the Middle East and West Africa, predisposes to long lasting depressed CMI responses to HCMV. This would agree with our present findings of extended HCMV excretion in the Cairene population compared with Londoners. Moreover it is consistent with more frequent reactivation of latent HCMV among older adults in Cairo and leading to the development of specific IgM. Such findings need further substantiating by more extensive follow up of CMI responses to HCMV and the extent and duration of intercurrent active infection with HCMV in different age groups. Differences between population groups in the capacity of individuals to control latent infection with HCMV may exert an influence on the outcome kidney, bone marrow and organ transplants.

REFERENCES

- Abou-El-Yazed, E., I.M. El-Hoseny, K. Kasim, A.A. El-Sadek and A.A. Morsy, 2008. Prevalence of cytomegalovirus infection among patients undergoing hemodialysis. *Egypt J. Immunol.*, 15: 33-41.
- Amer, K., M. El-Husseini, R. Hawas, S. Abdel Rahman and F. Shaltout, 1989. Prevalence of cytomegalovirus infection in Egyptian population with special reference to pregnant women. *Egypt. J. Haematol.*, 14: 103-108.
- Bello, C. and H. Whittle, 1991. Cytomegalovirus infection in Gambian mothers and their babies. *J. Clin. Pathol.*, 44: 366-369.
- Booth, J., G. Hannington, T. Bakir, H. Stern, H. Kangro, P. Griffiths and R. Heath, 1982. Comparison of enzyme-linked immunosorbent assay, radioimmunoassay, complement fixation, anticomplement immunofluorescence and passive haemagglutination techniques for detecting cytomegalovirus IgG antibody. *J. Clin. Pathol.*, 35: 1345-1348.
- Booth, J.C., S.A. Rashid and T.A.G. Aziz, 1987. Lymphocyte transformation test responses to cytomegalovirus in healthy adults from Britain and Iraq. *J. Med. Virol.*, 23: 265-272.
- Boppana, S.B., L.B. Rivera, K.B. Fowler, M. Mach and W.J. Britt, 2001. Intrauterine transmission of cytomegalovirus to infants of women with preconceptional immunity. *N. Engl. J. Med.*, 344: 1366-1371.
- Daboubi, M. and S. Al-Zaben, 2000. Cytomegalovirus infection in women of childbearing age in Jordan. *Jordan Med. J.*, 34: 106-108.
- Griscelli, F., M. Barrois, S. Chauvin, S. Lastere, D. Bellet and J.H. Bourhis, 2001. Quantification of human cytomegalovirus DNA in bone marrow transplant recipients by real-time PCR. *J. Clin. Microbiol.*, 39: 4362-4369.
- Hanshaw, J.B., 1971. Congenital cytomegalovirus infection: A fifteen-year perspective. *J. Infect. Dis.*, 123: 555-561.
- Kamar, N., C. Mengelle, L. Esposito, J. Guitard and M. Mehrenberger *et al.*, 2008. Predictive factors for cytomegalovirus reactivation in cytomegalovirus-seropositive kidney-transplant patients. *J. Med. Virol.*, 80: 1012-1017.
- Kangro, O.H., J.C. Booth, T.M.F. Bakir, Y. Tryhorn and S. Sutherland, 1984. Detection of IgM antibodies against cytomegalovirus: Comparison of two radioimmunoassays, enzyme-linked immunosorbent assay and immunofluorescent antibody test. *J. Med. Virol.*, 14: 73-80.
- Kim, S., J.J. Snider and M.J. Gill, 2006. Cytomegalovirus disease in HIV infection: Twenty years of a regional population's experience. *Clin. Infect. Dis.*, 42: 1808-1809.

- Morgan, M.A., E.S.M. El-Ghany, N.A. Khalifa and L.R. Rasslan, 2003. Prevalence of cytomegalovirus (CMV) infection among Neonatal Intensive Care Unit (NICU) and healthcare workers. *Egypt J. Immunol.*, 10: 1-8.
- Novak, Z., S.A. Ross, R.K. Patro, S.K. Pati, R.A. Kumbla, S. Brice and S.B. Boppana, 2008. Cytomegalovirus strain diversity in seropositive women. *J. Clin. Microbiol.*, 46: 882-886.
- Noyola, D.E., G.J. Demmler, W.D. Williamson, C. Griesser and S. Sellers *et al.*, 2000. Cytomegalovirus urinary excretion and long term outcome in children with congenital cytomegalovirus infection. Congenital CMV longitudinal study group. *Pediatr. Infect. Dis. J.*, 19: 505-510.
- Omisakin, K.A., F. Davidson, D.H. Bevan, W. al Nakib, K.S. Abdel Wahab and J.C. Booth, 1996. Lymphocyte responses to human cytomegalovirus in different groups of patients in Britain and in adults from West Africa and the Middle East. *J. Med. Virol.*, 49: 311-318.
- Pass, R.F., S. Stagno, W.J. Britt and C.A. Alford, 1983. Specific cell-mediated immunity and the natural history of congenital infection with cytomegalovirus. *J. Infect. Dis.*, 148: 953-961.
- Pass, R.F., 2005. Congenital cytomegalovirus infection and hearing loss. *HERPES Camb.*, 12: 50-55.
- Peckham, C.S., 1989. Cytomegalovirus in the neonate. *J. Antimicrob. Chemother.*, 23: 17-21.
- Saltzman, R.L. and P.K. Peterson, 1987. Immunodeficiency of the elderly. *Rev. Infect. Dis.*, 9: 1127-1139.
- Schopfer, K., E. Lauber and U. Krech, 1978. Congenital cytomegalovirus infection in newborn infants of mothers infected before pregnancy. *Arch. Dis. Child.*, 53: 536-539.
- Stagno, S., M.E. Dworsky, J. Torres, T. Mesa and T. Hirsh, 1982. Prevalence and importance of congenital cytomegalovirus infection in three different populations. *J. Pediatr.*, 101: 897-900.
- Stagno, S., R.F. Pass, G. Cloud, W.J. Britt and R.E. Henderson *et al.*, 1986. Primary cytomegalovirus infection in pregnancy: Incidence, transmission to fetus and clinical outcome. *J. Am. Med. Assoc.*, 256: 1904-1908.
- Steel, H.M., J.C. Booth, Y.S. Tryhorn and H. Stern, 1988. A simple immunoalkaline phosphatase method for the rapid diagnosis of cytomegalovirus (CMV) infection. *Serodiagn. Immunother. Infect. Dis.*, 2: 193-200.
- Stern, H. and S.D. Elek, 1965. The incidence of infection with cytomegalovirus in a normal population. A serological study in Greater London. *J. Hyg. Camb.*, 63: 79-87.
- Weymouth, L.A., I.H. Gomolin, T. Brennan, S.P. Sirpenski and D.R. Mayo, 1990. Cytomegalovirus antibody in the elderly. *Intervirology*, 31: 223-229.
- Yamamoto, A.Y., M.M. Mussi-Pinhata, S.B. Boppana, Z. Novak and V.M. Wagatsuma *et al.*, 2010. Human cytomegalovirus reinfection is associated with intrauterine transmission in a highly cytomegalovirus-immune maternal population. *Am. J. Obstet. Gynecol.*, 202: 297.e1-297.e8.