



International Journal of
**Zoological
Research**

ISSN 1811-9778



Academic
Journals Inc.

www.academicjournals.com

**The Effect of Substituting Fishmeal Diets with Varying Quantities of
Ensiled Parboiled Beniseed (*Sesamum indicum*) and Raw African
Locust Bean (*Parkia biglobosa*) on the Growth Responses and
Food Utilization of the Nile Tilapia *Oreochromis niloticus***

¹B.S. Audu, ²K.M. Adamu and ¹S.A. Binga

¹Department of Zoology, University of Jos, P.M.B. 2084, Jos, Nigeria

²Department of Zoology, Delta State University, P.M.B. 1, Abraka,
Delta State, Nigeria

Abstract: A mixture of ensiled parboiled beniseed (*Sesamum indicum*) and raw African locust bean (*Parkia biglobosa*) replaced fishmeal in the diet of *Oreochromis niloticus* at 0, 30, 70 and 100% test materials inclusion levels. Each dietary treatment was randomly duplicated in tanks stocked with twenty fingerlings (2.71±0.003 g) and fed three times (08:00, 14:00 and 18:00 h) to satiation for 56 days. The result revealed best live weight gain and feed utilization in the fish fed diet D₄ followed by diet D₁ and D₃ with the least value recorded in fish fed diet D₂ level. Generally, there was significant increase (p<0.05) in the weight gain in formulated diets (D₄) and control. The result supports the suggestion that tilapia can be fed with mixture of ensiled parboiled beniseed and raw African locust bean which is hoped will reduce tremendously the over dependence on fishmeal protein and human utilization of the very scarce fish meal.

Key words: Ensilation, parboiled, raw, *Sesamum indicum*, *Parkia biglobosa*, growth, Tilapia

INTRODUCTION

The high cost and scarcity of fishmeal have prompted world wide investigation into the ways of replacing fishmeal with less expensive foodstuff (Eyo, 1989). Lovel and Limsuwan (1982) have revealed the superiority of fishmeal in growth performance due mainly to its nutritional composition and its amino acid profile needed for optimum growth. However, it has been well established that in all fish diets, irrespective of the species, protein is the most important dietary nutrient and fish meal remain one of the major sources of dietary protein (Kaushik, 1989).

Plant protein sources such as Soya bean and groundnut cake have had favourable consideration as a replacement for fishmeal because of their high nutritive value and relative low cost as reported by Jackson *et al.* (1982). However, most of these plant protein sources are poor in one essential amino acids or the other. For instance, the protein content of beniseed was found to be rich in amino acids such as leucine, arginine and methionine but was relatively low in lysine (Arnon, 1972; Kochhar, 1981). Thus, no single oilseed meal can completely replace or supply the protein requirement of tilapia (Jauncey and Ross, 1982). Tacon *et al.* (1983), Nandeeshia *et al.* (1991), Ofojekwu and Kigbu (2002) Mukhopadhyay and Ray (2005), Abimorad *et al.* (2007), Sà *et al.* (2007) and Martinez-Llorens *et al.* (2007) have reported similar work to evaluate the suitability of replacing partially fishmeal in digestibility study. The present research was conducted to evaluate the suitability of replacing partially and/completely fishmeal diet with plant protein sources (Ensiled parboiled beniseed (*Sesamum indicum*) and Raw African locust bean (*Parkia biglobosa*) on the growth response and feed utilization of the Nile tilapia (*Oreochromis niloticus*) under laboratory conditions.

Corresponding Author: B.S. Audu, Department of Zoology, University of Jos, P.M.B. 2084, Jos, Nigeria
Tel: +2340804520805

Table 1: Preliminary crude protein composition of the test materials

Test materials	Crude protein (%)
Unparboiled/Unfermented locust bean	37.76
Parboiled fermented locust bean	43.00
Unparboiled/Unfermented beniseed	17.00
Parboiled fermented beniseed	26.09

MATERIALS AND METHODS

The experiment was conducted in the undergraduate Research laboratory, Department of Zoology, University of Jos, Jos, Nigeria. A 56 day (June to August) growth trial test was conducted with a sum total of 360 mixed sex fingerlings of *Oreochromis niloticus* means weight (2.71 ± 0.003) g obtained from Rock water fish farm, Rayfield, Jos, Plateau State, Nigeria. The fingerlings twenty were stocked in each four green circular plastic tanks with replicate measuring 48 cm diameter, 24 cm depth and 30 L capacity. Water volume was maintained at 20 L with de-chlorinated municipal water supply. Fish were fed to satiation three times daily (08:00, 14:00 and 1800 h). After the last feeding, the tanks were cleaned up of any unconsumed feeds and fecal matter while lost water was replaced with dechlorinated municipal water. The temperature, pH, dissolved oxygen, free carbon (iv) oxide were determined fortnightly (Stirling, 1985). The water temperature was $22 \pm 2^\circ\text{C}$; pH was 6.60 ± 0.25 , while dissolved oxygen was $2.01 \pm 0.30 \text{ mg L}^{-1}$. The photo period was natural and all tanks had the similar lighting condition. Fecal matters were collected by sieving through a sieve of 400 μm mesh size and oven dried. Pooled feces per treatment were kept for digestibility studies. Ten fishes were randomly selected and measured bi-weekly for weight verification to the nearest 0.1 g following method described by Ufodike and Matty (1993) and returned into their tanks before the next feeding. The mixed feeds ingredients were formulated into two batches. Batch A had 200 g of raw beniseed parboiled for 5 min, cooled and introduced into a bottle Stopped and was allowed to stand for 7 days at 25°C . The sample was then sun dried on screened surgical tray. Two hundred gram of batch B consisting of raw seeds of African locust bean were washed, screened sun dried and grinded using ceramic mortar before been sieved into fine powder with sieve of mesh size 400 μm .

Four diets including control were prepared (pelleted) to contain mixture of the test ingredients and were used to substitute fishmeal at 0, 30, 70 and 100% test material inclusion resulting to 35% crude protein content required for tilapia growth (Jauncey and Ross, 1982). The proximate composition of the test diets and control were determined according to AOAC (1980) procedures. 0.05 g of inert material (chromic oxide) was added to determine protein digestibility following the methods suggested by Furukawa and Tsukahara (1996). The percentage composition of the test diets and control and the proximate composition of the four diets are shown in Table 1. There 3 fishes each were sacrificed for proximate analysis before and after the experiment according to the method described by AOAC (1980).

The significance difference ($p = 0.05$) of the result of mean weight at different level of treatment were tested using One way analysis of variance. Significance difference in the data was tested with student t-test. Growth indices, such as Specific Growth Rate (SGR), Feed Conversion Ratio, (FCR), Protein Efficiency Ratio (PER), Net Protein Utilization (NPU) were analyzed using appropriate formulae (Halver, 1989) while corrected mortality (Finney, 1971) and percentage Live Weight Gain (LWG) were also determined.

RESULTS AND DISCUSSION

The results of the proximate composition of the test diets and control were significantly different ($p < 0.05$) ranging from 21.45-29.50% crude protein; 17.68-45.07% crude fat; 9.31-18.79% crude fibre; 4.42-5.91% ash; 3.34-4.27% moisture and 7.84-30.19% nitrogen free extract (Table 2).

Table 2: Ingredients and proximate compositions of experimental diets

Ingredients	D ₁ (Control)	D ₂ (70-30)	D ₃ (30-70)	D ₄ (0-100)
Fish meal	35.00	24.50	10.50	0.00
Corn flour	58.95	45.00	26.52	12.62
Cassava flour	5.00	5.00	5.00	5.00
Vitamin premix [#]	1.00	1.00	1.00	1.00
Chromic oxide (Cr ₂ O ₃)	0.05	0.05	0.05	0.05
Beniseed/Locust bean	0.00	24.40	56.93	81.33
Proximate composition (DM%)				
Moisture	3.34	4.27	3.74	3.89
Crude protein	29.50	26.12	21.45	23.43
Crude fat	17.18	24.20	34.87	45.07
Fiber	18.79	9.31	17.10	15.35
Ash	5.34	5.91	4.50	4.42
Carbohydrate (NFE)	25.35	30.19	18.34	7.84

[#]Vit. A, 5,000,000.00 I. U; Vit. B, 1,000,000.00 I. U; Vit E, 20,000 mg; Vit. K₃ 1,000.00 mg; pantothenic acid, 4,000.00 mg; Vit B₁, 1,200.00 mg; Vit. B₂, 2,400.00 mg; Vit. B₆, 2,400.00 mg; Niacin 16,000.00 mg; Biotin, 32.00 mg; Vit. B₁₂, 10.00 mg; Folic acid, 400.00 mg; Choline chloride, 120,000.00 mg, Manganese, 40,000.00 mg; Iron, 20,000.00 mg; Zinc, 18,000.00 mg; Copper, 800.00 mg; Iodine, 620.00 mg; Cobalt, 100.00 mg; Selenium, 40.00 mg

Table 3: Growth performance of *Oreochromis niloticus* exposed to the various formulated diet during the 8 weeks exposure period

Diets	Mean weight/week				
	0	2	4	6	8
D ₁	2.70±0.04	2.76±0.02	2.89±0.03	3.01±0.04	3.33±0.07
D ₂	2.71±0.03	2.71±0.02	2.73±0.05	2.76±0.02	2.78±0.04
D ₃	2.71±0.03	2.73±0.04	2.78±0.08	2.84±0.03	2.92±0.06
D ₄	2.71±0.01	2.79±0.05	2.93±0.02	3.21±0.01	3.49±0.04

Mean±SE value obtained from 10 fish each from an aquarium with replicate

Table 4: Growth and feed utilization of Tilapia *Oreochromis niloticus* fed different formulated feed for 8 weeks

Mean values	D ₁	D ₂	D ₃	D ₄
Initial weight (g)	2.70±0.04	2.71±0.03	2.71±0.03	2.71±0.02
Final weight (g)	3.33±0.04	2.78±0.04	2.92±0.06	3.49±0.04
Live Weight Gain (LWG%)	23.33±0.03	2.58±0.04	7.75±0.03	28.78±0.02
Specific Growth Rate (SGR%)	0.37±0.02	0.05±0.02	0.14±0.03	0.45±0.01
Food Conversion Ratio (FCR)	2.21±0.02	1.09±0.02	0.48±0.03	2.48±0.01
Protein Efficiency Ratio (PER)	2.00±0.01	0.30±0.02	0.90±0.02	3.30±0.01
ADC [#]	59.86±0.02	7.93±0.03	54.00±0.02	69.63±0.02
ANPU ^{###}	17.80±0.04	16.42±0.03	12.18±0.02	25.08±0.01
Survival rate (%)	85.00	65.00	80.00	85.00
Corrected survival rate (%)	100.00	64.71	94.12	100.00

#: Apparent Digestibility Coefficient, ###: Apparent Net Protein Utilization

At the end of the exposure period 15, 45, 20 and 15% of mortalities were recorded in D₁, D₂, D₃ and D₄ experimental tanks respectively. However, using Abbott's formula for corrected mortality revealed 35.29 and 5.88% mortalities in D₂ and D₃ experimental tanks only. Abbot's formula for corrected mortality eliminates the factor that led to mortality in the experimental control tank as it relates to other experimental setup, respectively.

As a result of feeding the fish with the different level of formulated diet there was a significant difference (p<0.05) in the fish weight (Table 3) as well as feed utilization efficiency (Table 4). Growth indices such as LWG, SGR, FCR and PER did not differ significantly (p>0.05) in the fish fed test diets and control. The Apparent Digestibility Coefficient (ADC) values obtain were not statistically different (p>0.05) with the control ADC (59.86±0.02). (Table 4) Similarly, Apparent Net Protein Utilization (ANPU) of different diet revealed insignificant difference (p>0.05) as compared to the control value (17.80±0.04). The final fish moisture content revealed significance difference (p<0.05) when compared to that of the control (Table 5).

Table 5: Initial and final body composition of Tilapia fed various formulated diets for 8 weeks

Mean value	Final				
	Initial	D ₁	D ₂	D ₃	D ₄
Moisture	10.45	10.93	16.99	10.84	10.80
Crude protein	50.53	53.24	45.28	52.30	54.08
Crude fat	20.80	24.24	18.23	23.35	25.08
Ash	16.93	10.51	18.31	12.42	9.03
NFE*	1.29	1.00	1.19	1.09	0.71

*: Nitrogen Free Extract

The temperature range in this experimental set up is under the tolerance range for the test fish *Oreochromis niloticus*. The result obtained from the preliminary analysis of the test materials showed that the parboiled ensiled beniseed has a higher level of good quality protein (26.09%) compared to the protein content of the un-parboiled ensiled (17.00%) at $p < 0.05$. The fermentation process is very essential because, it liberates more nutrients in the form of essential amino acids such as lysine, methionine and the production of more energy (Stainkraus, 1983). Though the fermentation lasted for only a week, it yielded a slight increase in the crude protein content of the un-ensiled beniseed to 1.4% and when extended, has been reported by Arnon (1972) and Kochhar (1981) to be within the range of 12.22 and 20.25%.

Highest Specific Growth Rate (SGR) was observed in D₄ (0.45±0.01), followed by D₁ (0.37±0.02) indicating better nutrient utilization. Yengkokpam *et al.* (2005) also deduce same from their findings when *Catla catla* fingerlings were fed different level of corn diet. The improved growth rate observed in D₄ then D₁ may also be ascribed to the good feed utilization by the experimental fish as evidenced by the recorded high values of FCR and PER. The high crude protein, crude fat with low moisture and fiber contents recorded in diet in D₄ may be responsible for the improved growth rate observed.

High crude fat is known to enhance energy production and hence better growth rate. Therefore, the possible reason for D₂ low performance could be due to its high content of crude fiber. Davies (1985) reported that higher fiber content impede fish growth.

Live Weight Growth (LWG) that follows D₄ > D₁ > D₃ > D₂ order in this experimental setup has been attributed to high fibre content of diet (Omoregie and Ogbemudia, 1993). However, this finding strongly accepts that deduction thus high LWG correlates high value of fibre content of diet.

Apparent Digestibility Coefficient (ADC) is thought to be an important tool to investigate the digestibility performance of food ingredients used in aquaculture. The value of ADC (D₄ > D₁ > D₃ > D₂) observed in this study was similar reported by Sá *et al.* (2007) when white sea bream *Diplodus sargus* juvenile were fed with varying level of dietary protein.

The need to use plant meal in combined form to produce the cheapest and required nutrient for fish cannot be overemphasized, thus the basis of this research which Kissil and Lupatsch (2002) reported less fish growth in fish fed an 80% soy protein concentration than in fish fed with a control diet without soy protein, but when the gilthead sea bream *Sparus aurata* was completely replaced by a plant material mixture, the results were excellent.

The replacement of fishmeal by alternate sources of protein has met with varied degree of success, depending on the nature and composition of ingredients, inclusion level and method of processing. This study revealed clearly that the nature, source and composition of ingredient and their inclusion level affect the degree of digestibility. Thus ensiled parboiled beniseed *Sesamum indicum* and raw African locust bean *Parkia biglobosa* have proven to be effective substitute for fish protein in *Oreochromis niloticus* after the 56 days of exposure.

REFERENCES

- Abimorad, E.G., D.J. Carneiro and E.C. Urbinati, 2007. Growth and Metabolism of Pacu (*Piaractus mesopotamicus*, Holmberg 1887) juvenile fed diets containing different protein, lipid and carbohydrate levels. *Aquacult. Res.*, 38: 36-44.
- AOAC (Association of Analytical Chemist), 1980. Official Methods of Analysis of Analytical Chemist. Hornitz, W. (Ed.), 13th Edn., Published by AOAC.
- Arnon, I., 1972. Crop Production in Dry Regions, Plant Science Monographs. Polurin, N. (Ed.), Published by ELBS, English Language Book Society/Macmillian, London, 2: 381-387.
- Davies, S.J., 1985. The role of dietary fiber in animal nutrition recent advances. *Aquaculture*, 2: 219-249.
- Eyo, A.A., 1989. Animal Research Report (ARR). Proceedings of the 13th Annual conference of the FISCON, New Bussa. National Institute for Fresh Water Fisheries and Feed Production Project Fisheries Societies of Nigeria (FISON), pp: 13.
- Finney, J.D., 1971. Probi Analysis. Cambridge University Press, Cambridge, pp: 125-128.
- Furukawa, A. and S. Tsukahara, 1966. Acidic digestion method for the determination of chromic oxide on an index substance in the study of the digestibility of fish feeds. *Jap. Soc. Sci. Fish.*, 32: 502-506.
- Halver, J.E., 1989. Fish Nutrition. 2nd Edn., Academic Press Inc., London.
- Jackson, A.J., B.S. Capper and A.J. Matty, 1982. Evaluation of some plant protein in complete diets for the tilapia *Sarotherodon mossambicus*. *Aquaculture*, 27: 97-109.
- Jauncey, K. and B. Ross, 1982. A Guide to feeds and farming. Institute of Aquaculture, University of Stirling, Scot. UK., pp: 345.
- Kaushik, S.J., 1989. Use of Alternative Protein Sources for Intensive Rearing of Carnivorous Fishes. In: Progress in Fish Nutrition. Shiau, S.Y. (Ed.), Mar. Food Sci. Series, No. 9, Keeking Roc, Taiwan, pp: 191-201.
- Kissil, G. Wm and I. Lupatsch, 2002. Preliminary results of fish meal replacement in gilthead sea bream *Sparus aurata* by three plant proteins. 10th Symposium on Nutrition and Feeding of Fish, Rhodes, Grecian, Book of Abstracts, pp: 127.
- Kochhar, S.L., 1981. Tropical Crops: A Textbook of Economic Botany. Published by the University of Texas at Austin, pp: 155-173.
- Lovel, R.T. and T. Limsuwan, 1982. Intestinal synthesis and dietary non-essentiality of vitamin B₁₂ for *Tilapia niloticus*. *Trans. Am. Fish. Soc.*, III: 485-490.
- Martinez-Llorens, S., A.V. Monino, V.J.M. Salvador, M.P. Torres and M.J. Cerda, 2007. Soyabean meal as a protein source in gilthead sea bream (*Sparus aurata* L.) diets: Effects on growth and nutrient utilization. *Aquacult. Res.*, 28: 82-90.
- Mukhopadhyay, N. and A.K. Ray, 2005. Effect of fermentation on Apparent total and nutrient digestibility of linseed *Linum usitatissimum* meal in Rohu *Labeo rohita* fingerlings. *Acta Ichthyol. et Piscatoria*, 35: 73-78.
- Nandeesh, M.C., G.K. Sirikanth, P. Kesharamath and S.K. Das, 1991. Protein and Fat Digestibility of Five Feed Ingredients by an Indian Major Carp (*Catla catla*). In: Proceedings of the 4th Asian Fish Nutrition Workshop. De Silva, S.S. (Ed.), Asian Fish. Soc. Manila Philippines, pp: 75 -81.
- Ofojekwu, P.C. and A.A. Kigbu, 2002. Effect of substituting fishmeal with *Sasamum indicum* (L.) cake on the growth and food utilization of the Nile tilapia, *Oreochromis niloticus*. *J. Aquacult.*, 17: 45-49.
- Omoregie, E. and F.I. Ogbemudia, 1993. Effect of substituting fishmeal with palm kernel meal on the growth and food utilization of the Nile tilapia, *Oreochromis niloticus*. *The Isr. J. Aquacult.*, 45: 113-119.

- Sá, R., P. Pousao-Ferreira and A. Oliva-Teles, 2006. Effect of dietary protein and Lipid levels in growth and feed utilization of white sea bream (*Diplodus sagus*) juvenile. *Aquacult. Nut.*, 12: 310-321.
- Sá, R., P. Pousao-Ferreira and A. Oliva-Teles, 2007. Growth performance and metabolic utilization of diets with different protein: Carbohydrate ratios by white sea bream (*Diplodus sagus*) juvenile. *Aquacult. Res.*, 38: 100-105.
- Stainkraus, K.H., 1983. Indonesian Tempe and Related Fermentation: Protein Rich Vegetarian Meat Substitute. In: *Hand Book of Indigenous Fermented Food*. Marcel Dekker Inc. New York.
- Stirling, H.P., 1985. *Water Analysis for Aquaculturist*. 1st Edn., Institute of Aqua, University of Stirling. Britain, pp: 47-50.
- Tacon, A.G.T., K. Jauncey, A. Falaye, M. Pantha, I. MacCowan and E.A. Stafford, 1983. The use of meat and bone meal, hydrolyzed feather meal and soya bean in practical fry and fingerlings diets for *Oreochromis niloticus*. Symposium on Tilapia in Aquaculture, proceedings of the Nazareth, Isreal. Tel Aviv University of Isreal, pp: 356-365.
- Ufodike, E.B.C. and A.J. Matty, 1983. Growth responses and nutrient digestibility in mirror carp, *Cyprinus carpio* fed different levels of cassava and rice. *Aquaculture*, 31:41-50.
- Yengkokpam, S., N.P. Sahu, A.K. Pal, S.C. Mukherjee and D. Debnath, 2005. Haematological and Hepatic changes in *Catla catla* fingerlings in relation to dietary sources and levels of gelatinized carbohydrate. *Acta Ichthyologica et Piscatoria*, 35: 87-92.