



Journal of
**Fisheries and
Aquatic Science**

ISSN 1816-4927



Academic
Journals Inc.

www.academicjournals.com

Collection of Broodstock and Juveniles of *Macrobrachium vollenhovenii* from the Wild

P.E. Anyanwu, I.K. Ayaobu-Cookey, B.I. Ebonwu, M.B. Hamzat, P.C.G. Pepple, A.F. Ihimekpen, M.A. Matanmi, G.O. Ikoyo-Eweto, E.S. Afolabi, M.A. Abass, B.L. Benson, M.R. Ajijo, M.M. Abdullahi, N.M. Achilike and A.S. Tafida

Nigerian Institute of Oceanography and Marine Research, PMB 12729, Victoria Island, Lagos, Nigeria

Corresponding Author: I.K. Ayaobu-Cookey, Nigerian Institute of Oceanography and Marine Research, PMB 12729, Victoria Island, Lagos, Nigeria

ABSTRACT

A total of 457 broodstock and 143,000 juveniles of *Macrobrachium vollenhovenii* were collected at Isheri Olofin area and Okunmanya River over a period of time. They were transported live to the Shrimp hatchery of Nigerian Institute for Oceanography and Marine Research (NIOMR) in Victoria Island Lagos. Live transportation of the broodstock of *M. vollenhovenii* with body weight ranging from 71.0-264.5 g in 50 L capacity plastic containers half filled with water recorded 98-100% survival rate between 80-110 min transportation time. However, live transportation of the juveniles in 50 L capacity plastic containers without water recorded survival rates ranging from 20-66% while those carried with water and aerated recorded 12.5-93.75%. An air conditioned vehicle was used during the live transport. Higher mortality rates were recorded for the juveniles than the broodstock due the large numbers transported in addition to the small-sized gear used for trapping the juveniles. The study has developed a simple technology for collection and transportation of live specimens of broodstock and juveniles of *M. vollenhovenii*. Efforts will be made to increase survival of the juveniles through use of larger containers and gears.

Key words: Transportation, broodstock, juveniles, *Macrobrachium vollenhovenii*

INTRODUCTION

The giant African River Prawn *M. vollenhoveni* is commonly found in fresh water which includes lakes, rivers, swamps, canals, ponds (and most cases those that have access to brackish water habitat) as well as in estuarine areas. The adults are available all year round while, the juveniles occur during the peak heavy rainy season. The species is hardy as it can thrive in murky waters and even survive in water with very low dissolved oxygen. These attributes make the species a good candidate for aquaculture purposes. In West Africa, small seasonal commercial fisheries exist for the capture of *M. vollenhovenii* and the main target species for fisheries in this area is the relatively large fresh water prawn (Willfuhr-Nast *et al.*, 1993; Abohweyere, 2008).

In Nigeria, *M. vollenhovenii* has been found to have high commercial potential (Ajuzie and Fagade, 1992; Abowei *et al.*, 2006; Anyanwu *et al.*, 2007). However, culture of the species is not well developed as there are very few prawn farms in the country. To kick start prawn farming in Nigeria, broodstock and juveniles can be collected from the wild prior to development and

domestication of technology for hatchery production and grow-out. The rearing of organisms through recruitment from the wild is an age long practice. This culture mode could be used to ascertain the propagation potential of the specie. The collection from the wild usually involves selection of berried females and mature males. Once collected, they are carefully transported live to a hatchery where they can be quarantined and acclimatized to the ambient environment. This could serve as a prerequisite for culture of the prawns as natural systems of rearing *Macrobrachium* species have been carried out in Asia and other parts of the world with great success (Ayoola *et al.*, 2009).

Broodstock and juveniles of *M. vollehovenii* are abundant in the fresh water environment of Nigeria especially in the coastal waters (Oluayo, 2014; Deekae and Abowei, 2010; Olele *et al.*, 2012). Location of sites for collection and standardizing the technology for collection and live prawn transportation will kick start the culture of the species in Nigeria, diversify protein source for the populace, create employment opportunities and generate wealth. The aim of this study was to locate collection sites of broodstock and juveniles *M. vollehovenii* and standardize technology for their live transportation.

MATERIALS AND METHODS

Area of collection: Adult berried females, mature males and juveniles of *M. vollehovenii* were collected from Ogun River at Isheri Olofin area and Okunmanya River (Fig. 1) in Ogun State. The broodstock were caught with woven cane traps, a non return valve trap as shown in Fig. 2 while the juveniles were caught with mosquito net trap (Fig. 3).

Acclimatization: The prawns were transported in 50 L plastic containers with or without water in an air conditioned car to NIOMR shrimp hatchery. On arrival at the hatchery, the prawns were treated with 5 ppm formalin for disinfection before they were transferred to quarantine tanks and acclimatized to the new environment. The water temperature, pH and ammonia of the transport water before and after transport were also taken. The total length in centimeters and body weight



Fig. 1: Fisherfolk returning with harvested prawns



Fig. 2: Trap for catching adult prawns *Macrobrachium vollenhovenii*



Fig. 3: Specially designed traps for harvesting live prawn juveniles from Ogun river

in grammes of samples of the broodstock and juveniles collected were also taken and recorded. The time taken during transportation and survival rates for each batch of prawns transported was taken.

RESULTS

A total of 457 broodstock of *M. vollenhovenii* were collected and stocked at different stages of maturation while 143,000 specimens of juveniles were also collected in the period under review. The morphometric measurement of samples of the broodstock collected showed that, body weight ranged from 71.0-264.5 g, total length 26.8-35.4 cm and carapace length 8.5-14.2 cm. The abdominal length ranged from 10.0-18.0 cm while, telson length ranged from 2.9-5.0 cm. The number of



Fig. 4: Harveasted live prawn juveniles

Table 1: Collection of broodstock of *Macrobrachium vollenhovenii* from the wild

Date	No. of broodstock collected	Mode of transportation	No. of hours of trip	Survival (%)
20/03/2012	50	Aerated in air-conditioned car	1 h, 30 min	100
11/04/2012	55	-	1 h, 45 min	100
15/05/2012	88	-	1 h, 25 min	100
27/08/2012	50	-	1 h, 35 min	100
05/10/2012	109	-	1 h, 39 min	98
12/10/2012	15	-	1 h, 45 min	100
16/10/2012	29	-	1 h, 30 min	100
30/10/2012	18	-	1 h, 35 min	100
08/11/2012	8	-	1 h, 50 min	100
29/11/2012	35	-	1 h, 40 min	100
Total	457			

broodstock collected was higher between March and October with a total catch of 352 specimens while, fewer numbers were collected between mid October and November with a total catch of 105 specimens.

The method of transportation of live broodstock using aeration in 50 L plastic containers with aeration and water was successful with records of 98-100% survival rates. This was carried out at a transportation time between 1 h, 20 min to 1 h, 50 min (Table 1). The water quality parameters of the transport water for the broodstock and juveniles before and after transport are presented in Table 2. The values were within the tolerance range for the species.

For the juveniles, a total of 143,000 specimens of *M. vollenhovenii* were collected from the various stations and transported to the institutes shrimp hatchery at times between 2 h, 10 min and 2 h, 45 mins. Live transportation of the juveniles in 50 L capacity plastic containers without water (Fig. 4) recorded survival rates ranging from 20-66% while those carried with water and aerated recorded 12.5-93.75%. The juveniles were stocked in earthen ponds for grow-out to table size (Fig. 5). The morphometric measurement of samples of the juveniles collected showed that,



Fig. 5: Stocking of juveniles in ponds

Table 2: Water quality parameter of transport water for broodstock and juveniles of *Macrobrachium vollenhovenii*

Parameters	Values before transport	Values after transport
Broodstock		
pH	7.2	7.3
Water temperature (°C)	26.1	27.0
Ammonia (mg L ⁻¹)	0.0	1.2
Nitrite (mg L ⁻¹)	0.0	0.1
Alkalinity (mg L ⁻¹)	52.0	40.0
Salinity (ppt)	0.0	0.0
Juveniles		
pH	7.0	6.8
Water temperature (°C)	28.0	27.5
Ammonia (mg L ⁻¹)	0.0	2.0
Nitrite (mg L ⁻¹)	0.0	0.2
Alkalinity (mg L ⁻¹)	56.0	34.2
Salinity (mg L ⁻¹)	0.0	0.0

body weight ranged from 0.1-0.8 g, total length 1.5-4.2 cm and carapace length 0.5-1.7 cm. The abdominal length ranged from 0.6-1.8 cm while telson length ranged from 0.2-0.6 cm. Details are presented in Table 3.

DISCUSSION

The presence of broodstock and juveniles of *Macrobrachium vollenhovenii* in the study area is a testament to their viability and preponderance. The 457 broodstock and the 143,000 juveniles that were collected during the study at peak rainy season. This is in consonance with studies done by Nwosu and Wolfi (2006), Abohweyere (2008) and Olele *et al.* (2012). The broodstock and juveniles were caught with cones haped bamboo baskets (non return valve traps) as described by Solarin *et al.* (2003) and Jimoh *et al.* (2011).

Table 3: Collection of *Macrobrachium vollenhovenii* juveniles from the wild

Date	Estimated No. of juveniles collected	Mode of transportation	No. of hours of trip	Survival (%)	No. of live juveniles
25/09/2012	20,000	In A/C car without water	2 h, 45 min	20.0	4,000
28/09/2120	8,000	-	2 h, 20 min	62.5	5,000
12/10/2012	3,000	-	2 h, 30 min	66.0	2,000
16/10/2012	8,000	Aerated plastic bags	2 h, 30 min	93.75	7,500
19/10/2012	48,000	-	2 h, 25 min	12.5	6,000
23/10/2012	30,000	-	2 h, 20 min	33.3	10,000
30/10/2012	6,000	-	2 h, 10 min	83.3	5,000
08/11/2012	20,000	-	2 h, 25 min	40.0	8,000
Total	143,000	-			47,500

The body weight of the broodstock caught ranged from 71.0g-264.5 g. This is much larger than those caught in other studies done by Oluayo (2014) and Lawal-Are and Owolabi (2012). The broodstock caught in this work represented total lengths ranging from 26.8 and 35.4 cm; this is at variance with the study of Marioghae and Ayinla (1995) and Olele *et al.* (2012) in other water bodies.

The sex ratio of males and females of 1:4 representing 91 males and 366 females in this study is accordance with the findings of (Olele *et al.*, 2012) who posited that there were more gravid females than males of *Macrobrachium vollenhovenii* that were caught during peak rainy season. Also, Nwosu and Wolfi (2006) reported that the species are available all year round during peak rainy season. George and Rao (1967) also reported similar sex differentials. On the contrary, Kingdom and Erondy (2013) in his study reported catch of more males than females. The study of Inyang (1981), Marioghae (1982) and Kavu (1985) showed that males and females are of equal ratio. Although, Tawari-Fufeyin *et al.* (2005) said that sex ratio may not always be static as they may vary from season to season and from year to year within the same population.

During transportation of the broodstock in 50 L plastic containers containing water with aeration 98-100% survival was recorded. This was achieved through travel times of approximately 1-2 h. For the juveniles, they were transported via two modes which were in 50 L containers in an air conditioned car without water and in aerated plastic bags with water. Both transportation modes showed 20-66 and 12.5-93.75% survival, respectively. The water quality parameters measured during transportation of both broodstock and juveniles of *Macrobrachium vollenhovenii* in this study were within the tolerance range for the species.

CONCLUSION

Availability of berried broodstock of *M. vollenhovenii* and the high survival rate recorded in this experiment showed that the species were abundant in the waters of Ogun State. The technology of collection and transportation developed in this study could be adopted for supply of live broodstock and juveniles. The presence of juveniles in the wild is an indication that aquaculture production of the species through wild recruitment and culture is possible. However, development of technology for successful wild collection of juveniles with increased survival rate should be addressed. The berried females stocked in the hatchery spawned easily in the plastic tanks and different developmental stages were recorded by identifying colour changes in the eggs, an indication of the possibility of hatchery propagation of the species. However, low survival of the larvae after 10 days is still a major challenge.

REFERENCES

- Abohweyere, P.O., 2008. Length-weight relationship and condition factor of *Macrobrachium vollenhovenii* in the Lagos-Lekki lagoon system, Nigeria. *Nig. J. Fish.*, 5: 98-105.
- Abowei, J.F.N., S.N Deekae, C.C Tawari, M.E. Allison and S.A. Ngodigha, 2006. A review of shrimp fishery in Nigeria. Pre-Joe Publishers, Port Harcourt, Nigeria, Pages: 53.
- Ajuzie, C.C. and S.O. Fagade, 1992. African river prawn has commercial potential. *Fish Farmer Int. File*, 6: 15-16.
- Anyanwu, P.E., B.C. Okoro, A.O. Anyanwu, M.A. Matanmi and B.I. Ebonwu *et al.*, 2007. Length-weight relationship, condition factor and sex ratio of African Mudcatfish (*Clarias gariepinus*) reared in indoor water recirculation system tanks. *Res. J. Biol. Sci.*, 2: 780-783.
- Ayoola, S.O. and A.A. Idowu and A.O. Lawal-Are, 2009. Monoculture of fresh water shrimps (*Macrobrachium* species). *Eur. J. Sci. Res.*, 29: 471-480.
- Deekae, S.N. and J.F.N. Abowei, 2010. The recruitment pattern of *Macrobrachium macrobrachion* (Herklots, 1851) from Luubara Creek, Ogoni Land, Niger Delta, Nigeria. *Res. J. Applied Sci. Eng. Technol.*, 2: 568-573.
- George, M.J. and P.V. Rao, 1967. Distribution of sex ratios of penaeid prawns in the trawl fishery off Cochin. *Proceedings of the Symposium on Crustacea: Part II*, January 12-16, 1965, MBAL, Ernakulam, India, pp: 698-700.
- Inyang, N.M., 1981. On the biology of *Macrobrachium felicinum* (Holthius) in the lower Niger River of Southern Nigeria. *Rev. Zool. Afr.*, 98: 440-449.
- Jimoh, A.A., E.O. Clarke, O.O. Whenu and H.B. Adeoye, 2011. Food and feeding habits of the African river prawn (*Macrobrachium vollenhovenii*, Herklots, 1857) in epe lagoon, Southwest Nigeria. *Int. J. Fish. Aquacult.*, 3: 10-15.
- Kavu, B.W., 1985. *Macrobrachium macrobrachium* (Herklots, 1851), its availability and effects of salinity on the growth of its juvenile. M.Sc. Thesis, Aquaculture, Rivers State University of Science and Technology, Port Harcourt, Nigeria.
- Kingdom, T. and E.S. Erondy, 2013. Reproductive biology of African river prawn *Macrobrachium vollenhovenii* (Crustacea, Palaemonidae) in the lower Taylor creek, Niger Delta, Nigeria. *Ecol. Balkanika*, 5: 49-56.
- Lawal-Are, A.O. and A.T. Owolabi, 2012. Comparative biology of the prawns *Macrobrachium macrobrachion* (Herklots) and *Macrobrachium vollenhovenii* (Herklots) from two interconnecting fresh/brackish water Lagoons in South-West Nigeria. *J. Mar. Sci. Res. Dev.*, Vol. 2. 10.4172/2155-9910.1000108
- Marioghae, I.E. and O.A. Ayinla, 1995. The reproductive biology and culture of *Macrobrachium vollenhovenii* (Herklots, 1857) and *macrobrachium macrobrachion* (Herklots, 1851) in Nigeria. NIOMAR Technical Paper No. 100, African Regional Aquaculture Centre/Nigerian Institute for Oceanography and Marine Research, Port Harcourt, Nigeria.
- Marioghae, I.E., 1982. Notes on the biology and distribution of *Macrobrachium vollenhovenii* and *Macrobrachion* in the Lagos Lagoon (Crustacea, Decapoda, Palamonidae). *Rev. Zool. Afr.*, 96: 493-508.
- Nwosu, F.M. and M. Wolfi, 2006. Population dynamics of the giant African river prawn *Macrobrachium vollenhovenii* Herklots 1857 (Crustacea, palaemonidae) in the cross river estuary, Nigeria. *West Afr. J. Applied Ecol.*, 9: 1-18.

- Olele, N.F., P. Tawari-Fufeyin and J.C. Okonkwo, 2012. Reproductive biology of fresh water prawn *Macrobrachium vollenhovenii* (Herklot, 1857) caught in Warri River. Arch. Zootechnica, 15: 41-57.
- Oluayo, B.O., 2014. Some aspects of the reproductive biology of freshwater prawn (*Macrobrachium vollenhovenii*, Herklot 1857). Federal University of Technology, Akure. <http://datad.aau.org:8080/jspui/handle/123456789/22483>.
- Solarin, B.B., R.E.K. Udolisa, N.O. Omotoyo, P.E. Lebo and E. Ambrose, 2003. Hook, line and sinker: The small scale fishing gear in Nigeria. ICSF Chennai India/Brussels Belgium. Samudra, 35: 41-46.
- Tawari-Fufeyin, P., S.A. Ekaye and U. Ogigirigi, 2005. Contribution to the biology of *Chrysichthys nigrodigitatus* (1803) in Ikpoba River, Benin City Nigeria. J. Agric. For. Fish., 6: 19-23.
- Willfuhr-Nast, J., H. Rosenthal, P.J. Udo and N. Fredrich, 1993. Laboratory cultivation and experimental studies of salinity effects on the larval development in the African river prawn *Macrobrachium vollenhovenii* (Decapoda, *Palaemonidae*). Aquat. Living Resour., 6: 115-137.