



Journal of
**Fisheries and
Aquatic Science**

ISSN 1816-4927



Academic
Journals Inc.

www.academicjournals.com



Research Article

Levels of Available Nitrogen-Phosphorus Before and After Fish Mass Mortality in Maninjau Lake of Indonesia

¹Hafrijal Syandri, ¹Azrita, ¹Junaidi and ²Ainul Mardiah

¹Department of Aquaculture, Faculty of Fisheries and Marine Science, Bung Hatta University, Padang, Indonesia

²Department of Aquaculture, Faculty of Marine and Fisheries Science, Nahdlatul Ulama University of West Sumatera, Padang, Indonesia

Abstract

Background and Objective: Maninjau lake is one of the important locations for aquaculture activity for many local people in Indonesia. The objective of the current research was to estimate the Nitrogen (N), Phosphorus (P) and Total Organic Matter (TOM) before and after fish mass mortality in Maninjau lake. **Materials and Methods:** This research was conducted in February, 2017, four months after mass mortality occurred. Nile tilapia (*Oreochromis niloticus*) and Common carp (*Cyprinus carpio*) mortality were occurred in August and September, 2016. Data were collected from four stations on Maninjau lake (Muko-Muko, Pasa, Pandan and Sungai Tampang). Water samples were taken from the surface (depth 0.1 m) and under floating net cages (depth 30 m) at each station and analyzed for N, P and TOM content. Water quality data taken in February, 2016 was used as a comparison. The differences between N, P, TOM, Particulate Organic Matter (POM) and Dissolved Organic Matter (DOM) levels before and after fish mass mortality were analyzed using a student t-test. Any differences between stations were analyzed using one-way ANOVA was performed using SPSS computer software. **Results:** The levels of N and P before and after fish mass mortality were significantly different ($p < 0.05$). The N levels at surface ranged from 1.83-2.30 mg L⁻¹. At 30 m, N levels ranged from 2.11-2.60 mg L⁻¹. The P levels ranged from 0.50-0.91 mg L⁻¹ and 0.81-0.92 mg L⁻¹ at 0.1 and 30 m depths, respectively. The N level tended to be the limiting factor for algae growth at each station (all N/P < 16). The TOM levels before and after fish mass mortality ranged from (Mean \pm SD) 4.55 \pm 0.02-16.33 \pm 0.01 mg L⁻¹ and 6.97 \pm 0.72-19.04 \pm 0.04 mg L⁻¹ at 0.10 and 30 m depths, respectively. **Conclusion:** The availability of N, P and TOM in the water was significantly higher ($p < 0.05$) after fish mass mortality and had a negative effect on the water quality of Maninjau lake.

Key words: Aquaculture, fish mass mortality, depth water, floating net cages, total organic matter

Received: March 22, 2017

Accepted: May 01, 2017

Published: June 15, 2017

Citation: Hafrijal Syandri, Azrita, Junaidi and Ainul Mardiah, 2017. Levels of available nitrogen-phosphorus before and after fish mass mortality in Maninjau lake of Indonesia. J. Fish. Aquat. Sci., 12: 191-196.

Corresponding Author: Hafrijal Syandri, Laboratory of Aquaculture, Faculty of Fisheries and Marine Science, Bung Hatta University, Padang, Indonesia
Tel: +62751-7051678

Copyright: © 2017 Hafrijal Syandri *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Nitrogen (N) and phosphorus (P) are very important elements in water bodies because they serve as a source of nutrients for plant-based organisms¹⁻³. In land water quality criteria are determined by the levels of N and P⁴. Increases in the levels of N and P in water bodies are largely related to aquaculture activity involving floating net cages^{5,6}. Although increase in P load have made lakes more productive, the ratio of N to P is a limiting factor that can be used to analyze the growth of phytoplankton^{7,8}. Compared to other micronutrients P plays an important role in the metabolism of lake biota⁹. The availability of P in water bodies (e.g., from PO₄) can often be used directly by vegetative components².

The surface area of Maninjau lake is 99.5 km²¹⁰. The aquaculture activity of *Oreochromis niloticus* and *Cyprinus carpio* by floating net cages is very intensive^{6,11}. In 2015, the total number of floating net cages at Maninjau lake was 20,608 units and the dominant species cultured was *O. niloticus*¹². The aquaculture activity related to *O. niloticus* and *C. carpio* has the potential to release significant amounts of N and P into water bodies^{2,13}, mainly from fish feed⁶. As a result, increasing aquaculture activity related to these fish species may lead to eutrophication and aquatic ecosystem changes^{3,14-17}.

Determining the availability of N, P and total organic matter in Maninjau lake is very important considering the intensive level of fish aquaculture activity involving floating net cages. From August-September, 2016, as much as 600 t of fish drowned in Maninjau lake¹⁸. Briefly, the fish mass mortality was occurred due to upwelling condition in Maninjau lake. The aim of the current study was to analyze the availability of N, P and TOM before and after fish mass mortality in Maninjau lake in order to increase the knowledge regarding the effects of this even on water quality.

MATERIALS AND METHODS

Sampling location: The present research was conducted in February, 2017, four months after mass mortality of *O. niloticus* and *C. carpio* occurred in Maninjau lake, West Sumatera, Indonesia. Fish mass mortality occurred in August and September, 2016. The N, P, TOM, Particulate Organic Matter (POM) and Dissolved Organic Matter (DOM) levels recorded in February, 2016, were compared with data take in February, 2017. The data were collected from four stations: Muko-Muko, Pasa, Pandan and Sungai Tampang. Water samples (250 mL) were taken at the surface (depth

0.10 m) and under floating net cages (depth 30 m) at each station using a Kemmerer Water Sampler (Wildco, USA) with diameter (D) 118 mm and sample volume 1200 mL. The sampler consists of a metal tube with stopper on each end that can be held open when the sampler is lowered by a line to a desired depth. After the stoppers close the ends of the tube, the sampler is retrieved with the desired samples of water being uncontaminated by other water. Then samples were placed in bottles and preserved with H₂SO₄.

Nitrogen and phosphorous analysis: The N and P levels were analyzed by (ultraviolet-visible) light spectrophotometry (UV 160 A. Japan) according to APHA¹⁹. The P was determined using the molybdovanadate method indicated by AOAC²⁰ at an absorbance of 400 nm. All samples were analyzed in triplicate. Redfield criteria were assessed by calculating the N/P ratio. An N/P<16 means that N is the limiting factor, while an N/P>16 means that P was the limiting factor, an N/P = 14-16 means that N and P were collectively at the limiting factor²¹.

Water quality analysis: Dissolved oxygen (O₂) levels were determined using a Yellow Spring oxygen meter model 52-Yellow Spring Instrument Co., Yellow Springs, OH, USA). The TOM, DOM and POM were analyzed from surface water samples using the titrimetric method²². The POM was calculated as the difference between TOM and DOM.

Statistical analysis: The N, P, TOM, POM and DOM levels before (February, 2016) and after (February, 2017) fish mass mortality were analyzed using a student t-test. The N, P, TOM, POM and DOM levels between stations were analyzed using one-way analysis of variance (ANOVA) followed by Tukey's HDS *post-hoc* test using SPSS software (Version 16.0 for Windows, SPSS Inc, Chicago IL)²³. All samples were performed in triplicates and data presented as Means ± Standard Deviation (SD). The treatment effects were considered to be significant at p<0.05.

RESULTS

Availability of N and P in surface water: The N and P levels in 0.10 m depth water before and after fish mortality occurred are shown in Table 1.

Availability of N and P in water under floating net cages: The levels of N and P in 30 m depth water before and after fish mass mortality occurred are presented in Table 2.

Table 1: Levels of nitrogen (N) and phosphorus (P) in Maninjau lake surface water

Stations	N (mg L ⁻¹)		P (mg L ⁻¹)	
	February, 2016	February, 2017	February, 2016	February, 2017
Muko-Muko	0.85±0.015 ^{aA}	1.83±0.015 ^{ab}	0.16±0.015 ^{aA}	0.91±0.010 ^{ab}
Pasa	1.16±0.025 ^{bA}	2.15±0.025 ^{bb}	0.23±0.030 ^{bA}	0.72±0.010 ^{bb}
Pandan	0.71±0.015 ^{cA}	2.09±0.020 ^{cb}	0.14±0.015 ^{cA}	0.50±0.020 ^{cb}
Sungai Tampang	1.13±0.030 ^{dA}	1.95±0.020 ^{db}	0.53±0.020 ^{dA}	0.74±0.010 ^{db}

Surface water samples were taken at a depth of 0.10 m, data are presented as the Mean±SD of triplicate samples, *The difference between mean with different lower case letters in a columns and the difference between means with different upper case letters for each parameter are statically significant (p<0.05)

Table 2: Levels of nitrogen (N) and phosphorus (P) in water under floating net cages in Maninjau lake

Stations	N (mg L ⁻¹)		P (mg L ⁻¹)	
	February, 2016	February, 2017	February, 2016	February, 2017
Muko-Muko	0.96±0.017 ^{aA}	2.11±0.035 ^{ab}	0.36±0.015 ^{aA}	0.72±0.010 ^{bb}
Pasa	1.23±0.015 ^{bA}	2.30±0.020 ^{bb}	0.37±0.030 ^{bA}	0.97±0.010 ^{bb}
Pandan	0.90±0.010 ^{cA}	2.60±0.030 ^{cb}	0.34±0.015 ^{cA}	0.81±0.020 ^{bb}
Sungai Tampang	1.43±0.010 ^{dA}	2.41±0.025 ^{db}	0.63±0.020 ^{dA}	0.85±0.010 ^{ab}

Surface water samples were taken at a depth of 30 m, data are presented as the Mean±SD of triplicate samples, *The difference between mean with different lower case letters in a columns and the difference between means with different upper case letters for each parameter are statically significant (p<0.05)

Table 3: Levels of TOM, DOM and POM in Maninjau lake

Stations	TOM (mg L ⁻¹)		DOM (mg L ⁻¹)		POM (mg L ⁻¹)	
	February, 2016	February, 2017	February, 2016	February, 2017	February, 2016	February, 2017
Muko-Muko	4.55±0.020 ^{aA}	6.97±0.720 ^{ab}	3.65±0.030 ^{aA}	4.56±0.036 ^{ab}	0.90±0.015 ^{aA}	2.41±0.020 ^{ab}
Pasa	9.12±0.025 ^{bA}	12.55±0.030 ^{bb}	6.58±0.035 ^{bA}	8.66±0.010 ^{bb}	2.55±0.015 ^{bA}	3.89±0.015 ^{bb}
Pandan	16.33±0.015 ^{cA}	19.04±0.035 ^{cb}	2.55±0.040 ^{cA}	3.89±0.020 ^{cb}	6.50±0.014 ^{cA}	6.82±0.035 ^{cb}
Sungai Tampang	13.59±0.035 ^{dA}	15.31±0.020 ^{db}	8.17±0.030 ^{dA}	9.48±0.015 ^{db}	5.43±0.035 ^{dA}	5.83±0.010 ^{db}

Data are presented as the Mean±SD of triplicate samples, *The difference between mean with different lower case letters in a columns and the difference between means with different upper case letters for each parameter are statistically significant (p<0.05), *TOM: Total organic matter, DOM: Dissolved organic matter, POM: Particulate organic matter




Fig. 1: Red field of nitrogen/phosphorus ratio (m mol⁻¹) in Maninjau lake before and after fish mass mortality

Red field criteria of the N/P ratio: The N/P values reported in February, 2016 ranged from 2.13±0.68-5.31±0.83, whereas those calculated here in (February, 2017) ranged from 2.01±0.42-4.18±0.58 (Fig. 1). Interestingly all sampling stations had an N/P<16.

TOM, DOM and POM levels before and after fish mass mortality: The levels of TOM, DOM and POM before and after fish mass mortality in Maninjau lake are presented in Table 3.

DISCUSSION

The current research showed the levels of N and P at 0.10 and 30 m depths significantly increased (p<0.05) after fish mass mortality (February, 2017). This increase was the result of as much as 600 t of *O. niloticus* and *C. carpio* drowning and sinking to the bottom of Maninjau lake. Previously, Asir and Pulatsu²⁴ reported that each ton of dead Rainbow trout (*Oncorhynchus mykiss*) released as much as 56.00 kg of N and 10.66 kg of P into water bodies. Other researchers have found similar result regarding *O. mykiss* production reporting 43.9 and 8.8 kg t⁻¹ of N and P, respectively¹⁶.

The N and P are end-products of fish load that can affect water quality in fish farming areas⁹. While the concentration of N and P in water can be estimated from the total amount of feed given to the fish²⁴, it is possible that increases in N and P may also result from fish mass mortality.

For example, the main end-product of protein metabolism in teleost fish is ammonia (NH_3) and most nitrogenous wastes from same fish species are excreted as urea²⁵. Riche and Brown²⁶ also reported that the end-products of protein catabolism in fish were dissolved NH_3 and urea. Meanwhile, the amount of N and P in fish feed and changing feed conversion ratios have been influenced the amount and types of nutrients excreted into the aquatic environment²⁷. Furthermore, Mallekh *et al.*²⁸ found that the fish size, species and feed type affect the amount of digestive residue.

The levels of N and P at 0.10 and 30 m depths after fish mass mortality in Maninjau lake was significantly different between water sampling stations ($p < 0.05$). The level of N ranged from 1.83-2.30 mg L^{-1} at 0.10 m and 2.11-2.60 mg L^{-1} at 30 m, while, P levels ranged from 0.50-0.91 mg L^{-1} at 0.10 m and 0.81-0.92 mg L^{-1} at 30 m. Fish mass mortality have an effect to N and P levels in Maninjau lake. The dead fish was decomposed by bacteria which can increase the level of N and P. The level of N and P at the surface and 30 m depth of Maninjau lake tend to higher than before fish mass mortality. In Toba lake, Indonesia, the N and P levels were 0,013-0,457 mg L^{-1} and 0,005-0.116 mg L^{-1} , lower compared than Maninjau lake. This condition was due to fish mass mortality occurred in this place¹. The N and P levels at Maninjau lake based on the regulations of the Ministry of Environment of the Republic of Indonesia Number 28/2009²⁹. The water status of Maninjau lake refer to hypereutrophic. Moreover, the levels of N and P at the bottom of Maninjau lake (under floating net cages, 30 m depth) was higher than at the surface (0.10 m depth). The P containing nutrients settle and accumulated in deeper parts of the lake most likely when orthophosphate in the epilimnion zone undergoes co-precipitation and absorption with particulates or macro metals such as iron, manganese, aluminum and/or other organic compounds¹.

The amount of fish mass mortality observed in Maninjau lake was correlated with the number floating net cages at each station. The number of floating net cages was highest at Pasa (133 units/10,000 m^2) and the lowest at Muko-Muko (69 units/10,000 m^2). The N levels at 0.10 and 30 m depths at Pasa in February, 2017 were 2.15 ± 0.025 and 2.30 ± 0.020 mg L^{-1} , respectively. While, the level of P was 0.72 ± 0.01 and 0.97 ± 0.01 mg L^{-1} , respectively. Furthermore, the productivity of Maninjau lake is relatively high as indicated by the brightness of water, which ranges from 1.30-1.80 m. Compared to that of Toba lake which is lower water brightness 7.0-15.0 m^1 .

In general, orthophosphate is a P containing nutrient whose availability for growth of phytoplankton. The

orthophosphate level at the surface of Maninjau lake was relatively high, ranging from 0.055-0.125 mg L^{-1} . The nutrient element of P is highly dependent on the dissolved O_2 concentration. The levels of dissolved O_2 at 0.10 m at each station after fish mass mortality were 5.89, 5.93, 6.17 and 5.85 mg L^{-1} , respectively. However, the dissolved O_2 levels at 30 m were 4.70, 4.92, 5.01 and 4.82 mg L^{-1} , respectively. Largely an aerobic conditions are may be found at depths greater than 50 m and the average depth of Maninjau lake is 105 m^{10} . More orthophosphate is found in particle form of dissolved O_2 decreases below 50 m. According to Nomosatryo and Lukman¹ the accumulation of P at the bottom of Toba lake is related to higher levels of orthophosphate at the water's surface. Moreover, Kelly³⁰ stated that orthophosphate compounds liberated from sediment diffuse at the water's surface. The levels of N tended to be a limiting factor for algae growth in Maninjau lake because all stations had an $\text{N/P} < 16$ (Fig. 1). This result was similar to that of Lake Sentani-Indonesia $\text{N/P} 10.24^{31}$ but different from that of other Indonesia lakes. Both N and P levels were found to limit algae growth in Toba lake ($\text{N/P} 1.9-46.3$)¹, while P was the limiting factor in Panglima Besar Soedirman Reservoir ($\text{N/P} = 19.13-65.82$)⁸.

Analysis of different types of organic matter in Maninjau lake revealed that although DOM, POM and TOM levels increased. The DOM levels were dominant (Table 3). The level of TOM was higher after fish mortality, the TOM was highest at Pandan and Sungai Tampang stations. Similarly, Lukman and Hidayat³² reported the level of DOM was higher than the POM in Toba lake. Analysis of the accumulation of DOM in aquaculture activity is very important³³, because the process of sedimentation and mineralization may contribute to the decline of organic values, especially in Maninjau lake and both processes are easily to occur in lentic waters.

Interestingly the level of TOM at Pasa, Pandan and Sungai Tampang stations after fish mortality were relatively similar mean TOM 15.63 mg L^{-1} or (0.01563 kg m^{-3}), whereas, that at Muko-Muko was 6.97 mg L^{-1} (0.00697 kg m^{-3}). These difference in the level TOM also confirm that the increase in TOM found in February, 2017 were largely the result of fish mass mortality. This condition may be correlated with water retention time of lake. According to Syandri *et al.*¹⁰, Maninjau lake has an area 99.5 km^2 (75.38%) from the catchment area, has water volume 10,226,001,629.2 m^3 , average depth was 105 m, the average of outflow water was 12.86 m^3/sec and the role of groundwater is quite large as a source of water at Maninjau lake. So Maninjau lake has water retention time during 25 years.

CONCLUSION

The current research found clear evidence that mass mortality of *O. niloticus* and *C. carpio* related to floating net cages in Maninjau Lake is a significant source of N, P and TOM. The level of these parameters was significantly higher in deeper water (30 m depths) compared to that at the surface (0.10 m depths) after fish mortality. It is concluded that a major side effect of fish mass mortality is that it decreased the water quality. Fish mass mortality can be reduced by adjusting the timing and stocking density of aquaculture fish. This will help to reduce the downstream negative effects on the lake and surrounding and in turn positively effect to many local people, as improving labor conditions lead to increased economic welfare.

SIGNIFICANCE STATEMENTS

This study discovered the increasing levels of N, P and TOM after fish mass mortality in Maninjau lake. This condition has negative effect on water quality which cause hypereutrophic condition. While N level tended to as a limiting factor for algae growth at Maninjau lake. This study is an emerging issue now days and require urgent action from the concerned authorities and awareness from fish farmers by adjusting rearing time and stocking density of aquaculture fish for the future.

ACKNOWLEDGMENT

This study was supported by a study grant (Riset Unggulan Perguruan Tinggi) from the Directorate of Research and Community Service of the Ministry of Research Technology and Higher Education, Republic of Indonesia, (Grant No. SP.DIPA-042.06-0/2017).

REFERENCES

1. Nomosatryo, S. and Lukman, 2011. The availability of nutrients nitrogen (N) and phosphorus (P) in Toba Lake, North Sumatra province. *Limnotek*, 18: 127-137.
2. Rahman, M.M., 2015. Effects of co-cultured common carp on nutrients and food web dynamics in rohu aquaculture ponds. *Aquacult. Environ. Interact.*, 6: 223-232.
3. Kaya, D. and S. Pulatsu, 2017. Sediment-focused environmental impact of rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792) cage farms: Almus reservoir (Tokat). *Turk. J. Fish. Aquat. Sci.*, 17: 345-352.
4. Jorgensen, S.E. and R.A. Vollenweider, 1989. Guidelines of Lake Management, Volume 1: Principles of Lake Management. 1st Edn., International Lake Environment Foundation, Shiga, Japan, ISBN: 4-906356-27-3, Pages: 197.
5. Junaidi, H. Syandri and Azrita, 2014. Loading and distribution of organic materials in Maninjau lake West Sumatra province-Indonesia. *J. Aquacult. Res. Dev.*, Vol. 5. 10.4172/2155-9546.1000278.
6. Syandri, H., Azrita and Niagara, 2016. Trophic status and load capacity of water pollution waste fish culture with floating net cages in Maninjau lake, Indonesia. *Ecol. Environ. Conserv.*, 22: 459-466.
7. Downing, J.A. and E. McCauley, 1992. The nitrogen: Phosphorus relationship in lakes. *Limnol. Oceanogr.*, 37: 936-945.
8. Widyastuti, E., Sukanto and N. Setyaningrum, 2015. Effect of organic waste on trophic status, N/P ratio and phytoplankton abundance in Waduk Besar Panglima Soedirman. *Biosfera*, 32: 35-41.
9. Lazzari, R. and B. Baldisserotto, 2008. Nitrogen and phosphorus waste in fish farming. *Bolet. Inst. Pesca*, 34: 591-600.
10. Syandri, H., Junaidi, Azrita and T. Yunus, 2014. State of aquatic resources Maninjau lake west Sumatra province, Indonesia. *J. Ecol. Environ. Sci.*, 5: 109-113.
11. Syandri, H., Elfiondri, Junaidi and Azrita, 2015. Social status of the fish-farmers of floating-net-cages in lake Maninjau, Indonesia. *J. Aquacult. Res. Dev.*, Vol. 7. 10.4172/2155-9546.1000391.
12. Syandri, H., Elfiondri, A. Mardiah and Azrita, 2016. Social status of Nile tilapia hatchery fish-farmers at Maninjau lake areas, Indonesia. *J. Fish. Aquat. Sci.*, 11: 411-417.
13. Abou, Y., A. Saidou, D. Mama, E.D. Fiogbe and J.C. Micha, 2012. Evaluation of nitrogen and phosphorus wastes produced by Nile tilapia (*Oreochromis niloticus* L.) fed *Azolla*-diets in earthen ponds. *J. Environ. Protect.*, 3: 502-507.
14. Prathumchai, N., C. Polprasert and A.J. Englande, 2016. Phosphorus leakage from fisheries sector-A case study in Thailand. *Environ. Pollut.*, 219: 967-975.
15. Dodds, W.K. and V.H. Smith, 2016. Nitrogen, phosphorus and eutrophication in streams. *Inland Waters*, 6: 155-164.
16. Kocer, M.A.T., M. Kanyilmaz, A. Yilayaz and H. Sevgili, 2013. Waste loading into a regulated stream from land-based trout farms. *Aquacult. Environ. Interact.*, 3: 187-195.
17. Herbeck, L.S., D. Unger, Y. Wu and T.C. Jennerjahn, 2013. Effluent, nutrient and organic matter export from shrimp and fish ponds causing eutrophication in coastal and back-reef waters of NE Hainan, tropical China. *Cont. Shelf Res.*, 57: 92-104.
18. DFMAR., 2017. The fishery statistic of Agam Regency, West Sumatera province of Indonesia. Department of Fisheries and Marine Agam Regency, Indonesia, pp: 87.

19. APHA., 1995. Standard Methods for Examination of Water and Wastewater. 19th Edn., American Public Health Association, Washington DC., USA.
20. AOAC., 1980. Official Methods of Analysis. 13th Edn., Association of Official Analytical Chemists, Washington, DC., USA.
21. Koerselman, W. and A.F.M. Meuleman, 1996. The vegetation N:P ratio: A new model tool to detect the nature of nutrient limitation. *J. Applied Ecol.*, 33: 1441-1450.
22. Greenberg, A.E., L.S. Clesceri and A.D. Eaton, 1992. Standard Methods for the Examination of Water and Wastewater. 18th Edn., American Public Health Association, Washington, DC., USA.
23. Tukey, M., 1949. Comparing individual means in the analysis of variance. *Biometrics*, 5: 99-114.
24. Asir, U. and S. Pulatsu, 2008. Estimation of the nitrogen-phosphorus load caused by rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792) Cage-Culture farms in Kesikkopru Dam Lake: A comparison of pelleted and extruded feed. *Turk. J. Vet. Anim. Sci.*, 32: 417-422.
25. Wood, C.M., 1993. Ammonia and Urea Metabolism and Excretion. In: *The Physiology of Fishes*, Evans, D.H. (Ed.). CRC Press, Boca Raton, FL., pp: 379-425.
26. Riche, M. and P.B. Brown, 1999. Incorporation of plant protein feedstuffs into fish meal diets for rainbow trout increases phosphorus availability. *Aquacult. Nutr.*, 5: 101-105.
27. Rodehutschord, M., S. Mandel and E. Pfeffer, 1994. Reduced protein content and use of wheat gluten in diets for rainbow trout: Effects on water loading with N and P. *J. Applied Ichthyol.*, 10: 271-273.
28. Mallekh, R., T. Boujard and J.P. Lagardere, 1999. Evaluation of retention and environmental discharge of nitrogen and phosphorus by farmed turbot (*Scophthalmus maximus*). *N. Am. J. Aquacult.*, 61: 141-145.
29. MERI., 2009. Regulations of the Ministry of Environment of the Republic of Indonesia number 28/2009 about the carrying capacity of lake water pollution and/or reservoir. Ministry of Environment of the Republic of Indonesia.
30. Kelly, L.A., 1992. Dissolved reactive phosphorus release from sediments beneath a freshwater cage aquaculture development in West Scotland. *Hydrobiologia* 235: 569-572.
31. Indrayani, E., K.H. Nitimulyo, Hadisusanto and Rustadi, 2015. Analysis of nitrogen, phosphorus and organic carbon content at Lake Sentani-Papua. *J. Manusia Lingkungan*, 2: 217-225.
32. Lukman and Hidayat, 2002. Loading and distribution of organic materials in Cirata reservoir. *J. Teknol. Lingkungan*, 3: 129-135.
33. Hambly, A.C., E. Arvin, L.F. Pedersen, P.B. Pedersen, B. Seredynska-Sobecka and C.A. Stedmon, 2015. Characterising organic matter in recirculating aquaculture systems with fluorescence EEM spectroscopy. *Water Res.*, 83: 112-120.