



Journal of
Plant Sciences

ISSN 1816-4951



Academic
Journals Inc.

www.academicjournals.com

Partial Purification of Polygalacturonase from Tomato Fruits Infected by *Rhizopus arrhizus* Fisher

A.A. Ajayi, A.O. Adejuwon and P.O. Olutiola
Department of Microbiology, Obafemi Awolowo University,
Ile Ife, Osun state, Nigeria

Abstract: The production of polygalacturonase during the deterioration of tomato (*Lycopersicon esculentum* Mill.) by *Rhizopus arrhizus* Fisher was investigated. The enzyme was partially purified by a combination of ammonium sulphate precipitation, gel filtration and ion-exchange chromatography. Two peaks of absorption, with molecular weight estimates of approximately 166 000 Daltons and 60 260 Daltons were obtained.

Key words: Polygalacturonase, tomato fruits, *Rhizopus arrhizus*, gel filtration, ion-exchange chromatography

INTRODUCTION

One of the essential substances primarily found in tomato fruits is lycopene. This substance, apart from the fact that it gives tomatoes their vibrant color, it is of health value as it helps to protect against diseases such as cancer and heart diseases (Rao *et al.*, 1999). Audisio *et al.* (1993) reported that tomato is very rich in vitamin C and this serves as a good index in estimating the freshness of the product. However, a large percentage of the tomato fruits produced in Nigeria is lost to post harvest infections caused by microorganisms. Fungi are most of these microorganisms (Ajayi *et al.*, 2003).

Rhizopus arrhizus is a known pathogenic fungus (Stevens, 1974; Orabi *et al.*, 1999). It establishes itself in its host by secreting extracellular enzymes which break down host tissues (Wheeler, 1975; Virto *et al.*, 1999; Elibol and Dursun, 2002). Martak *et al.* (2003) reported the semi continuous fungal fermentation of lactic acid by *Rhizopus arrhizus*. The ability of pathogenic fungi to produce enzymes responsible for the degradation of host tissues is one of the major factors influencing their virulence (Agrios, 1978; Lehtinen, 1993). Pectinases are a group of cell wall degrading enzymes produced by many phytopathogens (Hagar and McIntyre, 1972; Bruno *et al.*, 2001). Pectinases are enzymes useful in the degradation of such host tissues (Alabi and Naqvi, 1977; Walton and Cervone, 1990).

Very little has been reported about the involvement of *Rhizopus arrhizus* in the deterioration of tomato fruits in Nigeria. This study therefore describes the production of polygalacturonase, a pectic enzyme, during the infection of tomato fruits by *Rhizopus arrhizus*. Attempts were made to determine the molecular weight of the enzyme during purification. It is believed that this study would be very useful in providing solutions to the problems of tomato wastage in our markets.

MATERIALS AND METHODS

The experiment described in this paper was carried out between December, 1999 and January, 2005 in the Department of Microbiology, Obafemi Awolowo University, Ile Ife, Osun state, Nigeria.

Corresponding Author: A.A., Ajayi, Department of Microbiology, Obafemi Awolowo University,
Ile Ife, Osun state Nigeria

Organism

The isolate of *Rhizopus arrhizus* Fisher employed for this research work was isolated from tomato fruits in the Department of Microbiology, Obafemi Awolowo University, Ile Ife, Osun state, Nigeria. The organism was routinely grown and maintained on 1% (w/v) Sabouraud dextrose agar slants. The organism was subcultured from the old culture onto fresh Sabouraud dextrose agar plates whenever it was to be used. Seventy-two-hour-old culture was used as inoculum for this research.

Inoculation and Cultivation

The inoculation techniques employed were as earlier described (Ajayi *et al.*, 2003). The medium used in this case was Sabouraud dextrose agar. The experimental and control tomato fruits were placed individually in sterile Petri dishes under surface sterilized bell jars. Incubation was at room temperature. The fruits were examined daily for deterioration.

Extraction of Enzyme from Tomato Fruits

Ten days after inoculation, the deteriorated tomato fruits were weighed and chilled for 30 min inside a freezer and homogenized with an MSE homogenizer at full speed (25 cycles per second) with chilled liquid extractant (1:1w/v) for 2 min at 30 sec. interval. The extractant was 0.01M citrate phosphate buffer, pH 4.5 containing 5 mM NaN₃ to prevent microbial contamination. The homogenate was initially allowed to percolate through four layers of sterile glass fibre. This was used as the crude enzyme.

Preparation of Enzyme for Column Chromatography

The crude enzyme preparation was dialyzed using acetylated cellophane tubing (Whitaker *et al.*, 1963). Analysis was performed using a Multiple dialyser (Pope Scientific Inc., Model 220, U.S.A) at 4°C. Proteins in the crude enzyme preparation were precipitated by adding solid ammonium sulphate (Sigma) to 90% saturation.

Fractionation of Enzyme on Sephadex G-100

The vertical glass tube chromatography column (2.5×70 cm) of Sephadex G-100 (Particle size, 40-120 μ) was prepared and calibrated as previously described (Ajayi *et al.*, 2003). Ten millimeters of the enzyme concentrate was applied to the column and eluted with 0.05 M citrate phosphate buffer (pH 4.5). Each of the fractions was analyzed for polygalacturonase activity.

Fractionation by Ion-Exchange Chromatography

Fractions from the Sephadex G-100 column which showed appreciable polygalacturonase activity were pooled. Ten milliliter of the pooled enzyme was applied to a CM Sephadex C-50 column (2.5×40 cm) which was prepared as described above for Sephadex G-100 column. Fractions were eluted with 0.05 M citrate phosphate buffer (pH 4.5) containing 0.1, 0.2, 0.4 and 0.5M gradients of NaCl. Fractions (5 mL per tube) were collected and assayed for polygalacturonase activity.

Enzyme Assay

Polygalacturonase activity was assayed according to the method described (Olutiola, 1982a). The reaction mixture was 1 mL of 0.1% (w/v) pectin (Sigma) in 0.01 M citrate phosphate buffer (pH 4.5) and 0.5 mL of the enzyme. Each control tube contained 1 mL of the substrate. The experimental and control tube were incubated in a water bath at 37°C for 3 h. The total reducing sugar was determined by the Dinitrosalicylic acid (DNSA) method (Miller, 1959; Olutiola, 1983). One unit of polygalacturonase activity was defined as the amount of enzyme which released 1 μ mole galacturonic acid per minute.

RESULTS

Gel-filtration of the dialyzed enzyme on Sephadex G-100 column gave four peaks of absorption designated D, E, F and G (Fig. 1). Components of peaks D and E exhibited polygalacturonase activity whereas peaks F and G lacked polygalacturonase activity. The molecular weights of components of peaks D and E were approximately 166,000 and 60,260 Daltons respectively. Elution of components of peak D on CM Sephadex C-50 column produced four peaks of absorption designated Da, Db, Dc and Dd (Fig. 2). Components of peaks Da and Db possessed polygalacturonase activity while the components of peaks Dc and Dd lacked polygalacturonase activity. Purification folds of approximately 30 and 14 were obtained for the components of peaks Da and Db respectively (Table 1). Components of peak E produced three peaks of absorption when separated on CM Sephadex C-50 column. These peaks were designated Ea, Eb and Ec (Fig. 3). Components of peak Ea possessed polygalacturonase activity with a purification fold of approximately 12 (Table 1) while the components of peaks Eb and Ec possessed no detectable polygalacturonase activity.




Fig. 1: Separation by gel filtration of proteins obtained from tomato fruits deteriorated by *Rhizopus arrhizus* and the enzymic activity of the fractions towards pectin. (O,) protein (E_{280}); ---, polygalacturonase

Table 1: Partial purification steps of polygalacturonase from tomato fruits infected by *Rhizopus arrhizus*

Fraction	Total activity (units)	Total protein (mg)	Specific activity (units mg ⁻¹ protein)	Yield (%)	Purification fold
Crude extract	1840	390.0	4.7	100.0	1.00
(NH ₄) ₂ SO ₄	1600	155.0	10.3	87.0	2.19
Sephadex G-100					
Peak D	960	18.5	51.9	52.2	11.04
Peak E	290	11.8	24.5	15.8	5.21
CM Sephadex C-50					
Peak Da	380	2.7	140.7	20.7	29.90
Peak Db	110	1.7	64.7	6.0	13.80
CM Sephadex C-50					
Peak Ea	132	2.3	57.4	7.2	12.20




Fig. 2: Separation by ion-exchange chromatography (CM Sephadex C-50) of proteins (fraction D) separated by gel filtration, Fig. 1) and enzymic activity of the fraction towards pectin. (O), protein (E₂₈₀); ---, polygalacturonase




Fig. 3: Separation by ion-exchange chromatography (CM Sephadex C-50) of proteins (fraction E) separated by gel filtration, Fig. 1) and enzymic activity of the fractions towards pectin. (O), protein (E₂₈₀); ---, polygalacturonase

Table 2: Molecular weights of polygalacturonase from other microorganisms

Species investigated	Molecular weight	References
<i>Botryodiplodia theobromae</i>	12,000 Daltons (peak Aa) 56,200 Daltons (peak Ab)	Ajayi <i>et al.</i> (2003)
<i>Aspergillus niger</i>	82 kDa (exo-PG 1)	Sakamoto <i>et al.</i> (2002)
A polygalacturonase inhibitor protein (PGIP) characterized from tomato fruit.	34 kDa	Stotz <i>et al.</i> (1994)
Ripe tomatoes	42 kDa PG 1 (Heat stable form). 42 kDa PG 2 (Heat labile form).	Fachin <i>et al.</i> (2004)
<i>Aspergillus niger</i>	Multiple form of polygalacturonase with molecular mass ranging from 30-60 kDa	Jyothi <i>et al.</i> (2005)

DISCUSSION

During the deterioration of tomato fruits by *Rhizopus arrhizus*, proteins which exhibited polygalacturonase activity were produced. However, similar extracts from uninfected tomato fruits possessed traces of polygalacturonase activity. The occurrence of appreciable quantities of the enzyme in tomato tissues infected by *Rhizopus arrhizus* strongly suggest that the enzyme is of fungal origin. It has been reported that plant cell wall polysaccharide degrading enzymes play fundamental role in host-pathogen interaction (Alana *et al.*, 1990). Previous researchers have implicated the involvement of fungi in the deterioration of tomato fruits and fruits from other plants (Adejuwon and Olutiola, 2005; Olutiola, 1982b) but there is paucity of information on the deterioration of tomato fruits by *Rhizopus arrhizus* in Nigeria.

The protein and total reducing sugars of the tomato fruits decreased with infection, corroborating research findings from previous researchers, but the molecular weights of the enzyme fractions differ from the work of previous researchers (Table 2). It has been shown that the differences in molecular weights of enzymes have been associated with a number of factors (Sevillano and Zarra, 1997).

REFERENCES

- Adejuwon, A.O. and P.O. Olutiola, 2005. Protease in tomato fruits infected by *Fusarium oxysporum*. *Plant Pathol. J.*, 4: 43-45.
- Agrios, G.N., 1978. *Plant Pathology*. Academic Press, New York, pp: 703.
- Ajayi, A.A., P.O. Olutiola and J.B. Fakunle, 2003. Studies on polygalacturonase associated with the deterioration of tomato (*Lycopersicon esculentum* Mill.) fruits infected by *Botryodiplodia theobromae* PAT. *Sci. Focus*, 5: 68-77.
- Alabi, R.O. and S.H.Z. Naqvi, 1977. Production of cellulolytic and proteolytic enzymes by *Cercospora arachnidicola*. *Trans. Br. Mycol. Soci.*, 68: 296-298.
- Alana, A., I. Alkorta, J.B. Dominquez, M.J. Liama and J.L. Serra, 1990. Pectin lyase activity in a *Penicillium italicum* strain. *Applied Environ. Microbiol.*, 56: 3755-3759.
- Audisio, M., D. Dante, A. DeCica and C. Swaci, 1993. Vitamin C content in tomatoes in relation to modality and degree of maturity. *Revista Discienza Development Alimentazione*, 22: 507-512.
- Bruno, G., D. Geristo and G.E. Harman, 2001. Interaction of ammonium, glucose and chitin regulation on the expression of cell wall degrading enzymes in *Trichodema atroviride* strain Pl. *Applied Environ. Microbiol.*, 67: 5643-5647.
- Elibol, M. and O. Dursun, 2002. Response surface analysis of lipase production by freely suspended *Rizopus arrhizus*. *Process Bioche.*, 38: 367-372.
- Fachin, D., C. Smout, I. Verlent, B. Ly Nguyen, A.M. Van Loey and Hendrickx, 2004. Inactivation Kinetics of purified tomato polygalacturonase by thermal and high-pressure processing. *J. Agric. Food Chem.*, 52: 2697-2703.

- Hagar, S.S. and G.A. McIntyre, 1972. Pectic enzymes produced by *Pseudomonas fluorescens* and organism associated with "pink eye" disease of potato tubers. *Canadian J. Bot.*, 50: 2479-2488.
- Jyothi, T.C., S.A. Singh and A.G. Appu Rao, 2005. The contribution of ionic interactions to the conformational stability and function of polygalacturonase from *Aspergillus niger*. *Intl. J. Biol. Macromol.*, 36: 310-317.
- Lehtinen, U., 1993. Plant cell wall degrading enzymes of *Septoria nodorum*. *Physiol. Mol. Plant Pathol.*, 43: 121-134.
- Martak, J., S. Stefan, S. Erika, K. Ludmila and K. Michael, 2003. Fermentation of lactic acid with *Rhizopus arrhizus* in a stirred tank reactor with a periodical bleed and feed operation. *Process Biochem.*, 38: 1573-1583.
- Miller, G.L., 1959. Use of dinitrosalicylic acid reagent for the determination of reducing sugar. *Anal. Chem.*, 31: 426-432.
- Olutiola, P.O., 1982a. Extra cellular polygalacturonase complex from *Penicillium citrinum* Thom. associated with internal mouldiness of cocoa (*Theobromae cacao*) beans. *Acta Phytopathologica Academiae Scientiarum Hungaricae*, 17: 239-247.
- Olutiola, P.O., 1982b. Polygalacturonase and pectin lyase of *Penicillium sclerotigenum*. *Nig. J. Microbiol.*, 2: 154-167.
- Olutiola, P.O., 1983. Cell wall degrading enzymes associated with the deterioration of cocoa beans by *Penicillium steckii*. *International Biodeterioration Bull.*, 19: 27-36.
- Orabi, K.Y., E. Li, A.M. Clark and C.D. Huford, 1999. Microbial transformation of sampongine. *J. Natural sProducts*, 62: 988-992.
- Rao, A.V., N. Fleshner and S. Agarwal, 1999. Serum and tissue lycopene and biomarkers of oxidation in prostrate cancer patients. A case control study. *Nutrition and Cancer*, 33: 159-164.
- Sakamoto, T., E. Bonnin, B. Quemener and J.F. Thibaut, 2002. Purification and characterization of two exo-polygalacturonases from *Aspergillus niger* able to degrade xylogalacturonan and acetylated homogalacturonan. *Biochim. Biophys. Acta*, 1572: 10-18.
- Sevillano, M.J. and I. Zarra, 1997. Pectin depolymerase activities associated with cell walls from *Cicer arietinum* L. *Plant and Cell Physiol.*, 38: 1259-1263.
- Stevens, R.B., 1974. *Plant Disease*. The Ronald Press Company, New York, pp: 139-152.
- Stotz, H.U., J.J. Contos, A.L. Powell, A.B. Bennet and J.M. Labaxitch, 1994. Structure and expression of an inhibition of fungal polygalacturonases from tomato. *Plant Mol. Biol.*, 25: 607-617.
- Virto, C., S. Inzemar and P. Adbererentz, 1999. Enzymatic synthesis of lysophosphatidic acid. *Enzyme Microb. Technol.*, 24: 651-658.
- Walton, J.D. and I. Cervone, 1990. Endopolygalacturonase from the maize pathogen *Cochliobolus carbonum*. *Physiol. Mol. Plant Pathol.*, 36: 350-359.
- Wheeler, H., 1975. *Plant Pathogenesis*. Springer Verlag, New York, Heidleberg, Berlin, pp: 106.
- Whitaker, D.R., K.R. Hanson and P.K. Datta, 1963. Improved procedures for preparation and characterization of *Myrothecium cellulase*. Part 2. Purification procedures. *Can. J. Biochem. Physiol.*, 41: 671-696.