



Journal of  
**Plant Sciences**

ISSN 1816-4951



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)

## Palynological Studies on Five Species of Asteraceae

F.N. Mbagwu, E.G. Chime and C.I.N. Unamba  
Department of Plant Science and Biotechnology, Imo State University,  
Owerri, Imo State, Nigeria

**Abstract:** Palynological studies on five species of the family Asteraceae namely *Gutenbergia nigritiana*, *Emilia praetermissa*, *Vernonia guineensis*, *Lagera pterodonta* and *Chromolena odorata* was carried out. Results obtained from this investigation showed that the pollen shape is spheroidal in *G. nigritiana*, *E. praetermissa* and *C. odorata* while it is elliptic in *V. guineensis* and *L. pterodonta*. The pollen aperture is porate in all except in *L. pterodonta* where it is elliptic. The pollen wall is echinate in all except in *C. odorata* where it is smooth. The general appearance of the pollen grains is circular in *G. nigritiana*, *E. praetermissa* and *C. odorata* but longer than wide in *V. guineensis* and *L. pterodonta*. It is therefore likely that the nature of the pollen grains in these species could be an evolutionary modification often inherited to determine the mode of pollination and thereby perpetuate a particular group of plants. The similarities in structure showed interspecies relationships and reasons for them to be in the same family while the differences in structures showed reasons for them to exist as distinct species.

**Key words:** Palynomorphology, studies, pollen grains, structure, species, taxa, Asteraceae

### INTRODUCTION

Members of the Asteraceae family are herbs or shrubs, they are rarely twiners or trees. The leaves are simple, alternate or opposite, rarely compound. The inflorescence is a head or capitulum with an involucre of bracts. Flowers are of two kinds-the central ones called disc florets are tubular and the marginal ones called ray florets are ligulate (Dutta, 1974). For its many characters, Asteraceae is assigned an advanced position, the highest according to Engler, in systematic botany. Thus the family is of recent origin. Fossil records showed that the genus *Senecio* came into existence first and other genera developed from it in due course. It is likely Asteraceae and Rubiaceae have risen from a common ancestry. Asteraceae maintains a phylogenetic connection with Umbelliferae through inflorescence and floral mechanism. Asteraceae is remarkable in many respects. It has the maximum number of species, its worldwide distribution; its variety of forms and its very effective mechanism for cross-pollination (Dutta, 1974).

Pollen or flower sperm is a fine to coarse powder consisting of microgametophytes (pollen grains) which carry the male gametes of seed plants (Wagner *et al.*, 1990). Palynology is the science that studies contemporary and fossil palynomorphs, including pollen, spore, dinoflagellate cysts, acritarchs, chitinozoans and scolecodonts, together with particulate organic matter and kerogen found in sedimentary rocks and sediments (Walker, 1976). The palynological attributes of plants have attracted the attention of many scientists in recent time. For example, Nyananyo (1985), on pollen morphology of *Talineae* showed that palynology provides useful data for the intrageneric classification of the large genera. Nyananyo (1987) and Nyananyo and Olowokudejo (1986) also used seed coat morphology and other palynological features of *Talinum* and *Calandrinia* to produce a more acceptable classification of the species in these taxa.

Similarly, Edeoga *et al.* (1996, 1998), Edeoga and Gomina (2001), Nyananyo (1990) and Mbagwu and Edeoga (2006) have utilized pollen attributes to establish probable evidence of relationships among

certain groups of flowering plants in Nigeria. According to them, the main characters of taxonomic value in pollen grain are the number and position of furrows, pollen wall morphology, symmetry and shapes and sizes of pollen grains. Edeoga and Ikem (2002) also showed that *Boerhavia coccinea* is characterized by tricolpate pollen grains while *B. ejecta* and *B. diffusa* have alcopate pollen grains hence *B. coccinea* could therefore be distinguished from other Nigerian collections of *Boerhavia* based on pollen characters. The variation in shape, aperture, polar unit, symmetry and difference in wall sculpture of pollen grains have been used by many authors in the delimitation of various taxa (Agwu and Uwakwe, 1992; Agwu and Osibe, 1992; Edeoga *et al.*, 1996; Lezine and Edeoga, 1991; Angeles, 1992; Mbagwu and Edeoga, 2006).

This study is based on the hypothesis that palynological studies have played a vital role in the delimitation of various taxa but has not been used to delimit the above five species of Asteraceae to the author's knowledge hence the need for the study with the objectives of examining the pollen morphology of five species of Asteraceae and using the pollen characters for the establishment of interspecific relationships among the five species investigated.

## MATERIALS AND METHODS

Mature plants of the five species of Asteraceae namely *Gutenbergia nigriflora*, *Emilia praetermissa*, *Vernonia guineensis*, *Lagera pterodonta* and *Chromolaena odorata* were collected around the Agricultural Garden of Imo State University Owerri, Nigeria. This study was conducted in January, 2006 at the Science laboratory of the Department of Plant Science and Biotechnology of Imo State University, Owerri, Nigeria. The specimens were taken to Forestry Herbarium Ibadan (FHI) for proper identification. The voucher specimens were deposited at the Imo State University Owerri Herbarium Nigeria.

Mature flower head of each of the five species were collected and teased out on a slide. Samplings were made on fresh flowers from plants in their natural conditions as these do not undergo any form of deterioration (Edeoga *et al.*, 1996, 1998). Samples for light microscope were acetolysed (Erdtman, 1960) which involves the introduction of the acetolysed mixture comprising acetic anhydride mixed with concentrated sulphuric acid ( $H_2SO_4$ ) in the ratio of 9:1 to the centrifuged materials in the plastic test tubes. The tubes were immersed in boiling water bath for 3-5 min after which they were allowed to cool and the supernatant decanted. The residue was washed twice with water and decanted, about 5 drops of glycerine alcohol in the ratio of 2:1 was added to the precipitates and transferred into labeled specimen tubes for storage. Unstained acetolysed pollen grains were embedded in glycerine jelly and sealed with wax after covering with zero cover slip. Photomicrographs of the pollen grains were taken using a Leitz Wetzlar ortholux microscope fitted with vivitar-V-335 camera (Fig. 1).

## RESULTS AND DISCUSSION

The palynomorphological studies of the five species of Asteraceae investigated is quite different but of generalized types found among the dicotyledons (Table 1). The pollen shape is spheroidal in *Gutenbergia nigriflora*, *Emilia praetermissa* and *Chromolaena odorata* but elliptic in *Vernonia guineensis* and *Lagera pterodonta* (Fig. 1). The pollen aperture is porate in all except in *Lagera pterodonta* where it is elliptic. Polyporate number of aperture and porate type of aperture characterized all the species investigated. The pollen wall is echinate in all except in *Chromolaena odorata* where it is smooth. The general appearance of the pollen grains is circular in *G. nigriflora*, *E. praetermissa* and *C. odorata* but longer than wide in *V. guineensis* and *L. pterodonta*. The outcome of this result shows that variants with similar pollen characters are more closely related whereas those with different pollen characters are distantly related. In some of the species where the pollen grains are longer than wide, this is attributed as a structural adaptation for effective dispersal by wind while the circular nature of some of the pollen grains are related to structural adaptation for effective pollination by insects (Gimenes, 1991; Edeoga *et al.*, 1996; Mbagwu and Edeoga, 2006). From this




Fig. 1 : Pollen morphology of the five species of Asteraceae studied

- A: *Gutenbergia nigriflora* with reticulate pollen wall.
- B: *Emilia praetermissa* with polyporate aperture and echinate pollen wall.
- C: *Vernonia guineensis* with echinate pollen wall and polyporate apertural type.
- D: *Lagera pterodonta* with echinate pollen exine and polyporate aperture.
- E: *Chromolaena odorata* with smooth pollen exine

Table 1: Pollen morphology of the five species of Asteraceae studied

Pollen characters	<i>G. nigriflora</i>	<i>E. praetermissa</i>	<i>V. guineensis</i>	<i>L. pterodonta</i>	<i>C. odorata</i>
Pollen shape	Spheroidal	Spheroidal	Elliptic	Elliptic	Spheroidal
Pollen aperture	Porate	Porate	Porate	Elliptic	Porate
No. of aperture	Polyporate	Polyporate	Polyporate	Polyporate	Polyporate
Pollen wall	Echinate	Echinate	Echinate	Echinate	Smooth
Aperture type	Porate	Porate	Porate	Porate	Porate
Appearance of pollen grains	Circular	Circular	Longer than wide	Longer than wide	Circular

study, it is likely that the nature of pollen grains could be an evolutionary modification often inherited to determine the mode of pollination and thereby perpetuate a particular group of plants in a given environment (Lowe and Soladoye, 1990; Ogwal, 1990).

The differences and similarities in pollen morphology of the investigated species are significant and could be exploited for biosystematic purposes. Applying these variations in pollen morphology to the species investigated showed that species in the family with similar pollen characters are more closely related and thus exhibit interspecific relationships suggesting reasons for them to be in the same family while those with different pollen characters are not very closely related and suggest reasons for them to be as distinct species. These findings therefore are in support of the work of other scientists such as Mbagwu and Edeoga (2006) and Lowe and Soladoye (1990) etc who have used their generalizations from pollen characters to delimit various taxa.

## REFERENCES

- Agwu, C.O.C. and E.E. Osibe, 1992. Airborne palynomorphs of Nsukka during the months of February-April, 1990. *Nig. J. Bot.*, 5177-5185.
- Agwu, C.O.C. and G.O. Uwakwe, 1992. Melissopalynological studies of honey from Abia and Imo States of Nigeria. *Nig. J. Bot.*, pp: 585-591.
- Angeles, G., 1992. The periderm of flooded and non flooded *Ludwigia octovalvis* (Onagraceae). *IAWA-BULL.*, 13: 195-200.
- Dutta, A.C., 1974. Botany for Degree Students. Calcutta, Oxford University Press. Delhi, Bombay, Madras.
- Edeoga, H.O., N.O. Ogbebor and A.O. Amayo, 1996. Pollen morphology of some Nigerian species of *Aneilema* R. Br. and *Ludwigia* L. *New Bot.*, 23: 223-231.
- Edeoga, H.O., H.N. Ugbo and P.I. Osawe, 1998. Palynology of species of *Commelina* L. and *Senna* Tourn. Ex. Mill from Nigeria. *New Bot.*, 25: 1-10.
- Edeoga, H.O. and A. Gomina, 2001. Nutritional values of some non-conventional leafy vegetables of Nigeria. *J. Econ. Tax. Bot.*, 20: 7-15.
- Edeoga, H.O. and I.C. Ikem, 2002. Structural morphology of the pollen grains in three Nigerian species of *Boerhavia* L. *New Bot.*, 29: 89-95.
- Erdtman, G., 1960. The acetolysis method. *Svensk Bot. Tidskr.*, 54: 561-564.
- Gimenes, M., 1991. Some morphological adaptations in Bees (*Hymenoptera*, *Ludwigia elegans* Onagraceae). *Rev. Brasil. Entomol.*, 35: 413-422.
- Lezine, A.M. and T.M. Edoth, 1991. Modern pollen deposition in West African Sudanian environments. *Rev. Paleobot. Palyn.*, 67: 41-58.
- Lowe, J. and M.O. Soladoye, 1990. Some Changes and Corrections to Names of Nigerian Plants and Nigerian Trees since the Publication of Flora of West Tropical Africa. 2nd Edn., *Nig. J. Bot.*, 3: 1-24.
- Mbagwu, F.N. and H.O. Edeoga, 2006. Palynological studies on some Nigerian species of *Vigna savi*. *J. Biol. Sci.*, 6: 1122-1125.
- Nyananyo, B.L., 1985. Pollen Morphology in *Talineae* (Portulacaceae) *Sensu McNeill*. *Biol. Afr.*, 2: 41-46
- Nyananyo, B.L. and J.D. Olowokudejo, 1986. Taxonomic studies in the genus *Talinum* (Portulacaceae) in Nigeria. *Willdenowia Berlin*, 15: 455-463.
- Nyananyo, B.L., 1987. Seed coat morphology in *Calandrinia* (Portulacaceae) and its taxonomic significance. *J. Plant. Sci.*, 3: 93-97.
- Nyananyo, B.L., 1990. Tribal and generic relationships in Portulacaceae (Centrospermae). *Feddes Report*, 101: 237-241
- Ogwal, E.N.K., 1990. A taxonomic investigation of the genus *Commelina* in Uganda. *Proc. AETFAT, Hamburg*, 236: 573-592.
- Wagner, I.O., R.P. Woodhouse and D.R. Walters, 1990. Pollen Grains, Their Structures, Identifications and Significance in Science and Medicine. McGraw Hills. New York.
- Walker, I.W., 1976. Evolution of exine structure in the pollen of primitive Angiosperms. *Am. J. Bot.*, 61: 8-10.