



Journal of
**Pharmacology and
Toxicology**

ISSN 1816-496X



Academic
Journals Inc.

www.academicjournals.com

Effects of Receptor Polymodality on Co-transmission by Acetylcholine and Serotonin in the Afferent Neuronal Pathway of the Gut

R. Miftahof

Division of Applied Mathematics, KAIST, Daejeon, Republic of Korea

Abstract: Electrophysiological mechanisms of co-transmission by serotonin (5-HT) and acetylcholine (ACh) on mechano-electrical activity of the gut were studied numerically. A mathematical model of the bursting primary sensory (AH) and motor (S) neurons linked in sequence and smooth muscle syncytium mimicked the afferent pathway of the enteric nervous plexus of the organ. The role of different receptor types, i.e., 5-HT type 3 and 4, nicotinic (nACh) and muscarinic cholinergic (μ ACh) and the effects of selective and non-selective receptor agonists/antagonists on the dynamics of nerve signal transduction and mechanical response in the tissue were analyzed. Results showed that selective stimulation of the 5-HT₃ receptors by endogenous 5-HT reduced the threshold of activation of the mechanoreceptors by 17.6%. Conjoint excitation by serotonin of the 5-HT₃ and 5-HT₄ receptors located on the primary sensory (AH) and motor (S) neurons converted their regular firing pattern of electrical discharges to a beating mode. Activation confined to 5-HT₃ receptors located on the somas of the AH and S type neurons, could not sustain normal electrical signal transduction between them. It required acetylcholine as a co-transmitter and a subsequent co-activation of the nACh receptors. Selective 5-HT₃ receptor antagonists, e.g., *Ondansetron* and *Granisetron*, increased the threshold activation of the mechanoreceptors and inhibited dose-dependently the production of action potentials by AH neurons. 5-HT₄ receptor agonists, e.g., TS-591, prucalopride and ML10302, applied alone onto the longitudinal smooth muscle of the gut did not have any effects on its electromechanical activity. However, excitation of the 5-HT₄ in conjunction with μ ACh receptors evoked an increase in intensity of the electromechanical activity of the syncytium. GR113808A, a selective 5-HT₄ antagonist, acting alone strongly inhibited smooth muscle contractions but its effect was overcome through the activation of the 5-HT₃, nACh and μ ACh receptors. A non-selective strong 5-HT₃ and weak 5-HT₄ - receptor agonists, *Cisapride*, demonstrated a prominent effect on the AH neuron with no significant changes in the electrical activity of the S neuron. *Cisapride* depolarized the soma of AH neuron with the generation of high amplitude spikes. The drug caused twitch contractions of the longitudinal smooth muscle. Comparison of the theoretical results to *in vivo* and *in vitro* experimental data indicated satisfactory qualitative and quantitative agreement. The numerical investigations helped us reveal the intrinsic mechanisms of co-transmission by ACh and 5-HT and the role of receptor polymodality at the cellular and tissue levels that could not have been elucidated using the existing experimental *in vivo* or *in vitro* methods.

Key words: Co-localization, acetylcholine, serotonin, cholinergic receptors, 5-HT₃ and 5-HT₄ receptors, afferent neuronal pathway, mathematical model

Introduction

There has been a rapid increase of experimental data on the electro-pharmacological roles of acetylcholine (ACh) and serotonin (5-HT) as neurotransmitters, the effects of their co-transmission

and multiple receptor expression, in pathogenesis of various diseases. However, our understanding of the cellular mechanisms involved in the intricate pathways of processes responsible for the plethora of biological manifestations remains unsatisfactory. First, the combined processes of co-transmission cannot be studied effectively by any of the existing experimental techniques currently available to researchers for determining intrinsic mechanisms. Secondly, traditional *in vivo* and *in vitro* experimental approaches fail to provide desired quantitative information about synaptic neurotransmission. Thus, a new technique is needed that amalgamates interdisciplinary data and provides the basis for an integrated, rather than reductionist, analysis of complex biological phenomena.

Acetylcholine is a major neurotransmitter in the enteric nervous system and the gut *per se*. Its effects are mediated by nicotinic and muscarinic receptors. The nACh receptors are non-selective ionotropic receptors; whereas μ ACh receptors are part of the metabotropic - G - protein coupled receptors family. Activation of nACh receptors results in the generation of an inward calcium current with the production of fast (fEPSP) or slow (sEPSP) excitatory postsynaptic potentials, respectively. In contrast to the diverse effects of serotonin in the gastrointestinal tract, ACh always has an excitatory effect and enhances myoelectrical activity.

The ubiquitous biogenic amine 5-hydroxytryptamine (5-HT, serotonin) is present in the neurons of the enteric nervous system and to a greater extent in the enterochromaffin (EC) cells. Serotonin acts as a neurotransmitter and a paracrine messenger in the gastrointestinal tract to mediate a wide range of physiological functions. These effects are achieved through activation of 5-HT₁ - 5-HT₇ receptors.

A distinct neural receptor, 5-HT₃, belongs to the family of ligand-gated ion channels. Serotonin applied by ionophoresis to a neuron with 5-HT₃ receptors causes a short latency (< 100 ms) and a duration (< 2 s) depolarization by invoking a fast inward excitation current. The latter is due to an increase in permeability of calcium ($g_{Ca^{2+}}$), potassium (g_{K^+}) and sodium (g_{Na^+}) channels.

The 5-HT₄ receptors belong to a G α_s - protein - coupled family. They are positively linked to adenylyl cyclase in the second messenger signal transduction mechanism. The biological effect of their pharmacological activation is correlated with an increase in permeability of calcium-activated potassium (g_{Ca^{2+},K^+}) and sodium (g_{Na^+}) channels and a decrease in permeability of potassium (g_{K^+}) channels.

The 5-HT₃ and 5-HT₄ receptors and ACh-receptors are expressed on the somas and presynaptic nerve terminals of the primary sensory (AH - type), motor (S - type) and possibly interneurons (S and AH-types) of the myenteric nervous plexus (Johnson and Heinemann, 1995). Electrochemical coupling at the neuro-neuronal synapses entails the initiation of the cascade reactions of the second messenger system and activation of ligand-operated channels. The differences in dynamics of the implicated biochemical pathways, complex allosteric neurotransmitter-receptor interactions, species and regional organ dependence, result in highly variable biological responses.

The aim of this research is to gain insight into phenomenological mechanisms of co-transmission by ACh and 5-HT and their effects on the signal transduction process in the afferent pathway of the enteric neural network of the gut. We focus primarily on the analysis of: i) the concurrent effects of co-activation of muscarinic (μ ACh), nicotinic (nACh) and 5-HT types 3 and 4, receptors, on the dynamics of signal transduction and ii) the role of pharmacological agents with selective, nonselective and mixed 5-HT₃ and 5-HT₄ - receptor agonists/antagonists mechanisms of action on the electrical activity.

The study involves mathematical concepts to model electromechanical processes of the abdominal viscera. The discussion of general physiological facts, which support the scientific concepts, mathematical formulation of the model, derivation of basic equations, construction and validation of the numerical algorithm, software design, have been described by Miftakhov and Christensen (2001), Miftakhov *et al.* (1999 a,b).

Materials and Methods

A model of the bursting AH and S type neurons linked in sequence is based on the Hodgkin-Huxley formalism and includes activity of the voltage-dependent Na^+ , K^+ , Ca^{2+} , Ca^{2+} - K^+ and leak Cl^- channels (Tuladhar *et al.*, 1997; Bertrand *et al.*, 2000; Van Hooft *et al.*, 1998). The dynamics of the propagation of electrical signals along the unmyelinated axons satisfies the classical Hodgkin-Huxley model. The synapse and the drug - receptor interaction models are equivalent to the pharmacodynamic model proposed by Miftakhov and Christensen (2001).

As a tool in our theoretical investigations we exploit a novel computational platform, Gut Discovery[®] (www.aincompany.com). A one-dimensional model that is composed of the interconnected primary sensory (AH) and motor (S) neurons and the smooth muscle syncytium is designed to analyze the electromechanical events in the gut. The model reproduces the following sequence of electrophysiological processes in the afferent pathway and smooth muscle syncytium: i) deformation of the free nerve endings of the mechanoreceptors by mechanical stimuli of a known intensity and duration; ii) generation of action potentials (AP) and their propagation along the unmyelinated fiber towards the soma of the AH - neuron; iii) the action potential generation at the soma of the AH - neuron and the propagation of the wave of depolarization along the nerve axon towards the soma of the S - neuron; iv) electrochemical coupling at the axo-dendritic synapse on the soma of the secondary neuron and generation of the fEPSP; v) discharge of the soma of the S - neuron and the propagation of the wave of depolarization along the nerve axon towards the neuro-muscular synapse; vi) electrochemical coupling at the synapse and fEPSP generation; vii) activation of L-type Ca^{2+} channels of the smooth muscle membrane; viii) active force generation and deformation of smooth muscle. The standard parameter settings depend on the identity of the cells types and can be obtained from Miftakhov *et al.* (1999 a,b).

With the model we study the effects of 5-HT and ACh alone and as a result of their interactions through the process of co-transmission and co-activation of different receptor types. The effect of drugs is achieved by varying conductances for affected channels or parameters involved in the second messenger system pathway. Electrophysiological and mechanical parameters of the model are adjusted during each experiment to reproduce the species variability.

Results

Electrical Activity of the Mechanoreceptors

Physiological Conditions

Under normal physiological conditions, deformation (ϵ) of the wall of the gut with multiple stretches, $\epsilon = 0.17$, $n = 10$, of duration, $t_d = 0.9$ s, followed at intervals, $\Delta t = 1.5$ s, initiates the voltage-dependent inward sodium, I_{Na} , outward potassium, I_{K} , currents and leak chloride, I_{Cl} , current at the free nerve endings of the mechanoreceptors. The sodium influx has an average rate 1 nA s^{-1} and reaches the maximum amplitude 1.14 nA (Fig. 1a). It has almost instant recovery phase, during which the strength of the current reduces to 0.23 nA , followed by a prolonged period, 0.8 s, of slow decline to the resting value: $\min I_{\text{Na}} = 0.042 \text{ nA}$. An outward K^+ current shows longer duration. The dynamics of activation, I_{K} , is linear with the exponential inactivation (Fig. 1b). The current rises to its maximum, $I_{\text{K}} = 0.04 \text{ nA}$, at a constant rate 0.04 nA s^{-1} and declines rapidly to a level 0.075 nA . The balanced activity of the ion currents, results in the production of the dendritic action potentials (APs) (φ_0^d) at the free nerve endings of the mechanoreceptors. They have maximum amplitude 15.7 mV and duration $\sim 2.3 \text{ ms}$.

Effect of 5-HT₃ Receptors

The mechanical deformation causes the release of endogenous 5-HT from the EC cells. Free serotonin binds to the 5-HT₃ receptors on the free nerve endings of the mechanoreceptors with a




Fig. 1a




Fig. 1b

Fig. 1 a,b: Dynamics of the I_K and I_{Na} ion currents at the free nerve endings - mechanoreceptors, under normal conditions (A); after activation of the 5-HT₃ receptors (B) and; following application of the selective 5-HT₃ receptor antagonists (C), respectively

subsequent increase in permeability of ligand-gated ion channels. The I_{Na} current demonstrates nonlinear exponential behavior with an initial rapid rise, at a rate 2.04 nA s^{-1} , followed by a slow dynamic, 0.2 nA s^{-1} (Fig. 1b). It reaches $\max I_{Na} = 1.77 \text{ nA}$. The recovery phase of the channel remains unchanged from that seen above. The dynamics of potassium channel activity have an exponential phase of activation with a slow start at the beginning, 0.05 nA s^{-1} and a quick climb, 0.3 nA s^{-1} , to the maximum value, 0.188 nA (Fig. 1a). In the presence of endogenous serotonin the process of inactivation of the I_K goes on faster, 0.6 nA s^{-1} . The above change in the ionic currents reflects an increase in the amplitude of APs to 26 mV .

Presence of endogenous 5-HT reduces the threshold for mechanical activation of the mechanoreceptors by 17.6%. Multiple stretch impulses of $\epsilon = 0.14$ applied to the wall of the gut are sufficient to initiate the cascade of electrical events as described above.

Effect of Selective 5-HT₃ Receptor Antagonists

Treatment of the mucosa of the gut with selective 5-HT₃ receptor antagonists, *Ondansetron* (GlaxoSmithKline) or *Granisetron* (Roche), abolishes the potassium current ($I_K \simeq 0 \text{ nA}$). There is a significant reduction in the sodium influx. The I_{Na} current shows an activation rate of 0.16 nA s^{-1} and reaches the maximum amplitude of 0.38 nA . As a result, the dendritic action potentials of the amplitude 3 mV are generated. They are not strong enough to set off electrical signals at the free nerve endings and to sustain their propagation of along the unmyelinated fibers. However, an increase in the intensity of the mechanical stimulus ϵ by 35% ($\epsilon = 0.23$) results in an unexpected response from the mechanoreceptors. There is an increase in both the I_{Na} and I_K currents: $\max I_{Na} = 1.04 \text{ nA}$ and $\max I_K = 0.2 \text{ nA}$, are recorded (Fig. 1a,b). The mechanoreceptors produce action potentials: $\varphi_0^d = 22.9 \text{ mV}$.

Electrical Activity of the Primary (AH) Neuron Stimulation of Mechanoreceptors

The dendritic action potentials elicited at the free nerve endings propagate along the unmyelinated axon and reach the soma of the neuron. There they activate the voltage-dependent Na^+ , K^+ , Ca^{2+} , Ca^{2+} - K^+ and leak Cl^- channels. The dynamics of the inward P_{Na} and outward P_K currents demonstrate high frequency oscillatory behavior with fast alternations of activation and inactivation (Fig. 2a). The maximal currents, $P_{Na} = 81.5 \text{ nA}$ and $P_K = 29.8 \text{ nA}$ are registered immediately after excitation of the membrane. In 0.2 s both currents reduce in strength: $P_{Na} = 40.8 \text{ nA}$ and $P_K = 15 \text{ nA}$ are recorded. The sodium and potassium channels remain active for 0.8 s .

The dynamics of activation of Ca^{2+} channels has slow initial phase, 1.9 nA s^{-1} (Fig. 2b). It is followed by a fast phase, 14.8 nA s^{-1} , when the maximum value, 11.65 nA , is reached. There are high frequency oscillations, $0.8\text{-}3.8 \text{ (nA)}$, in the current behavior that occur when the channel starts approaching its peak of activation. The recovery of the P_{Ca} shows the linear dynamics at a rate $\sim 23 \text{ nA s}^{-1}$. A similar pattern is seen with the outward P_{Ca-K} . It quickly attains the maximum level and sustains it for 0.6 s . The $\max P_{Ca-K} = 10 \text{ nA}$. There are high fluctuations of the current, which are only present during activation of the channel.

In absence of the free endogenous serotonin the AH neuron responds only to odd excitatory signals arriving from the mechanoreceptors with the bursts of action potentials, φ_1 , of the maximum amplitude 90 mV (Fig. 2c). The pattern of discharges resemble a bursting chaos type of electrical activity at a variable frequency $\nu=8\text{-}10 \text{ (Hz)}$.

Effect of 5-HT₃ Receptors

A selective stimulation of the 5-HT₃ receptors by 5-HT has a profound effect on activity of all ion channels. The duration of the active state of the Na^+ and K^+ channels increases to 1.54 s . There is a significant rise in intensity of the currents: $\max P_{Na} = 154.3 \text{ nA}$ and $\max P_K = 52.3 \text{ nA}$. The brisk

increase in the currents is followed by a lengthy period of recovery when fluctuations of the currents of average amplitudes, 61.7 nA and 35 nA, respectively, are observed.

The amplitude of calcium activated potassium current is: $\max P_{Ca-K} = 14$ nA. It demonstrates a slight increased activation rate, 25 nA s^{-1} . The inactivation phase of the P_{Ca-K} remains intact from the one considered above (Fig. 4a). There are no changes in the P_{Ca} . However, the channel sustains its active state for a longer period.

As a result of the above alterations the primary sensory neuron generates $\max \varphi_1 = 111.3$ mV. It is important to note that the neuron requires a lower level of deformation, $\min \epsilon = 0.14$, to respond with the production of action potentials to each impulse of excitation arriving from the free nerve endings. Its electrical pattern of discharges resembles beating mode with a frequency 7 Hz.

Effect of Selective 5-HT₃ Receptor Antagonists

Ondansetron and *Granisetron* applied separately completely abolish any electrical activity of the neuron. The neuron remains silent after external periodic stimulations of intensities, 0.14 - 0.17, at the free nerve endings of the mechanoreceptors. However, larger deformation, $\epsilon = 0.24$, causes an unexpected production of action potentials of short duration, 0.3 s and $\max \varphi_1 = 91$ mV. A gradual increase in the concentration of the selective 5-HT₃ antagonists has a dose-dependent effect on the reduction of excitability of the neuron.

Effect of 5-HT₄ Receptors

Excitation of the 5-HT₄ receptors on the soma of the primary sensory neuron by the free serotonin results in the production of high amplitude, $P_{Na} = 166.6$ nA and $P_K = 46.1$ nA and short duration, 0.2 s, sodium and potassium currents. The Na⁺ channel returns quickly to its unexcited state, while K⁺ channels require ~ 1 s to recover. The calcium current shows fast activation, at a rate of 36 nA s^{-1} and reaches the maximal amplitude, $\max P_{Ca} = 21.7$ nA. The dynamics of its recovery has two distinct phases: the initial phase with a rate 5.6 nA s^{-1} and the faster phase - 8.64 nA s^{-1} . The P_{Ca-K} shows a relatively smooth rise to the maximum value, 12.6 nA. The P_{Ca-K} remains active for ~ 2.55 s (Fig. 2a).

There is an increase in depolarization of the soma with a shift of the resting membrane potential to 50 mV. The neuron fires a few high amplitude APs, $\max \varphi_1 = 96.2$ mV, followed by a slow process of decay of depolarization. There is also a reduction in duration, 0.6 s, of the period of after-hyperpolarization.

Effect of Co-expression by 5-HT₃ and 5-HT₄ Receptors

A concomitant stimulation of the co-localized 5-HT₃ and 5-HT₄ receptors on the soma of the primary neuron results in increase in its excitability. At the beginning of the process, when activity of the ligand-gated 5-HT₃ receptors dominates, the currents of amplitude $P_{Na} = 25$ nA and $P_K = 36$ nA are recorded. As a result of a full affect of the 5-HT₄ receptors the dynamics of the sodium and potassium channels change to a high frequency oscillatory mode. Ion current of intensity $P_{Na} = 45$ nA and $P_K = 12$ nA are generated. The Ca²⁺ channel reiterates the type of activity observed under a separate stimulation of the 5-HT₃ receptors. With concurrent excitation of the 5-HT₄ receptors the pattern of the P_{Ca} changes to a beating mode with constant amplitude of fluctuations, 4 nA (Fig. 2b).

The neuron produces spikes of high amplitude, $\max \varphi_1 = 81$ mV, followed by a long quiescent period of sustained depolarization: $-13 < \varphi_1 < 25$ mV. This response is characteristic of a separate activity of the fast ligand-gated 5-HT₃ receptors. A transitory period of irregular chaotic type discharges of APs corresponds to the gradual activation of the 5-HT₄ receptors. Amplitude of φ_1 increases from 5 to 85 (mV) and the frequency of firing varies between $18 < \nu < 37$ (Hz). As the 5-HT₄ receptors become fully active the pattern of electrical activity of the soma begins to resemble a beating mode. Action potentials of average amplitude $\varphi_1 = 50$ mV and a frequency $\nu = 28$ Hz are generated.

The selective 5-HT₃ receptor antagonist, *Ondansetron*, abolishes the production of spikes on the soma. Continuous activation of the 5-HT₄ receptors results in a translation of the electrical pattern of the neuron from the beating to a self-excitatory mode. Regular rhythmic undulations of the membrane potential of amplitude 30 mV with single spikes, max $\varphi_1 = 40$ mV, on crests of the waves are produced. This effect persists while the 5-HT₄ receptors remain active.

Effect of Cisapride

A non-selective strong 5-HT₃ and weak 5-HT₄- receptor agonists, *Cisapride*, increases the influx of calcium ions into the cell. The maximal amplitude of the P_{Ca} equals 17.2 nA. There are changes in the patterns of activity of the P_{Na} , P_{Ca-K} and P_K currents with a period of irregular high frequency oscillatory activity and an interim quiescent period. Thus the maximal values of the P_{Na} and P_K of 228.5 nA and 57.6 nA, respectively, are achieved immediately after the application of *Cisapride*. The second peak of activity of the duration 0.5 s shows max $P_{Na} = 38$ nA and $P_K = 18.8$ nA. The P_{Ca-K} current demonstrates a steady rise in amplitude with max $P_{Ca-K} = 13$ nA at the beginning of the process followed by the second peak of $P_{Ca-K} = 19.2$ nA (Fig. 2a).

Application of *Cisapride* depolarizes the soma of the primary neuron and elevates the resting membrane potential: $\varphi_{1(rest)} = -62.5$ mV. The neuron generates spikes of high amplitude, max $\varphi_1 = 111$ mV and frequency, $7 \div 18$ (Hz). There is a period of 0.65 s when no action potentials are produced. It is followed by a short period of bursting activity with max $\varphi_1 = 62$ mV and duration 0.5 s.

Electrical Activity of the Motor (S) Neuron

Effect of nACh Receptors

Depolarization of the presynaptic membrane at the neuro-neuronal synapse activates release of vesicular acetylcholine and its diffusion into the synaptic cleft. The main part of ACh reaches the postsynaptic membrane and reacts with the receptors on the soma of the motor neuron. The generation of fEPSP starts with the beginning of (ACh-R)-complex development. It increases as a step function and achieves its maximum 87.1 mV in 0.25 ms. The level of fEPSP is sufficient to excite the neuron. It discharges a single train of pulses of amplitude 102.1 mV at a frequency 8.4 Hz and duration 1.9 s. The observed pattern of electrical activity resembles a regular bursting mode.

Effect of the 5-HT₃ and 5-HT₄ Receptors

A selective stimulation of 5-HT₃ receptors by the free endogenous 5-HT in absence of ACh in the system does not have excitatory effect on the soma of the motor neuron.

In contrast, activation of the 5-HT₄ receptors changes the pattern of neuronal electrical activity into a long lasting excitatory mode. Action potentials, φ_2 , of constant amplitude 114 mV and a frequency 7 Hz are produced. The ion currents: I_{Na}^m of magnitude $22,5 \times 10^2$ nA, $I_K^m \simeq 148$ nA, $I_{Ca-K}^m \simeq 16.2$ nA and $I_{Ca}^m \simeq 6$ nA are recorded (Fig. 3).

Effect of Co-activation of the 5-HT₃ and nACh Receptors

Simultaneous excitation of the 5-HT₃ and nACh receptors on the soma of the motor neuron results in its hyperpolarization, $\varphi_{2(rest)} = 66.8$ mV. The neuron generates action potentials of the maximal amplitude 134 mV at a frequency 7.8 Hz. The observed electrical activity is due to an increase in the dynamics of ion currents: $I_{Na}^m = 18,5 \times 10^2$ nA, $I_K^m = 18$ nA and $I_{Ca-K}^m = 14.8$ nA. They demonstrate a regular high frequency mode of activation - inactivation. The calcium current shows an intensive oscillatory phase of activation followed by a slow recovery period. Max $I_{Ca}^m = 11.9$ nA is recorded (Fig. 4).





Fig. 2a





Fig. 2a (Continued)




Fig. 2b





Fig. 2b (Continued)





Fig. 2c








Fig. 2c

Fig. 2 a,b,c: Dynamics of the F_{Ca-K} , F_{Ca} ion currents and action potentials, f , on the soma of the primary (AH) sensory neuron after selective stimulation of: 5-HT₃ receptors (A); 5-HT₄ receptors (B); concomitant stimulation of the 5-HT₃ and 5-HT₄ receptors (C) and after treatment with *Cisapride* (D)








Fig. 3: Dynamics of the I_{Na}^m (A), I_K^m (B), I_{Ca}^m (C) and I_{Ca}^m (D) ion currents and action potential, (E), f on the soma of the motor (S) neuron after selective stimulation of the 5-HT₄ receptors




Fig. 4




Fig. 4 (Continued)




Fig. 4: Dynamics of the I_{Na}^m (A), I_K^m (B), I_{Ca}^m (C) and I_{Ca}^m (D) ion currents and action potential, (E), f on the soma of the motor (S) neuron after co-joint stimulation by 5-HT and ACh of 5-HT₃ and nACh receptors




Fig. 5: Changes in the dynamics of the I_{Ca}^m (A) ion current and action potential, ϕ (B), on the soma of the S-neuron after application of *Cisapride* and co-stimulation of the nACh receptors




Fig. 6: Changes in slow wave, f (A) and active force, T_a (B), dynamics in the longitudinal smooth muscle after co-joint activation of the 5-HT_4 and μACh receptors

Effect of co-activation of the 5-HT₄ and nACh Receptors

Concurrent stimulation of co-localized 5-HT₄ and nACh receptors initiates strong large magnitude sodium ion current, max $I_{Na}^m = 27 \times 10^2$ nA. There is a concomitant slight decrease in amplitude of the I_{K}^m , I_{Ca-K}^m and I_{Ca}^m currents, compare to a separate excitation of 5-HT₄ and/or nACh receptors. These changes lead to the generation of the spikes of amplitude, $\varphi_2 = 116.5$ mV.

Effect of Selective 5-HT₃ and 5-HT₄ Receptor Agonists and the Co-expression of nACh Receptors

Cisapride and *renzapride* added into the system have a pronounced effect on the dynamics of the calcium current. Thus max $I_{Ca}^m = 7.04$ nA is observed. The Ca²⁺ channel shows oscillatory behavior with fluctuations of the current of amplitude 4.2 nA and constant frequency: $\nu = 6.4$ Hz. Ion currents of intensity: $I_{K}^m = 210$ nA, $I_{Ca-K}^m = 16.7$ nA and $I_{Na}^m = 28.5 \times 10^2$ nA are recorded. The soma of the neuron is hyperpolarized to -70 mV. The neuron remains in a highly excitable state for the duration of 3.44 s and generates APs max $\varphi_2 = 130$ mV and a frequency of $\nu = 6.4$ Hz.

No significant changes are observed in the pattern of behavior of I_{K}^m , I_{Ca-K}^m and I_{Na}^m ion currents after simultaneous application of *Cisapride* and the release of ACh into the system. However, the dynamics of the I_{Ca}^m demonstrates two distinct peaks of activation (Fig. 5). Immediately after activation of the 5-HT₃, 5-HT₄ and nACh receptors max $I_{Ca}^m = 6.8$ nA is observed. The current slightly decreases in intensity to 6.45 nA during the following 1.2 s. The second maximum of the influx of calcium ions with max $I_{Ca}^m = 9.5$ nA is recorded 1 s later. The current quickly subsides to its inactive state.

Interestingly the duration of electrical excitability of the neuron reduces to 2.8 s. There is a decrease in the maximal amplitude of APs, max $\varphi_2 = 120$ mV and an increase in frequency of spikes: $\nu = 7.8$ Hz. The level of hyperpolarization of the soma remains unchanged.

Effects of Selective 5-HT₃ - Receptor Antagonist and Co-expression of nACh Receptors

Ondansetron and *Granisetron* block the serotonergic pathway between the primary sensory and motor neurons. The normal electrical signal transduction is maintained through the co-existing cholinergic mechanisms. The latter sustain the neuro-neuronal synaptic connectivity. They are responsible for the generation of short duration (< 100 ms) and 85-90 mV in amplitude fEPSP on the soma of the adjacent motor neuron.

Electromechanical Activity of Smooth Muscle

Effect of μ ACh receptors. fEPSP at the neuro-muscular synapse activate the L-type Ca²⁺ channels located on the smooth muscle membrane. Alterations of the permeability of the channels cause cyclic transitory changes in the myoelectrical pattern. The slow wave mode transforms to bursting chaos with the generation of fast APs at a frequency of $\nu = 17$ Hz. Their amplitude is 23 mV at the beginning and decreases towards the end of the burst, $\varphi_m = 8$ mV. The bursting chaos transforms to regular bursting with generation of spikes on the crests of slow waves. The burst amplitude, burst duration and number of spikes per burst all increase. The action potentials have maximum amplitude of 72 mV and oscillate at a maximum frequency of 19 Hz. The regular bursting converts back to bursting chaos and slow wave mode.

There is a gradual increase in the concentration of intracellular calcium ions, mainly due to the influx of ions through the activated slow Ca²⁺ channels. Maximum internal calcium concentration ($[Ca^{2+}]_i = 0.46$ μ M) is achieved. As a result of activation of the contractile protein system, rhythmic contractions of the longitudinal smooth muscle, max $T^a = 14.4$ mN cm⁻¹, are produced. The duration of electromechanical activity is 2.6 s.

Effect of 5-HT₄ Receptors

Selective 5-HT₄ receptor agonists, TS-591, prucalopride and ML10302, applied onto the longitudinal smooth muscle of the gut do not have any effects on its electromechanical activity.

Effect of co-activation of the 5-HT₄ and μ ACh Receptors

Excitation of the 5-HT₄ receptors with TS-591, prucalopride and ML10302 in conjunction with μ ACh receptors evokes an increase in intensity of the membrane ionic currents and electromechanical activity of the smooth muscle syncytium. The F_{Ca} demonstrates high frequency alternations, 18.5 Hz, of short duration. Average amplitude of the L-type Ca²⁺ current is 7 nA and max $F_{Ca} = 10.6$ nA. The dynamics of the transitory calcium current, F_{Ca} , remains unchanged. It oscillates at a frequency 21 Hz and has average amplitude 0.30 nA. The outward voltage-dependent K⁺ and Ca²⁺ - K⁺ currents are reduced in strength. Maximum $F_{Ca-K} = 1.37$ nA and $F_K = 4.6$ nA are recorded. The above changes in membrane ionic currents cause an increase in the frequency of the production of APs, $\nu = 20$ Hz. The amplitude of spikes is slightly decreased, $\phi_m = 68$ mV, compare to that observed when only μ ACh receptors are active.

The concentration of free cytosolic calcium is 0.46 μ M. As a result, the longitudinal smooth muscle responds with the production of phasic contractions. The maximum active force (T^a) generated equals 15.6 g. The contractions are of duration 3.45 s, concurrent in phase and time with the dynamics of the intracellular Ca²⁺ oscillations and have constant amplitude, 6 g. The calculated velocity of twitch is 5.8 g cm⁻¹ (Fig. 6A and B).

Effect of Selective 5-HT₄ Receptor Antagonists and the Co-activation of the 5-HT₃ and μ ACh Receptors

Addition of the selective 5-HT₄ antagonist, GR113808A, strongly inhibits contractions evoked by the free endogenous serotonin. However, electromechanical activity can be sustained through the function of the neuronal 5-HT₃ and nACh receptors and μ ACh receptors located to the smooth muscle membrane. Thus, application of *Cisapride* leads to twitch contractions of smooth muscle of amplitude $T^a = 13.8$ mN cm⁻¹.

Discussion

With our experiments we were able to reproduce numerically the pharmacological patterns of behavior and to study the interplay between ACh and 5-HT. We analyzed systematically their exact effects on electrical activity of the primary sensory (AH) neuron and the motor (S) neuron in the afferent neural pathway of the enteric nervous plexus.

Comparison of our numerical results to experimental data shows satisfactory qualitative and quantitative agreement. Thus, using the isolated guinea pig ileum, Tuladhar *et al.* (1997; 2000) convincingly demonstrated that activation of the 5-HT₃ receptors by mucosally applied free 5-HT reduces the mechanical threshold for initiation of the peristaltic reflex by 20-26%. Present results show a 17.6% decrease from the required min ϵ . Further authors showed that treatment of the preparation with selective 5-HT₃ receptor antagonists, *Ondansetron* and *Granisetron*, resulted on an average 44% increase in the level of mechanical stimulus needed, which is in a satisfactory agreement with our findings - 35%. Present results indicate that a possible mechanism is associated with an increase in the conductivity of the Na⁺ and K⁺ channels and changes in the dynamics of activation of the I_{Na} and inactivation of the I_K currents. Interestingly, the inhibitory effect of *Ondansetron* and *Granisetron* with a complete suppression of the potassium current ($I_K \simeq 0$ nA) and a significant reduction in strength of the I_{Na} current can be overcome by a higher level of mechanical deformation. This fact suggests that the 5-HT₃ receptors may be directly linked to the mechanosensitive Na⁺ channels or are located in close proximity to them. However, cautious interpretation of the proposed mechanism is necessary. There

is currently no experimental evidence on the dynamics of the Na⁺ and K⁺ currents at the mechanoreceptor level to support or disprove this view.

Experimental intracellular recordings from the soma of the primary neurons of the guinea-pig ileum enteric nervous plexus demonstrate that after application of 5-HT (Bertrand *et al.*, 2000) the neuron produces long lasting, 2 - 4 s, trains of action potentials of amplitude 75-80 mV and a frequency ranging from 0.1 to 10 (Hz). Numerical analysis of the dynamics of the ion currents helps suggest the mechanisms responsible for the observed response. Thus, major changes are registered in the P_{Na} and P_K currents with high-frequency fluctuations recorded during the recovery phase. It is possible to assume that the increased excitability of the mechanoreceptors and instability of the Na⁺ and K⁺ currents decrease the threshold of activation of the AH neuron.

Selective activation of the 5-HT₃ receptors on the soma of the S neuron of the guinea-pig ileum induces action potentials of 5 mV and of low frequency (Bertrand *et al.*, 2000). These experimental findings are of particular interest because the physiological saline in the experimental preparation contained nicardipine and scopolamine - drugs that affect cholinergic transmission. The numerical simulation of co-transmission by ACh and 5-HT shows that the motor neuron generates high amplitude, 134 mV and frequency, 7.8 Hz, action potentials. This response can be attributed to a heteromeric co-assembly of the 5-HT₃ receptors with the nicotinic ACh ($\alpha 4$ subunit) receptors (Van Hooft *et al.*, 1998; Barajas-Lopez *et al.*, 2001; Holler *et al.*, 1999; Pindon *et al.*, 2002). While these experimental facts concur with the results of our numerical simulations the putative role of the 5-HT₃/nACh ligand-channel co-assembly requires further pharmacological investigation.

We were able to analyze numerically the effects of co-expression of the 5-HT₃, 5-HT₄ and nACh receptors on the soma of the S neuron on the dynamics of the electrical signal transduction. Our results demonstrate the minor role played by 5-HT type 3, in contrast to the 5-HT type 4 receptors. Activation of 5-HT₄ receptors evoked a long-lasting train of fast action potentials. Co-stimulation of the (5-HT₃ - nACh) and (5-HT₄ - nACh) receptors has a considerable excitatory effect on the neuron. It resulted in a significant increase in the amplitude and frequency of firing rate. These findings are in line with the belief of possible co-assembly of serotonergic and cholinergic receptors (Legay *et al.*, 1984; Briejer and Schuurkers, 1996; Foxx-Orenstein *et al.*, 1998).

Conclusion

Comprehensive experimental data on the physiology and neuro-pharmacology of the myenteric nervous plexus, combined with modern computational modeling technology, have allowed us to study in a "virtual environment" the effects of co-transmission by 5-HT and ACh and of receptor polymodality, i.e., 5-HT₃, 5-HT₄, nACh and μ ACh type receptors, on its electrical activity. The numerical investigations helped us reveal intrinsic mechanisms of co-transmission at the cellular and tissue levels that could not have been elucidated using the existing experimental *in vivo* or *in vitro* methods.

References

- Barajas-Lopez, C., R. Karanjia and R. Espinosa-Luna, 2001. 5-Hydroxytryptamine and atropine inhibit nicotinic receptors in submucosal neurons. *Eur. J. Pharmacol.*, 414: 113-123.
- Bertrand, P.P., W.A.A. Kunze, J.B. Furness and J.C. Borstein, 2000. The terminals of myenteric intrinsic primary afferent neurons of the guinea-pig ileum are excited by 5-hydroxytryptamine acting at 5-hydroxytryptamine - 3 receptors. *Neuroscience*, 101: 459-469.
- Briejer, M.R. and J.A.J. Schuurkers, 1996. 5-HT₃ and 5-HT₄ receptors and cholinergic and tachykininergic neurotransmission in the guinea-pig proximal colon. *Eur. J. Pharmacol.*, 308: 173-180.

- Foxx-Orenstein, A.E., J.G. Jin and J.R. Grider, 1998. 5-HT₄ receptor agonists and δ -opioid receptor antagonists act synergistically to stimulate colonic propulsion. *Am. J. Physiol.*, 275: G979-G983.
- Holler, C., M. Freissmuth and C. Nanoff, 1999. G proteins as drug targets. *Cell Mol. Life Sci.*, 55: 257-270.
- Johnson, D.S. and S.F. Heinemann, 1995. Detection of 5-HT_{3R-A}, a 5-HT₃ receptor subunit, in submucosal and myenteric ganglia of rat small intestine using *in situ* hybridization. *Neurosci. Lett.*, 184: 7-70.
- Legay, C., M.J. Saffrey and G. Burnstock, 1984. Coexistence of immunoreactive substance P and serotonin in neurons of the gut. *Brain Res.*, 302: 379-382.
- Miftakhov, R.N., G.R. Abdusheva and J. Christensen, 1999a. Numerical simulation of motility patterns of the small bowel. I. Formulation of a mathematical model. *J. Theor. Biol.*, 197: 89-112.
- Miftakhov, R.N., G.R. Abdusheva and J. Christensen, 1999b. Numerical simulation of motility patterns of the small bowel. II. Comparative pharmacological validation of a mathematical model. *J. Theor. Biol.*, 200: 261-290.
- Miftakhov, R. and J. Christensen, 2001. A Physicochemical Basis of Synaptic Transmission in the Myenteric Nervous Plexus. In: *Biophysical Neural Networks*, Poznanski, R. (Eds.), Mary Ann Liebert Inc., NY, pp: 147-176.
- Pindon, A., G. Van Hecke, P. Van Gompel, A.S. Lesage, J.E. Leusen and M. Jurzak, 2002. Differences in signal transduction of two 5-HT₄ receptor splice variants: Compound specificity and dual coupling with G α s - and G α i/o - proteins. *Mol. Pharmacol.*, 61: 85-96.
- Tuladhar, B.R., M. Kaiser and R.J. Naylor, 1997. Evidence for a 5HT₃ receptor involvement in the facilitation of peristalsis on mucosal application of 5-HT in the guinea pig isolated ileum. *Br. J. Pharmacol.*, 122: 1174-1178.
- Tuladhar, B.R., M.D. Womack and R.J. Naylor, 2000. Pharmacological characterization of the 5-HT receptor mediated contraction in the mouse isolated ileum. *Br. J. Pharmacol.*, 1313: 1716-1722.
- Van Hooff, J.A., A.D. Spier, J.L. Yakel, S.C.R. Lummis and H.P.M. Vijverberg, 1998. Promiscuous coassembly of serotonin 5-HT₃ and nicotinic α 4 receptor subunits into Ca²⁺ - permeable ion channels. *Proc. Natl. Acad. Sci. USA*, 95: 11456-11461.