

Journal of
**Pharmacology and
Toxicology**

ISSN 1816-496X



Academic
Journals Inc.

www.academicjournals.com



Research Article

Antioxidant and *in vitro* Toxicity Assessment of Elegant Explosion Daylily Flower (*Heemerocallis* spp.) Extract in Hepatic Cell Line

¹J. Johnson, ²S. Ogutu and ^{1,3}J. Boateng

¹Laboratory of Food Safety and Toxicology, Alabama A and M University, Normal AI, 35872, United States

²Laboratory of Food Chemistry, Alabama A and M University, Normal AI, 35872, United States

³Departments of Food and Animal Sciences, Laboratory of Nutrition Biochemistry, Alabama A and M University, Normal AI, 35872, United States

Abstract

Background and Objective: Edible flowers such as the daylily (*Heemerocallis* spp.) have been shown to have beneficial effects against chronic diseases. This study determined antioxidant activity and polyphenol contents of two cultivars of daylily flowers Barbara Mitchell (BM) and Elegant Explosion (EE). The *in vitro* bioactivities of EE extract were further examined in HepG2 cell lines. **Materials and Methods:** BM and EE daylily flowers extracts were prepared with aqueous ethanol and used to determine Total Phenolic Content (TPC), Total Flavonoid Content (TFC), Total Monomeric Anthocyanins (TMA), Condensed Tannins (CT) and antioxidant assays. Selected phytochemicals were also determined via HPLC/DAD. *In vitro* toxicity, apoptosis and enzymatic activities of EE extracts were assessed in HepG2 cells. **Results:** Results indicated EE showed significantly ($p \leq 0.05$) higher polyphenol contents and antioxidant activity compared to BM. *In vitro* cytotoxicity indicated that EE extracts possess moderately weak toxicity towards HepG2 cells. Significant ($p \leq 0.05$) increases in antioxidant activities were noted. **Conclusion:** With limited studies on polyphenol content and bioactivity of EE daylily flower, these results provide preliminary evidence and contribute to the promotion of edible flowers as a potential dietary supplement and nutraceutical for use as functional foods.

Key words: Edible flowers, daylily, superoxide dismutase, cytotoxicity, nutraceutical, phenolic compounds, endogenous antioxidants

Citation: Johnson, J., S. Ogutu and J. Boateng, 2021. Antioxidant and *in vitro* toxicity assessment of elegant explosion daylily flower (*Heemerocallis* spp.) extract in hepatic cell line. *J. Pharmacol. Toxicol.*, 16: 22-36.

Corresponding Author: Judith Boateng, Laboratory of Food Safety and Toxicology, Alabama A and M University, Normal AI, 35872, United States
Tel: 256-372-8794 Fax: 256-372-5432

Copyright: © 2021 J. Johnson *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

A primary knowledge of edible flower preparations has been documented from accounts described in ancient literature¹⁻³. For example, in 16th century Central Europe, fried batter-coated black elder (*Sambucus nigra*) flowers were common, as were dandelion flowers boiled with ersatz honey²⁻⁴. Ancient Romans enjoyed roses and lavender in sauces^{4,5}. In Victorian-era England, roses were used as flavouring agents in teas, sweets, pastries, conserves and sauces^{4,5}. Besides the wholesomeness and suitability in most aspects of culinary applications, edible flowers are abundant sources of nutrients such as carbohydrates, protein, fat, minerals and vitamins^{6,7}. Additionally, edible flowers contain varying quantities of several bioactive compounds⁸⁻¹¹. These bioactive compounds with antioxidant properties contribute to the medicinal soundness of edible flowers and thus support increased worldwide demand.

The daylily (*Heemerocallis* spp.) is an edible flower, which dates back thousands of years in sub-tropical Asia. The main diversity of the genus is centred in China, Korea and Japan, where they are used as medicine and food¹². For culinary and consumption purposes, different parts of the flower are used as vegetables and incorporated into soups, stews, or eaten raw. Daylilies (*Heemerocallis* spp.) have shown strong antioxidant activities *in vitro* and *vivo*¹³⁻¹⁵. These beneficial effects are believed to be due to several novel bioactive compounds including naphthalene glycoside, stelladerol, caffeoylquinic acid derivatives, as well as other antioxidants such as gallic acid, rutin and catechin^{16,17}. According to Rodriguez-Enriquez and Grant-Downton¹⁷, *Heemerocallis* spp. could have significant potential as a nutraceutical and the source of new compounds with biomedical activities. Thus, the daylily could be considered as a functional food and used in natural products and supplements to improve health.

Although the consumption of flowers is on the rise, it is concluded that there are still numerous aspects that need to be evaluated. This is especially important if it is to be considered as a functional food or nutraceutical. As highlighted by Granato *et al.*¹⁸, assigning a functional label to a potential food product or nutritional supplement requires completion of a proper assessment to attest its functionality using *in vitro* and *in vivo* methods. Given this, Granato *et al.*¹⁸ indicated that a wide variety of tests, both *in vitro* and *in vivo*, must be performed to determine efficacy and bioactivity as well as to assess the physicochemical and chemical properties of any food before stating it is functional.

This study aims to determine the polyphenolic contents and antioxidative activity of daylily cultivars Barbara Mitchell (BM) and Elegant Explosion (EE) and also, to assess their acute *in vitro* toxicity potential (cytotoxicity, apoptotic and antiproliferative properties) in HepG2 hepatic cell line.

MATERIALS AND METHODS

Study area: The study was carried out in the Department of Food Science at Alabama A and M University, Normal AL, United States from September, 2016-December, 2018.

Materials and reagents

Flowers: Barbara Mitchell (BM) and Elegant Explosion (EE) daylily plantlets were obtained from a local nursery (Bennet Nurseries) that cultivates and sells edible flowers from organic production. (The plantlets were confirmed by Dr. Patti Coggins, who has a long history of working with *Heemerocallis* spp.). The plantlets were cultivated and cared for in the greenhouse at Alabama A and M University, United States until harvesting.

Chemicals: All chemicals and reagents that were used in this study were of pure analytical grade and purchased from Fisher Scientific (Suwannee, GA) and Sigma Aldrich (St. Louis, MO). Kits and other supplies were purchased from Promega, Thermo Fisher Scientific (Madison, WI) and Cayman Chemical Company (Ann Arbor, MI). Catechin, Rutin, Gallic acid and Chlorogenic acid standards were acquired from Indofine Chemical Company (Hillsborough, NJ).

Harvesting and extraction of polyphenols from daylily flowers

Flowers: Barbara Mitchell and Elegant Explosion daylily flowers were harvested at full bloom in the morning for a week. The pulling method was used to combine several batches over the days of harvest. Flowers were placed in individual plastic containers with absorbent papers at the bottom to prevent any lymph leakage due to guttation. The harvested flowers were placed on ice and transported to the laboratory in refrigerator bags and stored at -20°C immediately. The flowers were analyzed within 1 week from the collection day.

Extraction procedure: Approximately 70 g of flowers were weighed and homogenized in 200 mL of 80% ethanol. The homogenate was stored in the dark overnight. The supernatant was removed by centrifugation (3000 rpm at 4°C for 20 min). The extraction procedure was repeated two more

times and all the supernatants/extracts were pooled. The extracts were concentrated by evaporation and the concentrated extract was stored at -80°C until analysis. The extracts were prepared in triplicates.

Determination of total polyphenol contents in daylily flowers

Total Phenolics Contents (TPC) and Total Flavonoids Content (TFC): The TPC of daylily extracts was determined using the Folin-Ciocalteu method modified for the microplate Gajula *et al.*¹⁹ The TPC was expressed as mg Gallic acid equivalent per gram sample (mg GAE g^{-1} sample). The TFC was determined via colorimetric assay²⁰ with modifications¹⁹ and was expressed as mg catechin equivalent per gram sample (mg CE g^{-1} sample).

Condensed tannins and Total Monomeric Anthocyanins (TMA) contents:

Condensed tannins were determined by the Vanillin-HCl/vanillin H_2SO_4 assay method¹⁸ and expressed in mg catechin equivalent g^{-1} sample (mg CE g^{-1} sample). The TMA of daylily flowers was determined by the pH differential method with modification^{21,22}. Daylily extracts were diluted with two buffers, potassium chloride buffer (0.025 M, pH 1.0) and sodium acetate buffer (0.4 M, pH 4.5). Diluted samples are read against a blank at 520 and 700 nm. The TMA was expressed as mg cyanidin-3-glucoside equivalent g^{-1} sample (mg C3GE g^{-1} dwb).

Determination of antioxidant activities in daylily flowers

Trolox Equivalent Antioxidant Capacity (TEAC) assay: TEAC was determined following previous methods²³. The TEAC was expressed as a Mmol Trolox g^{-1} sample.

Ferric Reducing Antioxidant Power (FRAP) assay:

The FRAP of daylily extracts was measured using methods described by Benzie and Strain²⁴. The FRAP activity was expressed as an mM Fe^{2+} g^{-1} sample.

DPPH radical scavenging capacity assay:

The DPPH test was performed following the method of Brand-Williams *et al.*²⁵, with modifications²⁶. The IC_{50} was calculated and the decrease in absorbance at 515 nm was monitored at T_0 and T_{30} , T_{60} and T_{90} . The DPPH scavenging ability was calculated as a percentage using the following Eq.²⁶:

$$\text{DPPH scavenging ability (\%)} = \frac{A_{515(t=0)} - A_{515(t=90\text{ min})}}{A_{515(t=0)}} \times 100 \quad (1)$$

Metal chelating activity assay: The Metal chelating activity was measured using a modified method by Wong *et al.*²⁷ Various concentrations of extracts were prepared and 180 μL of methanol (100%) and 5 μL FeCl_2 (2 mM) were added and incubated for 3 min. About 10 μL of ferrozine (5 mM) was added to initiate the reaction and the mixture was incubated for 10 min at room temperature after shaking. Absorbance was read at 562 nm. The percentage of Fe^{2+} -ferrozine complex formation was calculated as Fe^{2+} (%) chelating ability²⁷.

Determination of selected polyphenols in daylily flowers via

HPLC: High-performance liquid chromatography (HPLC) analysis of polyphenols in daylily flowers was carried out on a Varian Modular Analytical HPLC System. Polyphenol's identification was determined using methods of Schutz *et al.*²⁸ with modifications. Samples and standards were prepared using 80% ethanol. Mobile phase A was 0.5% (v/v) acetic acid in the water and mobile phase B was 0.5% (v/v) acetic acid in acetonitrile. Separations of polyphenols were conducted at a flow rate of 0.8 mL min^{-1} at 25°C . Catechin and gallic acid were eluted at 284 nm, chlorogenic acid at 320 nm and rutin at 350 nm. Spectral data was collected for each standard and sample. Polyphenols (i.e., rutin, catechin, chlorogenic acid and gallic acid) were selected based on the previous studies²⁹⁻³².

Determination of *in vitro* cytotoxicity of Elegant Explosion (EE) extract

Cell line and culture conditions of Hep2G cells:

Human hepatocellular carcinoma cell line (HepG2), (ATCC, Manassas, VA) was cultured following previous methods and conditions³³. HepG2 was seeded at 10^4 cells cm^{-2} density and allowed to adhere for 48 hrs and cells were used for cytotoxicity determinations on the third day after seeding.

Determinations of cell viability of EE extract in HepG2 cells:

Cell viability was determined by measuring the metabolic activity of living cells using the PrestoBlue® (Life Technologies Corporation, Grand Island, NY) assay reagent. After the seeding period specified above, HepG2 cells were incubated with different concentrations (0.2, 0.4, 0.8, 1.0, 2.0, 4.0 and 8.0 mg mL^{-1}) of EE extract for 24 and 48 hrs. After the incubation period, the culture medium (supernatant) was removed and cell viability was determined.

Morphological evaluation of EE extract on HepG2 cells:

HepG2 cells (1×10^4) were grown on coverslips and treated with or without daylily extracts for 48 hrs. The medium was discarded and cells were washed once with PBS. Cells were

fixed with MeOH: acetic acid (3:1) and stained with NucBlue® Live reagent (Hoechst 33342) and NucGreen® Dead reagent (Life Technologies). After 15 min of incubation, the dye was removed and the cells were rinsed with PBS and imaged with the EVOS® FL Auto system (Thermo Fisher Scientific, Waltham, MA).

Determination of antioxidant/detoxification enzymes

Extraction of protein from cells: After treatment with daylily extract, cell lysate was collected from HepG2 cells following a protocol from our lab^{33,34}. Protein in the cell lysate was determined using a BCA protein assay kit from Pierce (Rockford, IL). The cell lysate was used for enzyme analysis.

Determination of endogenous antioxidant enzymatic activity:

Glutathione (GSH) was determined according to Griffith³⁵ and expressed as $\mu\text{mol mg}^{-1}$ protein. Superoxide Dismutase (SOD) was determined by the method outlined by Fernández-Urrusuno *et al.*³⁶. Xanthine oxidase was utilized to generate a superoxide flux and samples were measured at 440 nm after the addition of xanthine oxidase. Data were expressed as units of SOD activity per mg protein (U mg^{-1} protein). Glutathione-S-transferase (GST) activity was based on the method of Habig *et al.*³⁷ and 1-chloro-2, 4-dinitrobenzene was used as the substrate. GST was expressed as $\mu\text{mol mg}^{-1}$ protein. All samples were analyzed in triplicates.

Statistical analysis: All experiments were repeated three independent times and results were conducted in triplicates. ANOVA was used to determine any significant differences among daylily cultivars. Data for the antioxidant assays are presented as the mean \pm standard deviation of the mean (SD) and those for the cytotoxicity assays are presented as the mean percentages relative to the control \pm SD. Significant ($p \leq 0.05$ or 0.1), means were separated using Tukey's Studentized Range Test. Statistical analysis was conducted using SAS, 9.4 tools.

RESULTS AND DISCUSSION

Total polyphenol contents in daylily flowers: The results of TPC, TFC, TMA and CTC are shown in Table 1. The TPC was significantly ($p \leq 0.05$) higher in EE ($2.13 \text{ mg GAE g}^{-1}$ dwb) compared to BM ($1.11 \text{ mg GAE g}^{-1}$ dwb). TFC showed a parallel trend between the BM and EE flowers. The TFC was approximately two times higher in EE ($1.47 \text{ mg CE g}^{-1}$ dwb) than in BM ($0.64 \text{ mg CE g}^{-1}$ dwb). While the results for TFC in

EE and BM are within the range reported elsewhere for edible flowers³⁸⁻⁴¹, there is high variability in TPC. The TPC reported in this study was lower compared to other studies. This variation may be due to the extraction methodology used or the types of solvent employed. Additionally, several factors such as genotype, environment, geographical location and altitude, UV exposure, season, soil, maturity stage at harvest, post-harvest conditions, etc. could all impact TPC variation. It should be noted that the daylily cultivars utilized in this study were cultivated in the AAMU greenhouse under controlled culture conditions. Flavonoids are well-known antioxidants with importance in human health. It is reported that due to their strong antioxidant capacity, most flavonoids outperform natural and well-known antioxidants including ascorbate (vitamin C) and α -tocopherol (vitamin E) because of their strong ability to donate electrons or hydrogen atoms to neutralize free radicals⁴². This indicates their potential as important therapeutic agents and also for improving the quality of foods⁴³. The contents of anthocyanins in edible flowers have been well documented^{7,44,45}. Among the most prominent anthocyanins, cyanidin-3-glucoside (C3G) is the major anthocyanin found in most flowering plants. The data of Table 1 shows the TMA content in EE was significantly ($p \leq 0.05$) higher ($1.73 \text{ mg C3G g}^{-1}$ dwb) compared to BM ($1.20 \text{ mg C3G g}^{-1}$ dwb). These results are comparable to other studies⁴⁶⁻⁵⁰. Anthocyanins, besides their importance as natural colourants or pigments in foods, have various beneficial health benefits^{46,49,51-56}. A similar trend was observed whereby CTC was significantly ($p \leq 0.05$) higher in EE ($0.024 \text{ mg CE g}^{-1}$ dwb) than in BM ($0.004 \text{ mg CE g}^{-1}$ dwb) (Table 1). Condensed tannins (proanthocyanidins) have found various applications in foods. They are used as additives, they prevent oxidation in foods and enhance the shelf life of beverages including teas, fruit juices and wine⁵⁷. For health purposes, condensed tannins or proanthocyanidins are noted for their antibacterial activity against urinary tract infections, for beneficial regulation of gut microbiota^{58,59} as well as anti-inflammatory enhancing properties^{57,60,61}.

Antioxidant activities of daylily flowers: The result of Table 2 show the FRAP, TEAC, DPPH and Fe^{2+} chelating antioxidant activities of EE and BM flowers. When it comes to determining the antioxidant activity of plant extracts there's not a standardized method for evaluation since the extracts may exhibit different antioxidant mechanisms. As such, it is important to perform more than one type of antioxidant activity measurement to take into account the various




Fig. 1: DPPH (%) of daylily cultivars at T₉₀ min

Values are expressed as means ± SD and were conducted in triplicates in three independent experiments. Statistical significance was determined by one-way ANOVA followed by Tukey's range test. EE: Elegant explosion, BM: Barbara mitchell

Table 1: Total polyphenol contents in daylily cultivars

Samples	TPC (mg GAE g ⁻¹ dwb)	TFC (mg CE g ⁻¹ dwb)	TMA (mg C3GE g ⁻¹ dwb)	CTC (mg CE g ⁻¹ dwb)
BM	1.11 ± 0.06 ^b	0.64 ± 0.01 ^b	1.20 ± 0.15 ^b	0.004 ^b
EE	2.13 ± 0.01 ^a	1.47 ± 0.02 ^a	1.73 ± 0.08 ^a	0.024 ^a

Values (n = 3) are means ± SD, means in the column (^{ab}) letter differ (p ≤ 0.05) using Tukey's Studentized range test. BM: Barbara mitchell, EE: Elegant explosion, GAE: Gallic acid equivalent, DWB: Dry weight basis (dry flowers), TFC: Total flavonoid, TPC: Total phenolic, TMA: Total monomeric anthocyanin content, CTC: Condensed tannins content, CE: Catechin equivalent, GAE: Gallic acid equivalent, C3G: Cyanidin-3-glucoside equivalent

Table 2: Antioxidant activities of daylily cultivars

Samples	FRAP	TEAC	DPPH (IC ₅₀ mg mL ⁻¹)	Fe ²⁺ chelating activity (IC ₅₀ mg mL ⁻¹)
BM	0.97 ± 0.04 ^b	4.82 ± 0.02 ^b	24	43.08
EE	2.48 ± 0.15 ^a	16.15 ± 0.24 ^a	14.31	37.05

Values (n = 3) are expressed as means ± SD means in column (^{ab}) letter differ (p ≤ 0.05) using Tukey's Studentized range test. (1) FRAP is expressed in μmol FeSO₄ per g of dry flowers, (2) TEAC is expressed in mM Trolox per g of dry flowers. BM: Barbara mitchell, EE: Elegant explosion, DPPH: 2,2-diphenyl-1-picrylhydrazyl, FRAP: Ferric reducing antioxidant power, Fe²⁺: Ferrous ion, TEAC: Trolox equivalent antioxidant capacity

mechanisms of antioxidant action^{62,63}. The results in Table 2 show EE had a significantly (p ≤ 0.05) higher FRAP value (2.48 μmol FeSO₄ g⁻¹ dwb) compared to BM (0.97 μmol FeSO₄ g⁻¹ dwb). TEAC activity was higher (p ≤ 0.05) in EE (16.15 mM Trolox g⁻¹ dwb) compared to BM (4.82 mM Trolox g⁻¹ dwb). The DPPH assay is a rapid, direct and reproducible procedure widely used to determine the radical scavenging activity, via inhibition (%) or IC₅₀ of plant extracts. The IC₅₀ was 14.31 mg mL⁻¹ (R² = 0.94171) in EE compared to 24 mg mL⁻¹ (R² = 0.99363) in BM in Table 2. The data of Fig. 1 shows the DPPH expressed as percentage (%) inhibition after 90 min (T₉₀) was higher (p ≤ 0.05) in EE (75.37%) than in BM (51.56%). These results are comparable with other studies investigating the antioxidant activities of edible flowers^{3,64,65}. The IC₅₀ for Fe²⁺ capacity was indicated at 43.08 (R² = 0.82275) and 37.05 mg mL⁻¹ (R² = 0.70796) for BM and EE, respectively. In Fig. 2 the Metal chelating (Fe²⁺) capacity in the presence of

BM and EE extracts showed a linear activity with increasing concentration. Although this study did not record the statistical correlation between polyphenol contents and antioxidant activity, many authors^{6,66,67} have indicated a strong linear relation between the polyphenol contents and the antioxidant activities of edible flower species.

Selected polyphenols in daylily flowers: HPLC chromatogram of selected polyphenol standards are shown in Fig. 3. The standards are chlorogenic acid (RT = 6.62 min), (+)-catechin (RT = 4.43 min), gallic acid (RT = 4.45 min) and rutin (RT = 15.04 min). The peaks were confirmed by UV/Vis spectra obtained at λ_{max} 284 nm for (+)-catechin (Fig. 3a) and gallic acid (Fig. 3b), 320 nm for chlorogenic acid (Fig. 3c) and 350 nm for rutin (Fig. 3d). The chromatogram in Fig. 4 shows the peaks for the polyphenolic compounds identified in daylily flower extracts. In Fig. 4a, the chromatogram show (+)-catechin,




Fig. 2: Fe²⁺ chelating activity of daylily cultivars

Values are expressed as means ± SD and were conducted in triplets in three independent experiments. Statistical significance was determined by one-way ANOVA followed by Tukey's range test. EE: Elegant explosion, BM: Barbara mitchell




Fig. 3(a-d): HPLC Chromatographs of polyphenolic standards catechin, gallic acid, chlorogenic acid and rutin

UV spectrum was recorded for (a) Catechin (284 nm), (b) Gallic Acid (284 nm), (c) Chlorogenic acid (320 nm) and (d) Rutin (350 nm)




Fig. 4(a-b): HPLC Chromatograms and UV spectrums of selected polyphenols identified in daylily cultivars
 (a) Catechin, chlorogenic acid and rutin were identified in Elegiant Explosion and (b) Catechin and rutin were identified in Barbara Mitchell

Table 3: Concentrations of selected polyphenol compounds ($\mu\text{g g}^{-1}$ dwb) identified in daylily cultivars

	Catechin ($\mu\text{g g}^{-1}$)	Rutin ($\mu\text{g g}^{-1}$)	Chlorogenic acid ($\mu\text{g g}^{-1}$)	Gallic acid ($\mu\text{g g}^{-1}$)
BM	13 ^b	11 ^b	ND	ND
EE	83 ^a	42 ^a	37	ND

Values (n = 3) are expressed as means \pm SD of three replicates. Means in column (^{ab}) letter differ ($p < 0.05$) using Tukey's Studentized range test. EE: Elegiant explosion, BM: Barbara mitchell, ND: Not detected, dwb: Dry weight basis

chlorogenic acid and rutin were identified in EE. The data of Fig. 4b shows (+)- catechin and chlorogenic were identified in BM. Surprisingly; gallic acid was below the quantification limit and was not detected in any of the flowers in our study. The data in Table 3 shows concentrations of polyphenols in EE and BM. (+)-Catechin and rutin in EE were six times and four times higher compared to BM. Chlorogenic acid was only identified in EE ($37 \mu\text{g g}^{-1}$ dwb). The results show that (+)-catechin was the most represented polyphenol in both the daylily cultivars (83 and $13 \mu\text{g g}^{-1}$ for EE and BM, respectively). It is reported that in daylily flowers, (+)-catechin represents nearly 75% of all polyphenols^{14,68}. The high (+)-catechin, chlorogenic acid and rutin concentration in EE may be responsible for the

demonstrated antioxidant activity. Polyphenols such as chlorogenic acid, rutin and (+)-catechin possess several physiological activities. These compounds are shown to inhibit the proliferation of cancer cells⁶⁹⁻⁷². They also have cardioprotective^{73,74}, anti-obesity⁷⁵ and anti-hypertensive properties⁷⁶⁻⁷⁹. Due to the polyphenol contents, the daylily flowers used in this study could be considered as ingredients for several types of functional food applications with health benefits or as a nutraceutical in dietary supplements. Even though there were some variations in polyphenolic composition compared to other studies⁸⁰⁻⁸², these variabilities might be due to the daylily flower cultivars, experimental conditions and the type of solvent used for polyphenolic




Fig. 5: Cytotoxicity of elegant explosion extracts in HepG2 cells after 24 and 48 hrs exposure

Values (n = 3) are expressed as means \pm SD of three replicates. EE: Elegant explosion

extractions. All the same, the polyphenol contents estimated in our study are in the range or comparable to those reported by some investigators^{14,30,41,66,83,84}.

While gallic acid is one of the most common phenolic acids in edible plants, we did not detect this phenolic compound in our samples. Gallic acid and catechin share some structural similarities and as a result, both compounds could have co-eluted together under the HPLC conditions in this study. In any case, our results correspond with Zheng *et al.*⁸⁵ and Barros *et al.*⁸⁴ who reported similar findings in edible flowers from China and Brazil, respectively. Similar results were also observed with *Calendula officinalis* L., *Chrysanthemum morifolium*, *Jasminum sambac* (L.) Ait and other edible flowers⁸³. Wang *et al.*⁸⁶ reported that good separation of gallic acid and catechin is best achieved if the concentration of ethanol in the final sample solution for HPLC analysis is less than 15%. In this study, the final sample solution of ethanol was more than 15%.

Overall, the results show EE flowers contained high TPC, TFC, TMA CTC and strong antioxidant properties compared to BM flowers. As such, EE flowers were selected and used to perform *in vitro* toxicity experiments.

Cytotoxicity of Elegant Explosion (EE) extracts in HepG2 cells:

The cytotoxicity of EE extract in Hep2G cells was achieved by the PrestoBlue[®] test and results are presented in Fig. 5. The PrestoBlue[®] assay is a simple and fast live assay to monitor cell viability and cytotoxicity⁸⁷. It is a resazurin-based compound that is quickly reduced by metabolically viable cells^{88,89}. Based on the results in Fig. 5, EE extract was non-cytotoxic activity towards HepG2 cells after 24 hrs exposure, as cytotoxicity ranged from 73-100% (8-0.2 mg mL⁻¹). The

results also show cell viability was slightly decreased with prolonged exposure (48 hrs incubation). After 48 hrs incubation cytotoxicity was 59% at the highest (8 mg mL⁻¹) and the IC₅₀ was recorded at 10.13 mg mL⁻¹. The present results indicate that hydroethanolic EE extract possesses weak to moderate toxicity towards the HepG2 cell line used in this study. This assertion is based on ISO 10993-5⁹⁰ which specifies that cell viability above 80% is non-cytotoxic; 80-60% is weakly cytotoxic; 60-40% is moderately cytotoxic and <40% is strongly cytotoxic. The present results are comparable to the previous observations^{7,91,92}. The effect could be attributed to the specific phenolic compounds and antioxidant activity of EE. Studies conducted by Cichewicz *et al.*⁹³ and Lin *et al.*⁹⁴ showed that polyphenols compounds anthraquinones and caffeoylquinic acid derivatives from daylily flowers showed weak cytotoxic activities and prevented oxidative stress in the HepG2 cell line. Recent research indicated caffeoylquinic acid derivatives and flavanone glycosides in *Chrysanthemum morifolium* displayed hepatoprotective effects in HepG2 cells⁹⁵. Kwon *et al.*⁹⁶ also reported quercetin and kaempferol in *Begonia semperflorens* flowers proved to be protective against oxidative stress in t BHP induced HepG2.

Morphological changes in Hep2G cells after 48h incubation:

The result of Fig. 6a-d show microscopic observations in Hep2G cells exposed to EE extract after 48 hrs incubation. The results show EE extract induced high apoptosis with increasing concentration (0-4 mg mL⁻¹). Chromatin condensation and fragmentations were observed in the HepG2 cells treated with EE extract, while the control untreated cells displayed minimal intact nuclei or apoptosis-inducing effects. Besides, microscopic observation showed




Fig. 6(a-d): Elegant explosion (EE) extract induced morphological changes in Hep2G cells following 48 hrs incubation

(a) Phase-contrast photomicrographs of Hep2G cells treated with EE extract at 0 (control)-4 mg mL⁻¹ for 48 hrs, (b-c) Cytotoxicity of EE extract analyzed by live/dead assay. NucBlue® Live (Hoechst 33342) reagent stains the nuclei of all cells while NucGreen® Dead reagent stains only the nuclei of dead cells. A higher percentage of apoptotic cells (live (blue)/dead (green)) are observed in the treated cells than in the control. Green channel signal intensity (dead cell stain) is strong in the cells treated with a high concentration of EE extract. Images are representative of three independent experiments. Cells were imaged on the EVOS® FL Auto system. Images were captured at ×10 magnification. Scale bars: 400

cell confluence was markedly reduced with increasing concentration. Furthermore, EE extract induced obvious morphology changes including cell shrinkage, abnormality and death as compared with control untreated cells (Fig. 6a). It is important to point out that maintenance of membrane integrity is a standard criterion for cell viability. NucBlue® Live (Hoechst 33342) reagent stains the nuclei of all cells (Fig. 6b).

Live cells react with the fluorescent reactive dye only on their surface to yield weakly fluorescent cells. The nuclei of dead cells with compromised membranes react with NucGreen® Dead reagent stains only and produce brightly stained green cells (Fig. 6c). The results from cell viability analysis, i.e., presto blue are in line with cell images. The data of Fig. 6d depicts the merge or overlay of Fig. 6a-c.

Table 4: Effects of elegant explosion daylily flowers extracts on endogenous antioxidant enzymes in HepG2 cells

	SOD (U mg ⁻¹ protein)	GST (μmol mg ⁻¹ protein)	GSH (μmol mg ⁻¹ protein)
24 hrs endogenous enzyme activity			
Control	1.11±0.06 ^c	6.36±0.31 ^a	0.27±0.01 ^b
0.1	1.4±0.04 ^a	5.84±0.07 ^b	0.40±0.02 ^a
0.2	1.48±0.03 ^a	4.60±0.13 ^{bc}	0.30±0.01 ^a
0.4	1.46±0.03 ^a	4.11±0.12 ^{bc}	0.32±0.05 ^a
0.8	1.47±0.02 ^a	4.0±0.16 ^{bc}	0.40±0.01 ^a
1	1.45±0.05 ^a	3.57±0.15 ^c	0.34±0.02 ^a
2	1.43±0.02 ^a	3.0±0.11 ^c	0.35±0.02 ^a
4	1.32±0.07 ^b	3.54±0.23 ^c	0.37±0.03 ^a
48 hrs endogenous enzyme activity			
Control	0.88±0.07 ^c	4.08±0.53 ^{ab}	0.14±0.01 ^b
0.1	1.15±0.01 ^b	4.68±1.05 ^a	0.40±0.01 ^a
0.2	1.32±0.03 ^a	3.87±0.38 ^{ab}	0.38±0.01 ^a
0.4	1.28±0.04 ^a	3.43±0.24 ^b	0.35±0.01 ^a
0.8	1.22±0.06 ^a	3.11±0.18 ^b	0.31±0.01 ^a
1	1.17±0.05 ^b	3.01±0.05 ^b	0.41±0.01 ^a
2	1.13±0.03 ^b	2.83±0.15 ^b	0.48±0.01 ^a
4	1.02±0.06 ^{bc}	2.97±0.12 ^b	0.46±0.01 ^a

Values are expressed as means ± SD and were conducted in triplicates. ^{abc}Letters demonstrate the significant difference ($p \leq 0.05$), comparing the concentrations of the assay at 24 and 48 hrs. Statistical significance was determined by one-way ANOVA followed by Tukey's range test. SOD: Superoxide dismutase, GSH: Glutathione, GST: Glutathione-S-Transferase

Effect of EE extract on redox biomarkers in HepG2 cells:

The result of Table 4 shows the effect of EE extract on endogenous enzyme activity. SOD activity (U mg⁻¹ protein) was increased ($p \leq 0.05$) in HepG2 cells exposed to EE extract (1.32-1.48 U mg⁻¹ protein after 24 hrs incubation and 1.02-1.32 U mg⁻¹ protein after 48 hrs incubation) compared to control untreated cells (1.11 U mg⁻¹ protein and 0.88 U mg⁻¹ protein, respectively for 24 and 48 hrs). The results show SOD activity was significantly ($p \leq 0.05$) different after 24 and 48 hrs incubation. While the present data show a high SOD activity after treatment with EE extract, there was no linearity regarding concentrations. Other studies have indicated that SOD activity increase when treated with polyphenols^{97,98}. SOD is a family of antioxidant enzymes that convert harmful superoxide radicals into H₂O₂, which is subsequently metabolized to water and oxygen by catalase and glutathione peroxidase. The present results could be due to the high antioxidant activity of the EE extract directly retarding the superoxide radical's formation. It is also possible that the EE extract activated the SOD enzyme that catalyzes the conversion of superoxide radicals into hydrogen peroxide and oxygen for subsequent detoxification to water and oxygen.

GST activity (μmol mg⁻¹ protein) decreased ($p \leq 0.05$) in HepG2 cells exposed to EE extract for 24 hrs compared with control shown in Table 4. However, the exposure of cells for 48 hrs produced no significant changes compared to the control. The results are in line with others who reported decreased GST activity in HepG2 cells after acute incubation with basil and ginger extracts^{33,97} and grape pomace extract⁹⁹.

Also, previous studies have detected no change in GST activity after prolonged incubation with polyphenolic compounds in HepG2 cells¹⁰⁰⁻¹⁰².

The results of the GSH level shown in Table 4 followed a similar pattern observed for SOD activity. The results demonstrated no significant difference in GSH levels as EE extract concentration increased. A study by Franco and Cidowski¹⁰³ indicated that GSH depletion is a common feature of apoptotic cell death. GSH participates in the disposal of potentially harmful electrophiles and protects cells from the damaging effects of free radicals^{104,105}. Therefore, its depletion renders cells particularly vulnerable to oxidative stress. Since these results indicated no significant associations between cell viability and cellular GSH level, this phenomenon could be due to the tendency for cultured cells to increase protection as the concentration of extracts is increased¹⁰⁶. Some studies utilizing high concentrations of plant extracts have reported increased GSH levels⁹⁷. Others have suggested that GSH synthesis could be up-regulated during oxidative stress by phenolic compounds if their concentration does not compromise cell viability¹⁰⁷. This might finally contribute to the paradox that is associated with the protective mechanisms of GSH.

CONCLUSION

A preliminary study of herbal extracts from daylily flower cultivars Elegant Explosion (EE) and Barbara Mitchell (BM) were evaluated for their polyphenol content and antioxidant activity. Also, the *in vitro* assessment of the safety

(cytotoxicity/cell viability) and bioefficacy (effect on endogenous antioxidant enzymes) was evaluated for EE extract. According to results EE extracts exhibited higher TPC, TFC, TMA, polyphenol contents and strong free radical scavenging inhibitory activities. Furthermore, EE extracts induced weak to moderate cytotoxicity.

SIGNIFICANCE STATEMENT

The extent of this study is to utilize daylilies for use as functional food ingredients. The results generated from this study will help in gathering baseline information on the potential of daylily flowers to be utilized in the preparation of nutraceutical, functional food products and dietary supplements.

ACKNOWLEDGMENT

This study was funded by the USDA capacity Building grant No. 11343134, Evans Allen project No: ALAX-012-0812 and Alabama A and M University Agriculture Research Station, Normal AL 35762.

REFERENCES

1. Friedman, H., O. Agami, Y. Vinokur, S. Droby and L. Cohen *et al.*, 2010. Characterization of yield, sensitivity to *Botrytis cinerea* and antioxidant content of several rose species suitable for edible flowers. *Sci. Hortic.*, 123: 395-401.
2. Rop, O., J. Mlcek, T. Jurikova, J. Neugebauerova and J. Vabkova, 2012. Edible flowers-A new promising source of mineral elements in human nutrition. *Molecules*, 17: 6672-6683.
3. Fernandes, L., S. Casal, J.A. Pereira, J.A. Saraiva and E. Ramalhosa, 2017. Edible flowers: A review of the nutritional, antioxidant, antimicrobial properties and effects on human health. *J. Food Compos. Anal.*, 60: 38-50.
4. Newman, M. and C.L. Kirker, 2016. *Edible Flowers: A Global History*. 1st Edn., Reaktion Books Ltd., London, pp: 19-21.
5. Stefaniak, A. and M.E. Grzeszczuk, 2018. Nutritional and biological value of five edible flower species. *Not. Bot. Horti. Agrobi. Cluj-Napoca*, 47: 128-134.
6. Navarro-González, I., R. González-Barrio, V. García-Valverde, A.B. Bautista-Ortín and M.J. Periago, 2015. Nutritional composition and antioxidant capacity in edible flowers: Characterisation of phenolic compounds by HPLC-DAD-ESI/MSn. *Int. J. Mole. Sci.*, 16: 805-822.
7. Pires, T.C.S.P., M.I. Dias, L. Barros, R.C. Calhella and M.J. Alves *et al.*, 2018. Edible flowers as sources of phenolic compounds with bioactive potential. *Food Res. Int.*, 105: 580-588.
8. Pinedo-Espinoza, J.M., J. Gutiérrez-Tlahque, Y.O. Santiago-Saenz, C.L. Aguirre-Mancilla and M. Reyes-Fuentes *et al.*, 2020. Nutritional composition, bioactive compounds and antioxidant activity of wild edible flowers consumed in semi-arid regions of Mexico. *Plant Foods Hum. Nutr.*, 75: 413-419.
9. Cunningham, E., 2015. What nutritional contribution do edible flowers make? *J. Acad. Nut. Diet.*, Vol. 115. 10.1016/j.jand.2015.03.002.
10. Loizzo, M.R., A. Pugliese, M. Bonesi, M.C. Tenuta, F. Menichini, J. Xiao and R. Tundis, 2016. Edible flowers: A rich source of phytochemicals with antioxidant and hypoglycemic properties. *J. Agric. Food Chem.*, 64: 2467-2474.
11. Shi, J., J. Gong, J. Liu, X. Wu and Y. Zhang, 2009. Antioxidant capacity of extract from edible flowers of *Prunus mume* in China and its active components. *LWT-Food Sci. Technol.*, 42: 477-482.
12. Kao, F.J., W.D. Chiang and H.M. Liu, 2015. Inhibitory effect of daylily buds at various stages of maturity on nitric oxide production and the involved phenolic compounds. *LWT-Food Sci. Technol.*, 61: 130-137.
13. Liu, X.L., L. Luo, B.B. Liu, J. Li, D. Geng, Q. Liu and L.T. Yi, 2014. Ethanol extracts from *Hemerocallis citrina* attenuate the upregulation of proinflammatory cytokines and indoleamine 2,3-dioxygenase in rats. *J. Ethnopharmacol.*, 153: 484-490.
14. Fu, M., Z. He, Y. Zhao, J. Yang and L. Mao, 2009. Antioxidant properties and involved compounds of daylily flowers in relation to maturity. *Food Chem.*, 114: 1192-1197.
15. Que, F., L.C. Mao and X.J. Zheng, 2007. *In vitro* and *vivo* antioxidant activities of daylily flowers and the involvement of phenolic compounds. *Asia Pac. J. Clin. Nutr.*, 16: 196-203.
16. Mueller, D.S., J.L. Williams-Woodward and J.W. Buck, 2003. Resistance of daylily cultivars to the daylily rust pathogen, *Puccinia hemerocallidis*. *HortScience*, 38: 1137-1140.
17. Rodriguez-Enriquez, M.J. and R.T. Grant-Downton, 2013. A new day dawning: *Hemerocallis* (daylily) as a future model organism. *Ann. Bot. Plants*, Vol. 5. 10.1093/aobpla/pls055.
18. Granato, D., J.S. Santos, L.G. Maciel and D.S. Nunes, 2016. Chemical perspective and criticism on selected analytical methods used to estimate the total content of phenolic compounds in food matrices. *TrAC Trends Anal. Chem.*, 80: 266-279.
19. Gajula, D., M. Verghese, J. Boateng, L.T. Walker, L. Shackelford, S.R. Mentreddy and S. Cedric, 2009. Determination of total phenolics, flavonoids and antioxidant and chemopreventive potential of basil (*Ocimum basilicum* L. and *Ocimum tenuiflorum* L.). *Int. J. Cancer Res.*, 5: 130-143.
20. Kim, D.O., O.K. Chun, Y.J. Kim, H.Y. Moon and C.Y. Lee, 2003. Quantification of polyphenolics and their antioxidant capacity in fresh plums. *J. Agric. Food Chem.*, 51: 6509-6515.
21. Luna-Vital, D., Q. Li, L. West, M. West and E.G. de Mejia, 2017. Anthocyanin condensed forms do not affect color or chemical stability of purple corn pericarp extracts stored under different pHs. *Food Chem.*, 232: 639-647.

22. Lee, J., R.W. Durst and R.E. Wrolstad, 2005. Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants and wines by the pH differential method: Collaborative study. *J. AOAC Int.*, 88: 1269-1278.
23. Montoro, P., A. Braca, C. Pizza and N. De Tommasi, 2005. Structure-oxidant activity relationship of flavonoids isolated from different plant species. *Food Chem.*, 92: 349-355.
24. Benzie, I.F.F. and J.J. Strain, 1996. The Ferric Reducing Ability of Plasma (FRAP) as a measure of antioxidant power: The FRAP assay. *Anal. Biochem.*, 239: 70-76.
25. Brand-Williams, W., M.E. Cuvelier and C. Berset, 1995. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci. Technol.*, 28: 25-30.
26. Muñoz-Espada, A.C., K.V. Wood, B. Bordelon and B.A. Watkins, 2004. Anthocyanin quantification and radical scavenging capacity of concord, norton and marechal foch grapes and wines. *J. Agric. Food Chem.*, 52: 6779-6786.
27. Wong, F.C., A.L. Yong, E.P.S. Ting, S.C. Khoo, H.C. Ong and T.T. Chai, 2014. Antioxidant, metal chelating, anti-glucosidase activities and phytochemical analysis of selected tropical medicinal plants. *Iran. J. Pharm. Res.*, 13: 1409-1415.
28. Schutz, K., D. Kammerer, R. Carle and A. Schieber, 2004. Identification and quantification of caffeoylquinic acids and flavonoids from artichoke (*Cynara scolymus* L.) heads, juice and pomace by HPLC-DAD-ESI/MSⁿ. *J. Agric. Food Chem.*, 52: 4090-4096.
29. Kaisoon, O., S. Siriamornpun, N. Weerapreeyakul and N. Meeso, 2011. Phenolic compounds and antioxidant activities of edible flowers from Thailand. *J. Funct. Foods*, 3: 88-99.
30. Kaisoon, O., I. Konczak and S. Siriamornpun, 2012. Potential health enhancing properties of edible flowers from Thailand. *Food Res. Int.*, 46: 563-571.
31. Xiong, L., J. Yang, Y. Jiang, B. Lu and Y. Hu *et al.*, 2014. Phenolic compounds and antioxidant capacities of 10 common edible flowers from China. *J. Food Sci.*, 79: C517-C525.
32. Liu, W., Y. Zhao, J. Sun, G. Li, Y. Shan and P. Chen, 2017. Study the effects of drying processes on chemical compositions in daylily flowers using flow injection mass spectrometric fingerprinting method and chemometrics. *Food Res. Int.*, 102: 493-503.
33. Yang, M., R. Hardin, S. Ogutu, M. Verghese and J. Boateng, 2016. Preliminary analysis of *in vitro* digestion and bioactivity assessment of basil and ginger in human liver cancer cell line. *J. Biol. Sci.*, 16: 202-214.
34. Boateng, J. and M. Verghese, 2012. Protective effects of the phenolic extracts of fruits against oxidative stress in human lung cells. *Int. J. Pharmacol.*, 8: 152-160.
35. Griffith, O.W. and A. Meister, 1985. Origin and turnover of mitochondrial glutathione. *Proc. Natl. Acad. Sci. USA*, 82: 4668-4672.
36. Fernandez-Urrusuno, R., E. Fattal, J. Feger, P. Couvreur and P. Therond, 1997. Evaluation of hepatic antioxidant systems after intravenous administration of polymeric nanoparticles. *Biomaterials*, 18: 511-517.
37. Habig, W.H., M.J. Pabst and W.B. Jakoby, 1974. Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *J. Biol. Chem.*, 249: 7130-7139.
38. Barriada-Bernal, L.G., N. Almaraz-Abarca, E.A. Delgado-Alvarado, T. Gallardo-Velázquez and J.A. Ávila-Reyes *et al.*, 2014. Flavonoid composition and antioxidant capacity of the edible flowers of agave durangensis (Agavaceae). *CyTA J. Food*, 12: 105-114.
39. Tian, H., F.F. Yang, C.Y. Liu, X.M. Liu and R.L. Pan *et al.*, 2017. Effects of phenolic constituents of daylily flowers on corticosterone- and glutamate-treated PC12 cells. *BMC Comp. Altern. Med.*, Vol. 17. 10.1186/s12906-017-1582-x.
40. Wang, Y., T. Xu, B. Fan, L. Zhang, C. Lu, D. Wang and F. Wang, 2018. Advances in researches on chemical composition and functions of *Hemerocallis* plants. *Med. Plant*, 9: 16-21.
41. Wu, W.T., M.C. Mong, Y.C. Yang, Z.H. Wang and M.C. Yin, 2018. Aqueous and ethanol extracts of daylily flower (*Hemerocallis fulva* L.) protect HUVE cells against high glucose. *J. Food Sci.*, 83: 1463-1469.
42. Hernandez, I., L. Alegre, F. Van Breusegem and S. Munne-Bosch, 2009. How relevant are flavonoids as antioxidants in plants? *Trends Plant Sci.*, 14: 125-132.
43. Pérez-Jiménez, J. and F. Saura-Calixto, 2018. Fruit peels as sources of non-extractable polyphenols or macromolecular antioxidants: Analysis and nutritional implications. *Food Res. Int.*, 111: 148-152.
44. Landi, M., B. Ruffoni, L. Guidi, M. Savona and D. Salvi, 2015. Cold storage does not affect ascorbic acid and polyphenolic content of edible flowers of a new hybrid sage. *Agrochimica*, 59: 348-357.
45. Yang, H. and Y. Shin, 2017. Antioxidant compounds and activities of edible roses (*Rosa hybrida* spp.) from different cultivars grown in Korea. *Appl. Biol. Chem.*, 60: 129-136.
46. Cendrowski, A., I. Ścibisz, M. Mitek, M. Kieliszek and J. Kolniak-Ostek, 2017. Profile of the phenolic compounds of *Rosa rugosa* petals. *J. Food Qual.*, Vol. 2017. 10.1155/2017/7941347.
47. Chensom, S., Y. Shimada, H. Nakayama, K. Yoshida and T. Kondo *et al.*, 2020. Determination of anthocyanins and antioxidants in 'titanicus' edible flowers *in vitro* and *in vivo*. *Plant Foods Hum. Nutr.*, 75: 265-271.
48. Chaiyasut, C., B.S. Sivamaruthi, N. Pengkumsri, S. Sirilun, S. Peerajan, K. Chaiyasut and P. Kesika, 2016. Anthocyanin profile and its antioxidant activity of widely used fruits, vegetables and flowers in thailand. *Asian J. Pharmaceut. Clin. Res.*, 9: 218-224.

49. Grajeda-Iglesias, C., M.C. Figueroa-Espinoza, N. Barouh, B. Baréa, A. Fernandes, V. de Freitas and E. Salas, 2016. Isolation and characterization of anthocyanins from *Hibiscus sabdariffa* flowers. *J. Nat. Prod.*, 79: 1709-1718.
50. Benvenuti, S., E. Bortolotti and R. Maggini, 2016. Antioxidant power, anthocyanin content and organoleptic performance of edible flowers. *Sci. Hort.*, 199: 170-177.
51. Wang, L.S. and G.D. Stoner, 2008. Anthocyanins and their role in cancer prevention. *Cancer Lett.*, 269: 281-290.
52. He, J. and M.M. Giusti, 2010. Anthocyanins: Natural colorants with health-promoting properties. *Annu. Rev. Food Sci. Technol.*, 1: 163-187.
53. Azzini, E., J. Giacometti and G.L. Russo, 2017. Antiobesity effects of anthocyanins in preclinical and clinical studies. *Oxid. Med. Cell Longev.*, Vol. 2017. 10.1155/2017/2740364.
54. Tsang, C., N.F. Smail, S. Almoosawi, G.J.M. McDougall and E.A.S. Al-Dujaili, 2018. Antioxidant rich potato improves arterial stiffness in healthy adults. *Plant Foods Hum. Nutr.*, 73: 203-208.
55. Hester, S.N., A. Mastaloudis, R. Gray, J.M. Antony, M. Evans and S.M. Wood, 2018. Efficacy of an anthocyanin and prebiotic blend on intestinal environment in obese male and female subjects. *J. Nutr. Metab.*, Vol. 2018. 10.1155/2018/7497260.
56. Vugic, L., N. Colson, E. Nikbakht, A. Gaiz, O.J. Holland, A.R. Kundur and I. Singh, 2020. Anthocyanin supplementation inhibits secretion of pro-inflammatory cytokines in overweight and obese individuals. *J. Funct. Foods*, Vol. 64. 10.1016/j.jff.2019.103596.
57. Rauf, A., M. Imran, T. Abu-Izneid, lahtisham-Ul-Haq and S. Patel *et al.*, 2019. Proanthocyanidins: A comprehensive review. *Biomed. Pharmacother.*, Vol. 116. 10.1016/j.biopha.2019.108999.
58. Tao, W., Y. Zhang, X. Shen, Y. Cao, J. Shi, X. Ye and S. Chen, 2019. Rethinking the mechanism of the health benefits of proanthocyanidins: Absorption, metabolism and interaction with gut microbiota. *Compr. Rev. Food Sci. Food Saf.*, 18: 971-985.
59. Unusan, N., 2020. Proanthocyanidins in grape seeds: An updated review of their health benefits and potential uses in the food industry. *J. Funct. Foods*, Vol. 67. 10.1016/j.jff.2020.103861.
60. Coleman, S.L. and O.M. Shaw, 2017. Progress in the understanding of the pathology of allergic asthma and the potential of fruit proanthocyanidins as modulators of airway inflammation. *Food Funct.*, 8: 4315-4324.
61. Lai, R., D. Xian, X. Xiong, L. Yang, J. Song and J. Zhong, 2018. Proanthocyanidins: Novel treatment for psoriasis that reduces oxidative stress and modulates Th17 and Treg cells. *Redox Rep.*, 23: 130-135.
62. González-Barrío, R., M.J. Periago, C. Luna-Recio, F.J. Garcia-Alonso and I. Navarro-González, 2018. Chemical composition of the edible flowers, pansy (*Viola wittrockiana*) and snapdragon (*Antirrhinum majus*) as new sources of bioactive compounds. *Food Chem.*, 252: 373-380.
63. Frankle, E.N. and A.S. Meyer, 2000. The problems of using one-dimensional methods to evaluate multifunctional food and biological antioxidants. *J. Sci. Food Agric.*, 80: 1925-1941.
64. Teerarak, M., K. Changsawake, K. Pilasombut and C. Laosinwattana, 2018. Antioxidant activities and heat stability of edible flowers of *Telosma minor* and *Sesbania javanica*. *J. Herbs Spices Med. Plants*, 24: 87-98.
65. Zheng, J., X. Yu, M. Maninder and B. Xu, 2018. Total phenolics and antioxidants profiles of commonly consumed edible flowers in China. *Int. J. Food Prop.*, 21: 1524-1540.
66. Li, A.N., S. Li, H.B. Li, D.P. Xu, X.R. Xu and F. Chen, 2014. Total phenolic contents and antioxidant capacities of 51 edible and wild flowers. *J. Funct. Foods*, 6: 319-330.
67. Garzón, G.A., D.C. Manns, K. Riedl, S.J. Schwartz and O. Padilla-Zakour, 2015. Identification of phenolic compounds in petals of nasturtium flowers (*Tropaeolum majus*) by high-performance liquid chromatography coupled to mass spectrometry and determination of oxygen radical absorbance capacity (ORAC). *J. Agric. Food Chem.*, 63: 1803-1811.
68. Mlcek, J. and O. Rop, 2011. Fresh edible flowers of ornamental plants-A new source of nutraceutical foods. *Trends Food Sci. Technol.*, 22: 561-569.
69. Mayr, C., A. Wagner, D. Neureiter, M. Pichler and M. Jakab *et al.*, 2015. The green tea catechin epigallocatechin gallate induces cell cycle arrest and shows potential synergism with cisplatin in biliary tract cancer cells. *BMC Comp. Altern. Med.*, Vol. 15. 10.1186/s12906-015-0721-5.
70. Bimonte, S., M. Cascella, M. Leongito, R. Palaia, D. Caliendo, F. Izzo and A. Cuomo, 2017. An overview of pre-clinical studies on the effects of (-)-epigallocatechin-3-gallate, a catechin found in green tea, in treatment of pancreatic cancer. *Recenti progressi Med.*, 108: 282-287.
71. Hou, N., N. Liu, J. Han, Y. Yan and J. Li, 2017. Chlorogenic acid induces reactive oxygen species generation and inhibits the viability of human colon cancer cells. *Anti-cancer Drugs*, 28: 59-65.
72. Yan, Y., N. Liu, N. Hou, L. Dong and J. Li, 2017. Chlorogenic acid inhibits hepatocellular carcinoma *in vitro* and *in vivo*. *J. Nut. Biochem.*, 46: 68-73.
73. Prince, P.S.M. and S. Priya, 2010. Preventive effects of rutin on lysosomal enzymes in isoproterenol induced cardio toxic rats: Biochemical, histological and *in vitro* evidences. *Eur. J. Pharmacol.*, 649: 229-235.
74. Mills, C.E., A. Flury, C. Marmet, L. Poquet and S.F. Rimoldi *et al.*, 2017. Mediation of coffee-induced improvements in human vascular function by chlorogenic acids and its metabolites: Two randomized, controlled, crossover intervention trials. *Clin. Nut.*, 36: 1520-1529.
75. Santana-Gálvez, J., L. Cisneros-Zevallos and D. Jacobo-Velázquez, 2017. Chlorogenic acid: Recent advances on its dual role as a food additive and a nutraceutical against metabolic syndrome. *Molecules*, Vol. 22. 10.3390/molecules22030358.

76. Cai, E.P. and J.K. Lin, 2009. Epigallocatechin gallate (EGCG) and rutin suppress the glucotoxicity through activating IRS2 and AMPK signaling in rat pancreatic β cells. *J. Agric. Food Chem.*, 57: 9817-9827.
77. Khalesi, S., J. Sun, N. Buys, A. Jamshidi, E. Nikbakht-Nasrabadi and H. Khosravi-Boroujeni, 2014. Green tea catechins and blood pressure: A systematic review and meta-analysis of randomised controlled trials. *Eur. J. Nutr.*, 53: 1299-1311.
78. Onakpoya, I.J., E.A. Spencer, M.J. Thompson and C.J. Heneghan, 2015. The effect of chlorogenic acid on blood pressure: A systematic review and meta-analysis of randomized clinical trials. *J. Hum. Hypertens.*, 29: 77-81.
79. Ward, N.C., J.M. Hodgson, R.J. Woodman, D. Zimmermann and L. Poquet *et al.*, 2016. Acute effects of chlorogenic acids on endothelial function and blood pressure in healthy men and women. *Food Funct.*, 7: 2197-2203.
80. Szewczyk, K., M. Miazga-Karska, W. Pietrzak, Ł. Komsta, B. Krzemińska and A. Grzywa-Celińska 2020. Phenolic composition and skin-related properties of the aerial parts extract of different *Hemerocallis* cultivars. *Antioxidants*, Vol. 9. 10.3390/antiox9080690.
81. Lin, S.H., H.C. Chang, P.J. Chen, C.L. Hsieh, K.P. Su and L.Y. Sheen, 2013. The antidepressant-like effect of ethanol extract of daylily flowers (*Jin Zhēn Huā*) in rats. *J. Traditional Complementary Med.*, 3: 53-61.
82. Joshi, R. and A. Gulati, 2011. Biochemical attributes of tea flowers (*Camellia sinensis*) at different developmental stages in the Kangra region of India. *Sci. Hortic.*, 130: 266-274.
83. Chen, G.L., S.G. Chen, Y.Q. Xie, F. Chen, Y.Y. Zhao, C.X. Luo and Y.Q. Gao, 2015. Total phenolic, flavonoid and antioxidant activity of 23 edible flowers subjected to *in vitro* digestion. *J. Funct. Foods*, 17: 243-259.
84. Barros, R.G.C., J.K.S. Andrade, U.C. Pereira, C.S. de Oliveira and Y.R.R.S. Rezende *et al.*, 2020. Phytochemicals screening, antioxidant capacity and chemometric characterization of four edible flowers from Brazil. *Food Res. Int.*, Vol. 130. 10.1016/j.foodres.2019.108899.
85. Zheng, J., M. Meenu and B. Xu, 2019. A systematic investigation on free phenolic acids and flavonoids profiles of commonly consumed edible flowers in China. *J. Pharm. Biomed. Anal.*, 172: 268-277.
86. Wang, H., K. Helliwell and X. You, 2000. Isocratic elution system for the determination of catechins, caffeine and gallic acid in green tea using HPLC. *Food Chem.*, 68: 115-121.
87. Emter, R. and A. Natsch, 2015. A fast resazurin-based live viability assay is equivalent to the MTT-test in the KeratinoSens assay. *Toxicol. in vitro.*, 29: 688-693.
88. O'Brien, J., I. Wilson, T. Orton and F. Pognan, 2000. Investigation of the alamar blue (resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity. *Eur. J. Biochem.*, 267: 5421-5426.
89. Boncler, M., M. Rozalski, U. Krajewska, A. Podsedek and C. Watala, 2014. Comparison of PrestoBlue and MTT assays of cellular viability in the assessment of anti-proliferative effects of plant extracts on human endothelial cells. *J. Pharmacol. Toxicol. Meth.*, 69: 9-16.
90. Wallin, R.F. and E.F. Arcscott, 1998. A practical guide to ISO 10993-5: Cytotoxicity. *Med. Dev. Diag. Ind.*, 20: 96-98.
91. Sagdic, O., L. Ekici, I. Ozturk, T. Tekinay and B. Polat *et al.*, 2013. Cytotoxic and bioactive properties of different color tulip flowers and degradation kinetic of tulip flower anthocyanins. *Food Chem. Toxicol.*, 58: 432-439.
92. López-García, J., Z. Kuceková, P. Humpolíček, J. Mlček and P. Saha, 2013. Polyphenolic extracts of edible flowers incorporated onto atelocollagen matrices and their effect on cell viability. *Molecules*, 18: 13435-13445.
93. Cichewicz, R.H., Y. Zhang, N.P. Seeram and M.G. Nair, 2004. Inhibition of human tumor cell proliferation by novel anthraquinones from daylilies. *Life Sci.*, 74: 1791-1799.
94. Lin, Y.L., C.K. Lu, Y.J. Huang and H.J. Chen, 2011. Antioxidative caffeoylquinic acids and flavonoids from *Hemerocallis fulva* flowers. *J. Agric. Food Chem.*, 59: 8789-8795.
95. Yang, P.F., Y.N. Yang, C.Y. He, Z.F. Chen and Q.S. Yuan *et al.*, 2019. New caffeoylquinic acid derivatives and flavanone glycoside from the flowers of *Chrysanthemum morifolium* and their bioactivities. *Molecules*, Vol. 24. 10.3390/molecules24050850.
96. Kwon, J.H., H.J. Oh, D.S. Lee, S.J. In and K.H. Seo *et al.*, 2019. Pharmacological activity and quantitative analysis of flavonoids isolated from the flowers of *Begonia semperflorens* Link et Otto. *Appl. Biol. Chem.*, Vol. 62. 10.1186/s13765-019-0416-6.
97. Offei-Oknye, R., J.L. Patterson, L.T. Walker, J. Boateng and M. Verghese, 2015. Chemopreventive potential of ginger on Hep2G cells. *Int. J. Cancer Res.*, 11: 52-66.
98. Cheraghi, O., G. Dehghan, M. Mahdavi, R. Rahbarghazi and A. Rezaabakhsh *et al.*, 2016. Potent anti-angiogenic and cytotoxic effect of conferone on human colorectal adenocarcinoma HT-29 cells. *Phytomedicine*, 23: 398-405.
99. Goutzourelas, N., D. Stagos, A. Housmekeridou, C. Karapouliou and E. Kerasioti, 2015. Grape pomace extract exerts antioxidant effects through an increase in GCS levels and GST activity in muscle and endothelial cells. *Int. J. Mol. Med.*, 36: 433-441.
100. Lima, C.F., M. Fernandes-Ferreira and C. Pereira-Wilson, 2006. Phenolic compounds protect HepG2 cells from oxidative damage: Relevance of glutathione levels. *Life Sci.*, 79: 2056-2068.
101. Singh, S., D.K. Singh, A. Meena, V. Dubey, N. Masood and S. Luqman, 2019. Rutin protects t-butyl hydroperoxide-induced oxidative impairment via modulating the Nrf2 and iNOS activity. *Phytomedicine*, 55: 92-104.

102. Crespo, R., B.E. Rodenak-Kladniew, M.A. Castro, M.V. Soberón and S.M.L. Lavarías, 2020. Induction of oxidative stress as a possible mechanism by which geraniol affects the proliferation of human A549 and HepG2 tumor cells. *Chem. Biol. Interact.*, Vol. 320. 10.1016/j.cbi.2020.109029.
103. Franco, R. and J.A. Cidlowski, 2009. Apoptosis and glutathione: Beyond an antioxidant. *Cell Death Differ.*, 16: 1303-1314.
104. Anderson, M.E., 1998. Glutathione: An overview of biosynthesis and modulation. *Chem. Biol. Interac.*, 111-112: 1-14.
105. Du, Z.X., H.Y. Zhang, X. Meng, Y. Guan and H.Q. Wang, 2009. Role of oxidative stress and intracellular glutathione in the sensitivity to apoptosis induced by proteasome inhibitor in thyroid cancer cells. *BMC Cancer*, Vol. 9. 10.1186/1471-2407-9-56.
106. Kasala, E.R., L.N. Bodduluru, R.M. Madana, K.V. Athira, R. Gogoi and C.C. Barua, 2015. Chemopreventive and therapeutic potential of chrysin in cancer: Mechanistic perspectives. *Toxicol. Lett.*, 233: 214-225.
107. Ortega, A.L., S. Mena and J.M. Estrela, 2011. Glutathione in cancer cell death. *Cancers*, 3: 1285-1310.