



# Asian Journal of Crop Science

ISSN 1994-7879

**science**  
alert  
<http://www.scialert.net>

**ANSI***net*  
an open access publisher  
<http://ansinet.com>



## Research Article

# Fastest Base Population Formation of Maize Through Exploitation of Xenia and Metaxenia Phenomena Derived from Hybrid Maize Varieties as Parents

<sup>1</sup>Suaib Suaib and <sup>2</sup>Darwis Suleman

<sup>1</sup>Department of Plant Breeding, Faculty of Agriculture, Halu Oleo University, Kendari, Southeast Sulawesi, Indonesia

<sup>2</sup>Department of Soil Sciences, Faculty of Agriculture, Halu Oleo University, Kendari, Southeast Sulawesi, Indonesia

## Abstract

**Background and Objectives:** Higher production of maize (*Zea mays* L.) could be obtained by breeding in a much faster method by using the xenia and metaxenia phenomena as the main observations. The objectives of this research were to study the possibility of maize hybrid varieties seeds as crossing parents in exploiting both xenia and metaxenia phenomena in creating the base population materials for breeding the new promising candidate of higher production variety of maize. **Materials and Methods:** Three seeds from hybrid varieties as crossing parents were Bonanza and Ganebo varieties (sweet maize) and Arumba variety (waxy maize). These varieties were separated and sown into the 500 cm long and 300 cm wide plot, 75 cm between rows and 30 cm within rows, with one seed per hill, so that there would be 56 individual plant per plot. The authors employed diallel crossing method and obtained three of each of regular  $F_1$  kernels, reciprocal  $F_1$  kernels and open pollinated  $F_1$  kernels of female parents as data sources.  $F_1$ 's kernels colour, 100 kernels weights (g) and the number of kernels per ear were the xenia components and diameter (mm) and lengths of ears (cm) were the metaxenia components. One way analysis of variance according to completely randomized design was used to analyze the observed variables and the Tukey t-test was subjected in judging the difference between  $F_1$ 's. **Results:** The research results showed that the 100 kernels weight, the number of kernels per ear and the diameter and the lengths of ears of some  $F_1$ 's as xenia and metaxenia phenomena, respectively, resulted better than their maternal parents under open pollination. **Conclusion:** There was a possibility to use these both hybrid varieties and xenia and metaxenia phenomena in producing the seeds materials in a faster way as base population for the next selection step in creating a promising new higher production of maize variety.

**Key words:** Diallel cross, higher production, maize hybrid variety, metaxenia, xenia

**Citation:** Suaib Suaib and Darwis Suleman, 2018. Fastest base population formation of maize through exploitation of xenia and metaxenia phenomena derived from hybrid maize varieties as parents. Asian J. Crop Sci., 10: 207-213.

**Corresponding Author:** Suaib Suaib, Department of Plant Breeding, Faculty of Agriculture, Halu Oleo University, Kendari, Southeast Sulawesi, Indonesia Tel: 081392227644

**Copyright:** © 2018 Suaib Suaib and Darwis Suleman. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

**Competing Interest:** The authors have declared that no competing interest exists.

**Data Availability:** All relevant data are within the paper and its supporting information files.

## INTRODUCTION

The term “Xenia” has already been referred to many species in 1881 by Focke<sup>1</sup> and “metaxenia” in corn by Kellerman and Swingle<sup>2</sup> and Denney<sup>3</sup>. These terms were two of the most widely discussed topics among the scientists during the period of 1881-1900's. Both phenomena were defined in disparities by some scientists in that the understanding in this aspect remain unclear, conflicted and overlapping from one to another. The term of “xenia” refers to the immediate effect of pollen on endosperm as a result of double fertilization (union of sperm with egg and two polar nuclei) in seed parent, while “metaxenia” refers to the direct effect of pollen on parts of seed and fruit outside embryo sac<sup>4</sup>. According to Denney<sup>3</sup>, the term of xenia was defined as the phenomenon of direct differential effects produced by pollen from different sources on the discernible characteristics, especially of size, colour, shape, chemical composition and developmental timing of part of seeds and other fruits, including the embryo, endosperm and maternal tissues, in the period from fertilization until germination. Whereas, the term of metaxenia was defined as the effect of pollen on fruit shape and other characteristics<sup>5</sup>.

From the explanations above the xenia phenomenon could be simplified as some characteristics that related to the seeds and the metaxenia may be addressed as some fruit characteristics, so that the kernel and ear characteristics of maize plant are strongly related to the xenia and metaxenia, respectively. Some attributes of kernel and ear characteristics are determining the production of maize other than fibre production. The heaviest the kernel and the longest the ear of some maize cultivars, the higher production would be obtained. Another maize kernel and ear characteristics are also determining the maize production, for example, the number of kernels per ear and the diameter of ear. Chemical composition of maize kernel would also be a xenia phenomenon like.

Following Kellerman dan Swingle<sup>2</sup>, observation about xenia and metaxenia in maize plant had been done in much more intensive, either in frequencies or in researcher team. Some of these were Kiesselbach and others<sup>1,5-13</sup>. As a result, an existing in plant breeding application in form of “top-cross” procedure has been applied in this plant species<sup>14-16</sup>. Xenia and metaxenia studies in another plant species were date palm by Swingle and others<sup>17-19</sup>, *Vicia faba* (L.) by Duc *et al.*<sup>20</sup>, *Mandarin* by Weingartner *et al.*<sup>21-23</sup>, *Hylocereus polyrhizus* by Mizrahi *et al.*<sup>24</sup>, *Selenicereus meianthus* by Dag and Mizrahi<sup>25</sup>, *Gossypium hirsutum* by Pahlavani and Abolhasani<sup>26</sup> and tomato plant (*Solanum lycopersicum* L.) by Piotto *et al.*<sup>27</sup>.

Possessing the higher production of maize, more precise and faster method of its breeding have already been proposed and implemented. Forming inbred line through selfing method for several generations, generally 7-8 generations selfing was the usual way<sup>28</sup>. The higher production of crossing pairing parents was also the most parents candidate pre-requisite<sup>29</sup>. Meanwhile, the plant hybrid vigour or heterosis achieved in their filials must be in higher level even though this was not the exact event. It could be in the mid parent heterosis or in the high parent heterosis or in the heterobeltiosis grades<sup>30,31</sup>. To this days, no reports had successfully been published on a more precise and faster way derived from hybrid variety as crossing pairing parents.

Exploration of both xenia and metaxenia phenomena in forming the base population of a candidate of superior maize variety as selection material was rarely published, except the scientific studies under genetical aspect of some crop plants. One of the usefulness result from similar study was the existence and the application of “top cross” method, while the materials used was from the inbred line as the crossing pairing parents<sup>14-16</sup>. Even though this was a preliminary study in using a hybrid varieties as crossing parents, the speculation of their fruitfulness should be proved. Faster method to determine the promising basic population for selection materials was the primary reason in observing the xenia and metaxenia by crossing of parentals pairing as hybrid variety. From this aspect, the colour of kernel, 100 kernel weight (g) and the number of kernels as xenia component were chosen, whereas the diameter (mm) and the shoot length (cm) as metaxenia element were picked-out. All variables mentioned above could be considered in determining the maize production level in as early as of the time.

This article will reported and deeply discussed in relations to the application of both xenia and metaxenia phenomena in creating a new promising superior variety of maize in a short period of time. Exploitation of both phenomena as a way to form of basic population for material selection further was also discussed.

## MATERIALS AND METHODS

**Location and plant materials:** The field experiment was conducted at Faculty Agriculture Experimental Station II, Halu Oleo Campus, Anduonohu Kendari, Southeast Sulawesi Indonesia, from March-July, 2018. A set of tools for land preparation, plant management and harvesting and variable measurements were used, while the seeds as a source of parental crossings of the three hybrid varieties, namely Bonanza, Ganebo and Arumba were used. The two first

varieties were sweet maize and dent type kernels and the latter was waxy maize type with flint type kernels. Days to harvest of these varieties were varied according to the variety but they were 60-70 days based on variety identity.

**Field experimental procedure and design:** Parental crossing sources were separately sown each in the 500 cm length and 300 cm wide plot and 200 cm between plots. The length of 75 cm between rows and 30 cm within row were applied as plants space and one seed per hill, so that there were seven rows for the length side and eight holes for the wide side and a total of 56 plants per plot. To avoid natural crossing pollination with different varieties the planting time arrangement was applied as an isolation method and each variety was sown two times, four rows at the first time and the remaining four rows after 2 weeks. The plants were managed according to the standard of maize hybrid plantation. Crossing parent pairings was done by artificial hands pollination to get both regular and reciprocal kernel and ear of  $F_1$  and by natural open pollination within the variety to get the  $F_1$  kernel and ear of female parent as comparison data. An artificial hands crossing was conducted when the shoot of female parent were silking and bagged by shootbag and the pollen source of male parent was bagged by tasselbag as soon before anthesis. The diallel method of Griffing<sup>32</sup> was used in parents crossing pattern.

**Research variables:** Those kernels and ears were harvested when the shoots achieved the physiological mature state in which the shootleaf was browning in colour. The kernel colour, 100 kernel weight (g) and the number of kernel per ear were xenia variables, whereas the ear diameter (mm) and ear length (cm) were metaxenia variables. Three times of each variable were subjected as replication.

**Data analysis:** The colour of kernels as a qualitative measurements was analyzed by qualitative method then qualitatively described and the quantitative measurement was analyzed by one way analysis of variance based on a completely randomized design (CRD). Mean separation between  $F_1$ 's was judged through LSD (Least Significant Difference) test of Tukey<sup>33</sup> at alpha 0.05 of significant level. All data analysis procedures was conducted by statistical package of SAS for Windows version 6.13<sup>34</sup>.

## RESULTS

By using complete diallel crossing scheme of the three parents, those three  $F_1$  kernels and ears, three  $F_1$  reciprocals kernels and ears and three  $F_1$  open pollinated kernels and ears

families were obtained. All these families were used as data analyzed in studying the xenia and metaxenia phenomena.

**Xenia observation:** Both Bonanza and Paragon parents were having the yellow kernel colour expression and the Arumba parent was white kernel colour. The kernel colour of qualitative character as xenia component which immediately expressed in female parent plant ( $F_1$  kernel) was observed as a result of kernel colour inheritance from male parent plant. Results of the research indicated that the yellow kernel colour of the male parent crossing with the yellow kernel colour of female parent generated the only yellow of kernel colour of  $F_1$ . Meanwhile, when the male and female parents had different kernel colour like white-yellow or yellow-white, respectively, the kernel colour of female parent plant ultimately expressed with both white and yellow colour kernels (Table 1).

The 100 kernels weight (g) and the number of kernels per ear of  $F_1$ 's resulted from crossing between parents were generally expressed the similar ear performances of female parent obtained to open pollination. One way analysis of variance of the effect of parental pollen on 100 kernel weight and number of kernels per ear of male parents showed the highly significance of both xenia component characters. Mean squared between kernels of male and female parents and their  $F_1$ 's 100 kernels weight and the number of kernel per ear were significantly differed. Ganebo and Arumba varieties as male parents and Bonanza variety as female parent were significantly differed from 100 kernel weight (Table 2).

Table 1: Male parent effect on  $F_1$  kernel colour of female parent

Parents*		$F_1$ kernel colour	
Female	Male	Yellow	White
Bonanza	Bonanza	+	-
Bonanza	Ganebo	+	-
Bonanza	Arumba	+	+
Ganebo	Ganebo	+	-
Ganebo	Bonanza	+	-
Ganebo	Arumba	+	+
Arumba	Arumba	-	+
Arumba	Bonanza	+	+
Arumba	Ganebo	+	+

\*Note of kernel colour, Bonanza: Yellow, Ganebo: Yellow and Arumba: White  
-: Not exist, +: Exist

Table 2: Variance analysis of the effect of male parent on the 100 kernel weight and the number of kernel per ear of Bonanza, Ganebo and Arumba female parents

Source of variation	Mean square	
	100 kernel weights (g)	No. of kernel per ear
Female Bonanza	3.866*	2067.111*
Female Ganebo	14.109*	1002.778*
Female Arumba	1.315*	303.444*

\*: Statistically significant differences at the 0.05 probability level

Table 3: Effect of male parents on the 100 kernel weight (g) and the number of kernels per ear of Bonanza, Ganebo and Arumba female parents

Parents			
Female	Male	100 kernel weight (g)	No. of kernel per ear
Bonanza	Bonanza	11.580 <sup>a</sup>	260.000 <sup>b</sup>
Bonanza	Ganebo	13.050 <sup>b</sup>	264.000 <sup>b</sup>
Bonanza	Arumba	13.813 <sup>b</sup>	216.667 <sup>a</sup>
Ganebo	Ganebo	16.203 <sup>i</sup>	241.667 <sup>k</sup>
Ganebo	Bonanza	11.880 <sup>m</sup>	250.000 <sup>k</sup>
Ganebo	Arumba	13.813 <sup>k</sup>	215.000 <sup>l</sup>
Arumba	Arumba	16.413 <sup>x</sup>	212.667 <sup>y</sup>
Arumba	Bonanza	16.223 <sup>x</sup>	231.000 <sup>x</sup>
Arumba	Ganebo	15.183 <sup>y</sup>	229.000 <sup>x</sup>

Different letters in a column within the same female indicate the significant differences at alpha 0,05 level according to the LSD-test

Mean separation of 100 kernel weight of  $F_1$ 's between parents pairing and their open pollination of female parents was significantly differed only the crossing between Bonanza as female parent and Ganebo and Arumba varieties as male parents. While, crossing pairings on both Ganebo and Arumba varieties as female parents were not significantly differed in 100 kernels weight. Another crossing pairings between Bonanza and Ganebo varieties as male parents and Arumba variety as female parent was also the only pairings that indicated the significantly differed in kernel number per ear. Whereas, crossing pairings on Bonanza and Ganebo varieties as female parent, statistically was not significantly differed in producing kernels number of  $F_1$ 's compared to their open pollinated of female parents (Table 3).

**Metaxenia observation:** One way analysis of variance on both ear diameter (mm) and length (cm) were significantly differed in all crossing pairings (Table 4). Comparison analysis in ear diameter and length between the open pollinated male parents and their  $F_1$ 's resulted from crossing parents were significantly differed at alpha 0.05 level according to Tukey's test. Crossing pairings on Bonanza variety as female parent and Ganebo variety as male parent and between Arumba variety as female parent and Bonanza and Arumba varieties as male parent were significantly differed on ear diameter. Ganebo variety as female parent was not affected by Bonanza and Arumba varieties as pollen parents in expressing the ear diameter. Crossing on Ganebo variety as female parents did not produce higher values of ear diameter and significantly differed when crossed with Bonanza and Arumba varieties as male parents, respectively. Another crossing pairing between Ganebo variety as female parent and Bonanza variety as male parent and between Arumba variety as female parent and Ganebo variety as male parent were significantly affected

Table 4: Variance analysis of the effect of male parent on ear diameter (mm) and ear length (cm) of Bonanza, Ganebo and Arumba female parents

Source of variation	Mean square	
	Ear diameter (mm)	Ear length (cm)
Female Bonanza	7239.000*	17.898*
Female Ganebo	10143.444*	6.564*
Female Arumba	7250.333*	19.223*

\*:Statistically significant differences at the 0.05 probability level

by their ear length compared to their open pollinated female parents. Whereas, crossing to the Bonanza variety as female parent was not significantly differed in yielding the ear length of their  $F_1$ 's compared to the ear length of open pollinated female (Table 5).

## DISCUSSION

Basically, there were four layers from outer to inner layer that composed the maize kernel, i.e., (1) Pericarp or outer layer, (2) Aleuron layer, (3) Endosperm layer and (4) Embryo. The kernel colour expression was due to the aleuron layer pigmentation, while the non-colour kernel expression was resulted from the lack of pigments produced by aleuron. Commonly, there are four genes that are responsible for the kernel colour expression also called colour genes. Symbolically, they are *Pr1* gene that expressed the aleuron red colour, *Y1* gene for white colour, *C1* gene for coloured aleuron1 and *R1* gene for coloured1. These two genes are similar but their name and symbols are different and encoded transcription activators that interact to influence the regulation of several structural genes within the anthocyanin pathway<sup>35</sup>.

In this field experiment, the kernels colour expression indicated the existency of xenia phenomenon on pattern as listed in Table 1, particularly in crossing between yellow and white kernels in which the  $F_1$  ear or kernel of female parents

Table 5: Effect of male parents on ear diameter (mm) and ear length of Bonanza-F<sub>1</sub>, Ganebo and Arumba female parents

Parents			
Female	Male	Ear diameter (mm)	Ear length (cm)
Bonanza	Bonanza	517.000 <sup>a</sup>	21.500 <sup>a</sup>
Bonanza	Ganebo	560.000 <sup>b</sup>	18.700 <sup>b</sup>
Bonanza	Arumba	462.000 <sup>c</sup>	16.633 <sup>c</sup>
Ganebo	Ganebo	560.000 <sup>k</sup>	18.700 <sup>l</sup>
Ganebo	Bonanza	536.667 <sup>l</sup>	20.500 <sup>k</sup>
Ganebo	Arumba	449.667 <sup>m</sup>	17.567 <sup>m</sup>
Arumba	Arumba	454.000 <sup>z</sup>	17.600 <sup>y</sup>
Arumba	Bonanza	524.333 <sup>y</sup>	16.967 <sup>y</sup>
Arumba	Ganebo	548.667 <sup>x</sup>	21.633 <sup>x</sup>

Different letters in a column within the same female indicate the significant differences at alpha 0,05 level according to the LSD-test

produced the yellow and white colours in a proportionally comparable comparison. Unfortunately, the crossing between those parents that had the same colour kernels will be resulting an F<sub>1</sub> which had the same kernels colour with their male and female parents which are naturally open pollinated. This findings was in line with Vancetovic *et al.*<sup>16</sup> observation on top cross scheme of five commercial hybrid varieties (ZP-1, ZP-2, ZP-3, ZP-4 and ZP-5) with yellow colours as female parents and one open pollination variety with dark blue kernels colour as male parent. Vancetovic *et al.*<sup>16</sup> reported that all the F<sub>1</sub> ears (female ears, other than open pollinated) were similar to the male ears colour namely the dark blue colour. Another identical observation reported that the crossing between IHO variety with low carotene content and yellow colour kernels as female parent and PR variety with high anthocyanin content and dark purple colour kernels would result in the F<sub>1</sub> kernels with which is similar to the female kernel colour i.e., dark purple colour kernels<sup>13</sup>.

Kernel colour expression in F<sub>1</sub> female parent was the indication of xenia existency as a result of male parent pollen effect. By this pattern, another xenia elements like carbohydrates, proteins, lipids and other derivatives kernel contents, 100 kernel weight and number of kernels per ear, might easily be expected. Sprague<sup>36</sup> reported that the kernel colour inheritance pattern of maize showed that the red, yellow and purple colours were the result of four pairing factors, namely the respective symbols S<sub>1</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>2</sub>, S<sub>3</sub>, S<sub>3</sub>, S<sub>4</sub> and S<sub>4</sub>. Later, Ford<sup>35</sup> explained that the fourth colours of maize kernel above were determined as alleles C<sub>1</sub>, C<sub>1-1</sub>, c<sub>1</sub>, Pr<sub>1</sub>, pr<sub>1</sub>, R<sub>1</sub>, r<sub>1</sub> and Y<sub>1</sub> coded by red, yellow and purple colours, respectively, whereas the y<sub>1</sub> was the white colour kernel. The expressions of each maize kernel colour in F<sub>1</sub> ear female was combination between their crossing pairing parents and the alleles number dependent<sup>35,36</sup>.

The weight of 100 kernels and the number of kernels per ear as another xenia elements generally indicated the decleaning on both variables of F<sub>1</sub> female parent resulting

from crossing between parents compared to the open pollination female parent. It means that the female parents plays an important role in the increasing the weight of 100 kernels and the number of kernel per ears. Only some crossing pairings which could be increased in these variables so that the compatibility of xenia phenomenon of both parents or the expression of xenia phenomenon in this observation was female parents dependent. From all pairing of parental crossings, there were two pairs of crossing which pointed the increasing of 100 kernels weight that are crossing between Bonanza female parent and Ganebo and Arumba varieties as male parents (Table 4). Meanwhile, the increase of kernels number per ear was expressed in the crossing between Arumba variety as female parent and Bonanza and Ganebo varieties as male parents. This expression showing the existence of paternal pollen effect of male plant in partial and the absence of maternal effect of female plant in remaining.

The important finding in this observation have shown that the parallele reports of another researchers<sup>12,13,16,37-39</sup> in which either the 100 kernels weight or the kernels number per ear were reported the same pattern. This means that the inheritance of the 100 kernel weight and the kernel number per ear depended upon the heritability traits of crossing pairings between male and female parents. In this connection, the inheritance of some xenia element is strongly depended by pollen parent effect or male parent so called paternal effect. In contrast, the unexpressed xenia phenomenon may caused by the dominance female effect or maternal effect. So, the increasing of yield was indicated by the higher production of F<sub>1</sub> female kernels derived from crossing between parents compared with the F<sub>1</sub> female kernels by open pollination.

The ear diameter and length were metaxenia components that are closely related to the maize variety production. Ear length and diameter are two of some characters that are tightly associated in maize production and both of these characters are metaxenia elements. The bigger the ear diameter and the longer the ear length, the most



probability of reaching the higher production of a maize variety as a result of the higher number of kernel per ear. As figured out at xenia phenomenon which showed only some crossing pairs that heritable, the metaxenia phenomenon in this findings was followed the above pattern. These could be seen in the  $F_1$  ear length and diameter obtained at the female parents in crossing pairing between male and female parents compared to the  $F_1$  open pollinated female parents (Table 5). Observation results from this field experiment showed the similar indication to the xenia effect that had been reported by several researchers from different regions<sup>12,13,16,37</sup>.

Both xenia and metaxenia in hesitancies are not always followed the increasing value of some characters but the decreasing value of some measures would be existed depending on the male parent potency of that character. In the case of the rising yield of some crop plant variety, the higher potential yield of male parent will be needed but in the case of lowering yield such as decreasing unsaturated lipid content of kernels, may substantially be needed the lower content of it component to the male parent. Then, the  $F_1$  kernels content obtained of female parent by crossing the male parent may be lower than the  $F_1$  kernels of female parent resulted from natural open pollination. Thus, the formation of the higher production of new superior maize variety through xenia and metaxenia exploitation, the higher production of male parent must be provided the higher values of 100 kernel weight, kernel number per ear and ear length and diameter.

### CONCLUSION

The kernel colour as xenia element was fully inherited at crossing pairings between white and yellow colour kernels. Another xenia elements i.e., 100 kernels weight was expressed when Ganebo and Arumba varieties were as male parents and the kernels number per ear was indicated when Bonanza and Ganebo varieties were as male parents. The ear diameter as metaxenia element was observed when Ganebo and Bonanza varieties as male parents. Whilst, the ear lengths as another metaxenia element was pointed out when Bonanza and Ganebo varieties were as male parents.

### SIGNIFICANCE STATEMENT

This study reports results of a study on xenia and metaxenia on kernel and ear characteristics as production variables derived from hybrid maize varieties as parents. Results generated here will eventually help maize breeders use the xenia and metaxenia without lengthy procedure in

forming the base population of maize as well as producing the new promising higher production of maize variety. The results presented will also give indication of appropriate use of both phenomena in generating the new superior varieties of maize.

### ACKNOWLEDGMENT

We would like to extend our gratitude to the Ministry of Research, Technology and Higher Education of Republic of Indonesia for the financial support provided for this research project Contract No. 056/SP2H/LT/DRPM/2018. Also to Miss Nur Rizky Alfiany Suaib, English Language and Applied Linguistics, Faculty of Arts and Humanity, Coventry University, Priory Street, CV1 5FB, Coventry, United Kingdom, who provided proofreading for the English expressions.

### REFERENCES

1. Focke, W.O., 1881. Die Pflanzen-Mischlinge: Ein Beitrag Zur Biologie Der Gewachse. Bomtraeger, Berlin, pp: 510-518, (Abstract in English).
2. Kellerman, W.A. and W.T. Swingle, 1888. Experiments in cross-fertilization of corn. Annu. Rep. Kansas Exp. Stat., 1: 316-337.
3. Denney, J.O., 1992. Xenia includes metaxenia. HortScience, 27: 722-728.
4. Soule, J., 1985. Glossary for Horticultural Crops. Wiley, New York.
5. Olfati, J.A., Z. Sheykhtaher, R. Qamgosar, A. Khasmakhi-Sabet, G. Peyvast, H. Samizadeh and B. Rabiei, 2009. Xenia and metaxenia on cucumber fruit and seed characteristics. Int. J. Vegetable Sci., 16: 243-252.
6. Kiesselbach, T.A., 1960. The significance of xenia effects on the kernel weight of corn. Research Bulletin: Bulletin of the Agricultural Experiment Station of Nebraska, No. 191.
7. Pixley, K.V. and M.S. Bjarnason, 1994. Pollen-parent effects on protein quality and endosperm modification of quality protein maize. Crop Sci., 34: 404-409.
8. Letchworth, M.B. and R.J. Lambert, 1998. Pollen parent effects on oil, protein and starch concentration in maize kernels. Crop Sci., 38: 363-367.
9. Bulant, C., A. Gallais, E. Matthys-Rochon and J.L. Priul, 2000. Xenia effects in maize with normal endosperm: II. Kernel growth and enzyme activities during grain filling. Crop Sci., 40: 182-189.
10. Dong, H., 2007. Effect of high oil corn pollinator on kernel quality of common corn and their physiological and biochemical basis. Master's Thesis, Shandong Agricultural University, China.
11. Ahuja, M. and N.S. Malhi, 2008. Genetic analysis of xenia effects in high oil maize lines. Indian J. Genet. Plant Breed., 68: 391-397.

12. Kahrman, F., C.O. Egesel and E. Sorlu, 2015. Effects of open- and self-pollination treatments on genetic estimations in maize diallel experiment. Spanish J. Agric. Sci., 13: 1-11.
13. Kahrman, F., M. Serment, M. Haslak and M.S. Kang, 2017. Pollen effect (Xenia) for evaluating breeding materials in maize. J. Genetika, 49: 217-234.
14. Thompson, P.R., A.B. Geyer, L.D. Lotz, H.J. Siegrist and T.L. Dobbels, 2002. Top cross high-oil corn production: Agronomic performance. Agron. J., 94: 290-299.
15. Thomison, P.R., A.B. Geyer, L.D. Lotz, H.J. Siegrist and T.L. Dobbels, 2003. Top cross high oil corn production: Select grain quality attributes. Agron. J., 95: 147-154.
16. Vancetovic, J., S. Zilic, S. Bozinovic and D. Ignjatovic-Micic, 2014. Simulating of top-cross system for enhancement of antioxidants in maize grain. Spanish J. Agric. Res., 2: 467-476.
17. Swingle, W.T., 1928. Metaxenia in the date palm: Possibly a hormone action by the embryo or endosperm. J. Hered., 19: 257-268.
18. Nixon, R.W., 1928. The direct effect of pollen on the fruit on the date palm. J. Agric. Res., 36: 97-128.
19. Nixon, R.W., 1928. Immediate influence of pollen: In determining the size and time of ripening of the fruit of the date palm. J. Heredity, 19: 241-255.
20. Duc, G., A. Moessner, F. Moussy and C. Mousset-Declas, 2001. A xenia effect on number and volume of cotyledon cells and on seed weight in faba bean (*Vicia faba* L.). Euphytica, 117: 169-174.
21. Weingartner, U., O. Kaeser, M. Long and P. Stamp, 2002. Combining cytoplasmic male sterility and xenia increases grain yield of maize hybrids. Crop Sci., 42: 1848-1856.
22. Weingartner, U., T.J. Prest, K.H. Camp and P. Stamp, 2002. The plus-hybrid system: A method to increase grain yield by combined cytoplasmic male sterility and xenia. Maydica, 47: 127-134.
23. Weingartner, U., K.H. Camp and P. Stamp, 2004. Impact of male sterility and xenia on grain quality traits of maize. Eur. J. Agron., 21: 239-247.
24. Mizrahi, Y., J. Mouyal, A. Nerd and Y. Sitrit, 2004. Metaxenia in the Vine Cacti *Hylocereus polyrhizus* and *Selenicereus* spp. Ann. Bot., 93: 469-472.
25. Dag, A. and Y. Mizrahi, 2005. Effect of pollination method on fruit set and fruit characteristics in the vine cactus *Selenicereus megalanthus* ("yellow pitaya"). J. Hortic. Sci. Biotechnol., 80: 618-622.
26. Pahlavani, M.H. and K. Abolhasani, 2006. Xenia effect on seed and embryo size in cotton (*Gossypium hirsutum* L.). J. Applied Genet., 47: 331-335.
27. Piotto, F.A., K.D. Batagin-Piotto, M.D. Almeida and G.C.X. Oliveira, 2013. Interspecific xenia and metaxenia in seeds and fruits of tomato. Scient. Agricola, 70: 102-107.
28. Allard, R.W., 1960. Principles of Plant Breeding. 1st Edn., John Wiley and Sons Inc., New York.
29. Sleper, D.A. and J.M. Poehlman, 2006. Breeding Field Crops. 5th Edn., Wiley-Blackwell, New York, USA., ISBN-13: 9780813824284, Pages: 424.
30. Fehr, W.R., 1993. Principles of Cultivar Development: Theory and Technique. Vol. 1. MacMillan, New York.
31. Ghahal, G.S. and S.S. Gosal, 2002. Principles and Procedures of Plant Breeding: Biotech-Nological and Conventional Approaches. Alpha Science International Ltd., Pangbourne, UK.
32. Griffing, B., 1956. Concept of general and specific combining ability in relation to diallel crossing systems. Aust. J. Biol. Sci., 9: 463-493.
33. Steel, R.R.D. and J.H. Torrie, 1980. Principles and Procedures of Statistics. 3rd Edn., McGraw-Hill International Book Co., London, Pages: 633.
34. SAS., 2002. SAS for windows version 6.13. SAS Institute Inc., Cary, NC, USA.
35. Ford, R.H., 2000. Inheritance of kernel color in corn: Explanations & investigations. Am. Biol. Teach., 62: 181-188.
36. Sprague, G.F., 1932. The inheritance of colored scutellums in maize. Techn. Bull., 292: 1-44.
37. Weiland, R.T., 1992. Cross-pollination effects on maize (*Zea mays* L.) hybrid yields. Can. J. Plant Sci., 72: 27-33.
38. Alvarez-Prado, S., C.G. Lopez, M.L. Senior and L. Borrás, 2014. The genetic architecture of maize (*Zea mays* L.) kernel weight determination. G3: Genes Genet. Genomes, 4: 1611-1621.
39. Liu, Z., A. Garcia, M.D. McMullen and S.A. Flint-Garcia, 2016. Genetic analysis of kernel traits in maize-teosinte introgression populations. G3: Genes Genomes Genet., 6: 2523-2530.