

# Asian Journal of Crop Science





### ට OPEN ACCESS

#### **Asian Journal of Crop Science**

ISSN 1994-7879 DOI: 10.3923/ajcs.2019.40.45



# Research Article Effect of Ratooning on Growth and Nutritional Quality of *Amaranthus* (*Amaranthus tricolor*) in Alfisol South Western Nigeria

<sup>1</sup>Ayeni Leye Samuel, <sup>2</sup>Oyebamiji, Kehinde Johnson, <sup>2</sup>Morakinyo-Fasipe Olutoye Temitope and <sup>3</sup>Agbona Ademola Isaac

<sup>1</sup>Department of Agricultural Science, Adeyemi College of Education, Ondo, Nigeria <sup>2</sup>Department of Crop Science, College of Agricultural Science, Ayo Babalola University, Arakeji, Ikeji, Osun State, Nigeria <sup>3</sup>Department of Agricultural Technology, Federal Polytechnic, Ile Oluji, Ondo State, Nigeria

## Abstract

**Background and Objective:** Increase in human population that results in food scarcity and malnutrition necessitates the need to increase growing of vegetable that is early maturing with well ratooning capacity. This study aimed to examine the effect of ratooning on growth, yield and nutrient uptake of *Amaranthus tricolor*. **Materials and Methods:** Two field experiments were conducted in 2017 at 2 different locations in Okegun situated at Ondo and Adeyemi College of Education Ondo, Teaching and Research Farm Ondo, South Western Nigeria. The treatments were harvesting by uprooting the plant or total harvesting of the plant at once (1 time harvest), twice (2 times harvest) and thrice (3 times harvest). The treatments were generated and used to discuss the findings. **Results:** Relative to total harvest, 2 and 3 times harvests significantly increased (p<0.05) plant height, fresh weight, dry weight, number of leaves, fresh weight, dry weight and moisture content. Three times harvests recorded highest K. Two times harvest had highest Ca, Mg, Na, N and P. One time harvest significantly (p<0.05) recorded the highest Mn, Fe, Cu and Zn. Relative to 1 time harvest, 2 times harvest recorded highest increase in crude protein and fat content of *Amaranthus tricolor*. **Conclusion:** This study reported that the 3 times harvest recorded the highest growth parameters while 2 times harvest recorded the highest nutritional value of Amaranth in this experiment.

Key word: Agronomic parameters, amaranth, ratooning, crude protein, malnutrition, uprooting

Citation: Ayeni Leye Samuel, Oyebamiji, Kehinde Johnson, Morakinyo-Fasipe Olutoye Temitope and Agbona Ademola Isaac, 2019. Effect of ratooning on growth and nutritional quality of *Amaranthus (Amaranthus tricolor*) in Alfisol south western Nigeria. Asian J. Crop Sci., 11: 40-45.

Corresponding Author: Ayeni Leye Samuel, Department of Agricultural Science, Adeyemi College of Education, P.M.B. 520, Ondo, Ondo State, Nigeria

Copyright: © 2019 Ayeni Leye Samuel *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

#### INTRODUCTION

Vegetables are succulent crops that have the tendency of re-growth if harvested when they are still at their tender stage of development. According to Prasanna and Pattar<sup>1</sup> ratooning is common with crops like banana, red gram, cotton, Jowar, sugarcane but it has also been extended to vegetables like Amaranths. The ultimate aim of a farmer is to obtain optimum yield. African diets among the rural dwellers are mainly carbohydrates; hence, they need supplements<sup>2</sup>. Chadha et al.<sup>3</sup> observed that African leafy vegetables could make a positive contribution to world food security. The optimum yield of crops depends on the fertility of the soil, pests and diseases, cultural practices, season of planting, time and method of harvesting of such crop. Many farmers majorly focus on how to improve soil fertility and control pests and diseases with little emphasis on harvesting technique. Many research works have shown that harvesting has significant effect on growth, nutritional quality and yield of crops. Foidl et al.4 conducted research on harvest frequency on the growth and leaf yield of moringa (Moringa oleifera lam), a leafy vegetable crop and obtain positive result. Ayub et al.5 conducted an experiment on the effect of nitrogen application and harvesting intervals on forage yield and quality of pearl millet (Pennisetum americanum L.). Currently, there is a dearth of information on the growth performance, yield and nutritional quality of *A. cruentus*<sup>6</sup>. Different methods that are commonly used by the local farmers include total uprooting of the plant, cutting of the top of the plant to give room for side shoot development known as ratooning and tipping. Tipping is one of the methods used in harvesting Amaranthus and it is done by plucking off the growing points of the shoots, while cutting is done by first cutting the plants at a height not less than 15 cm, with sequential cutting of 5 cm above the previous cutting<sup>7</sup>. The increase in human population that results in food scarcity and malnutrition necessitates the need to increase growing of Amaranths that is early maturing with well ratooning capacity and good nutritive value. There is little or no known experiment that test the harvesting interval on nutritional quality of Amaranthus tricolor in Ondo, southwestern Nigeria. The objectives of this study were to compare the effect of total harvesting to partial harvesting on growth, nutrient uptake and nutritional guality of Amaranthus tricolor in Ondo south western Nigeria.

#### **MATERIALS AND METHODS**

**Study area:** The experiments were carried out at Okegun Farm in Ondo and the Teaching and Research Farm of Agricultural

Science Department, Adeyemi College of Education, Ondo between August and October, 2017. Ondo is located in the latitude 70°, 05°N and longitude 40°, 55°E and at elevation of 381.3 m above the sea level. It belongs to the tropical rain forest zone. The soil belongs to the order Alfisol (U.S.D.A.) or Luvisol<sup>8</sup>. Evidence has shown that the land had been in use for planting various crops in the past. The lands were fairly flat, slightly sloppy and contained well drained loamy soil particles with small pebbles.

**Soil sampling:** Surface soil samples (0-20 cm) were randomly collected before the conduct of the experiment at the experimental sites, bulked, air-dried and sieved through 2 mm mesh. The bulked soil samples were analyzed for OM, total N, available P, K, Ca, Mg, Fe, Cu, Zn and Mn.

Experimental layout and design: The lands were manually cleared, stumped, mapped, pegged out and made into beds of 4 m by 4 m size with 0.5 m discard area. The experiments were laid out in Randomized Complete Block Design (RCBD) and replicated 3 times. The experimental treatments were total harvest (total uprooting-1 time harvest), 1st ratoon (2 times harvest) and 2nd ratoon (3 times harvest). The total harvest means that the tagged crops were uprooted and the roots were severed from the shoot exactly 28 days after sowing, 1st ratoon means that the tagged Amaranthus tricolor were partially cut at 28 days and finally uprooted at 42 days after sowing while the 2nd ratoon means that the tagged Amaranthus tricolor were partially harvested at 28 and 42 days and finally uprooted at 56 days after sowing. Ratoon in this context means, the shoots of the selected Amaranthus tricolor were harvested manually by cutting the stem with a local sharp knife at 20 cm above ground level and the remaining shoots of the plant were allowed to develop side shoots.

**Planting of** *Amaranthus* **and cultural operation:** *Amaranthus tricolor* seeds were purchased from the local market in Ondo. Sowing of seeds was done by broadcasting. The seeds were mixed with light dried soil to enhance even distribution of the seeds. The seeds were thinned to 20 stands per plot.

About 10 t ha<sup>-1</sup> of fresh poultry manure was cured for 2 weeks and incorporated into all the plots 2 weeks before planting to improve the soil N, P and K content when the initial soil sample showed that the soils were deficient in N, P and K (Table 1). Weeding was done manually with hand-held hoe 2 weeks after planting.

**Plant parameters determination:** The parameters determined were plants height, number of leaves, number

of branches, leaf area, fresh weight, dry weight and moisture content. The N, P, K, Ca, Mg, Fe, Cu, Zn and Mn contents of the leaves were determined. Crude protein, fat, NFE, fibre and ash contents were also determined<sup>9</sup>.

Five stands of amaranth were tagged per plot for data collection. The agronomic parameters determined were plant height, number of branches, number of leaves, leaf fresh and dry weight. Plant height was measured from the ground base of each tagged plant to the top using a meter rule. Number of leaves and branches were carried out by visual counting. The leaf area was determined using graphical method. Fresh weight was recorded by weighing the freshly harvested Amaranthus tricolor with weighing balance. The freshly harvested plants were oven dried and weighed with electric balance to determine the dry weight and moisture content of Amaranthus tricolor. For total harvest, Amaranthus tricolor was uprooted and the shoots were cut from the root at 28 days after sowing. For the 1st ratoon, (1 time harvest) the agronomic parameters determined at 28 and 42 days were added together to give cumulative yield while agronomic parameters determined at 28, 42 and 58 days were also summed up to give cumulative yield for the 2nd ratoon (2 times harvest).

The harvested plants from each plot were washed and transported in a well labeled envelope to the laboratory. The harvested plants were washed with distilled water and air-dried. The air-dried samples were packed inside well labeled envelopes and oven dried, at a low temperature until constant weights were recorded. The oven dried samples were allowed to cool inside a desiccator and grounded into powder. The grounded samples were then packed inside a cellophane nylon. The leaf N, P, K, Ca, Mg, Fe, Cu, Zn and Mn were analyzed from the wet digest of the leaf samples<sup>9</sup>. The nutritional quality such as crude protein, NFE, fibre and ash content were determined using AOAC<sup>9</sup> method. The nutrients uptake and nutritional quality of Amaranthus tricolor were determined at their final stages i.e., total harvest at 28 days, 1st ratoon at 42 days (1 time harvest) and 2nd ratoon at 58 days (2 times harvest) after sowing.

**Statistical analysis:** Data collected were subjected to one way analysis of variance using the Genstat software. The data collected from the two locations were combined and their means were used as result since the two locations had similar nutrient characteristics and belonged to the same group (Alfisol). Means were separated using Least Significant Difference at p<0.05 where F-ratio was significant.

#### RESULTS

The chemical characteristics of the soil used for the conduct of the experiment were presented in Table 1. The chemical properties of the soil showed that the soil were acidic, low in OM, N, P, K, Ca and Mg. The soil were high in Fe, Cu and Mn.

**Effect of ratooning on agronomic parameters of** *Amaranthus tricolor*: The effect of harvest at once, twice and thrice on growth parameters of *Amaranthus tricolor* was shown in Table 2. Compared to one time harvest, two times harvest and three times harvest significantly increased (p<0.05) plant height, number of leaves, number of branches, fresh and dry weight, moisture content and leaf area (except 2 times harvest).

The results showed that compared with 1 time harvest (total uprooting), plant height, number of leaves, fresh weight and moisture content of Amaranth were significantly increased (p<0.05). It was observed that 3 times harvest had the highest increase in all the agronomic parameters that were determined in this experiment.

**Effect of ratooning on nutrient uptake of** *Amaranthus tricolor.* Table 3 showed the result of ratooning on nutrient uptake of *Amaranthus tricolor.* Compared to one time harvest, two times harvest significantly increased (p<0.05) Ca, Mg, K and N while one time harvest recorded highest Mn, Fe, Cu and Zn.

Relative to 1 time harvest, *Amaranthus tricolor* harvested at 2 times recorded the highest increase in crude protein and crude fat (Table 4).

Table 1: Initial mean soil chemical properties of Okegun and Research Farm, Ondo

Soil properties	Values
pН	4.60
OM (%)	0.97
N (%)	0.04
P (mg kg <sup>-1</sup> )	5.79
Exchangeable bases (Cmol kg <sup>-1</sup> )	
К	0.08
Ca	0.67
Mg	0.81
Na	0.12
Micro nutrients (mg kg <sup>-1</sup> )	
Fe	23.40
Zn	4.86
Cu	8.70
Mn	7.19

Asian J. Crop Sci., 11 (2	2): 40-45, 2019
---------------------------	-----------------

Table 2: Cumulative effect of ratooning on growth and yield of Amaranthus tricolor

	Plant h	neight				Fresh	Dry			
Treatments	(cr	n)	No. of leaves	No. of br	anches	weight/plant (g	) weight/pla	nt (g) Mois	ture content	Leaf area (cm <sup>2</sup> )
3 times harvest	48.	77ª	44.86ª	4.44ª		221.36ª	46.56	3	174.86ª	18.04ª
2 times harvest	35.2	21 <sup>b</sup>	28.44 <sup>b</sup>	4.11ª		196.83 <sup>b</sup>	23.30 <sup>i</sup>	D	173.53ª	8.26 <sup>b</sup>
1 time harvest	22.	44 <sup>c</sup>	11.99°	2.60	5 <sup>b</sup>	37.00 <sup>c</sup>	6.10	:	30.90 <sup>b</sup>	9.74 <sup>b</sup>
Lsd (0.05)	4.5	56	7.67	1.2	1	10.34	10.00		10.12	3.89
Table 3: Cumulativ	e effect of rat	ooning on	nutrient uptake	of <i>Amaranth</i>	us tricoloi	r				
Treatments	Ca (%)	Mg (%)	K (%)	Na (%)	N (%)	P (%)	Mn (mg kg <sup>-1</sup> )	Fe (mg kg <sup>-1</sup> )	Cu (mg kg <sup>-1</sup> )	Zn (mg kg <sup>-1</sup> )
3 times harvest	1.44 <sup>b</sup>	1.09 <sup>b</sup>	6.70ª	1.15ª	2.93 <sup>b</sup>	0.39ª	106.59 <sup>b</sup>	403.56 <sup>c</sup>	5.84 <sup>b</sup>	54.09°
2 times harvest	1.90ª	1.70ª	6.24ª	1.28ª	3.69ª	0.42ª	115.26 <sup>b</sup>	1283.45 <sup>b</sup>	7.65 <sup>b</sup>	66.89 <sup>b</sup>
1time harvest	1.64 <sup>b</sup>	1.07 <sup>b</sup>	3.85 <sup>b</sup>	0.83 <sup>b</sup>	2.92 <sup>b</sup>	0.39ª	230.34ª	4012.26ª	61.43ª	185.33ª
Lsd (0.05)	0.27	0.12	1.34	0.27	1.11	NS	16.78	21.12	4.56	5.64

Table 4: Cumulative effect of ratooning on nutritional quality of *Amaranthus* tricolor

Treatments	Crude protein	Crude fat	NFE	Crude fibre	Ash
3 times harvest	11.90 <sup>b</sup>	3.11ª	40.02ª	17.57ª	9.11ª
2 times harvest	19.56ª	3.48ª	41.35ª	15.89ª	9.00ª
1 time harvest	16.79 <sup>b</sup>	3.21ª	45.15ª	15.26ª	8.67ª
Lsd (0.05)	4.52	NS	NS	NS	NS

#### DISCUSSION

Agboola and Ayodele<sup>10</sup> recommended 3, 0.15%, 8-10 mg kg<sup>-1</sup>, 0.2, 2.0 and 0.26 C mol kg<sup>-1</sup> for OM, total N, available P, K, Ca and Mg, respectively as critical level of nutrients required for optimum production of arable crop in southwestern Nigeria. Going by Agboola and Ayodele<sup>10</sup> assertion, the chemical properties of the soils showed that the soils were acidic, low in OM, N, P, K, Ca and Mg; hence, additional soil nutrients were required to increase the fertility of the soil.

The result obtained in this research showed that harvesting interval affected the growth and yield of amaranths. This was in line with the work of Foidl et al.4, who observed that average number of leaves of Moringa oleifera produced per plant increased with time in the experiment performed to show the effect of spacing and harvest frequency on growth and leaf yield of moringa (Moringa oleifera Lam), a leafy vegetable crop. Rambau et al.<sup>11</sup> also observed that stages of maturity had significant effects on leaf area and plant height in the experiment conducted to show the effect of three stages of maturity on the productivity, chemical composition and ruminant degradability of Napier grass (Pennisetum purpureum). The experiment conducted by Bashir et al.<sup>12</sup> to show the ratooning potential of different sugarcane genotypes at varying harvesting dates in Pakistan indicated that the growth and yield of cane varied significantly among all the harvesting dates examined.

The increase in the agronomic parameters in 2nd and 3rd time harvest in this experiment might be as a result of the nutrient absorbed by the plants more than the *Amaranthus tricolor* harvested at once. Three times harvest recorded the highest growth and yield parameters and was in line with the work of Ayeni *et al.*<sup>13</sup>. Though the cumulative agronomic parameters increased at highest harvest interval, yet there was reduction in nutritional quality of *Amaranthus* at 3rd time harvest. This might be partially adduced to senescence but might not be absolutely correct as the leaves harvested at the 3rd stage of harvest still remained green and fresh. Nouman *et al.*<sup>14</sup>, also opined that cutting frequency and height significantly affected both fresh and dry vegetable biomass like moringa.

It was observed that the cumulative yield of the Amaranthus tricolor harvested at intervals were higher than Amaranthus tricolor harvested at once by uprooting. The better performance of Amaranthus harvested at different stages than the Amaranthus harvested at once might be as a result of the method of harvesting that gave room for the regrowth of the side shoots since all the plants were subjected to equal treatments in terms of soil fertility and cultural practices. Maboko and du Plooy<sup>7</sup> reported that harvesting amaranths by cutting resulted in a delay of plant maturity compared to tipping in an experiment conducted to show the effect of plant density and harvesting method on yield components of hydroponically grown amaranth. Ratooning might have caused the plants to absorb more nutrients from the soil. Also, the nutrients absorbed might have translocated to the new leaves that were formed. Ojeniyi and Odedina<sup>15</sup> and Conrad<sup>16</sup> classified N, P, K and Mg as mobile plant nutrients while Fe, Mn, Cu and Ca were classified as immobile plant nutrients. Based on the results obtained, the higher concentration of N, P, K and Mg in 2 and 3 times harvests (especially in 2 times harvest) might be as a result of migration of these plant nutrients to the newly formed shoots. The higher concentration of the plant nutrients in the 2 and 3 times harvests might have increased the plant growth. The lower, protein, fat and NFE content in *Amaranthus tricolor* harvested three times interval showed the reduction in the nutritional quality of the crop. Ansah *et al.*<sup>17</sup> observed reduction in crude protein content of crops as harvesting intervals increased as equally observed in this work. Okubena-Dipeolu<sup>18</sup>, Mabapa *et al.*<sup>19</sup> and Dev *et al.*<sup>20</sup> argued that nutritional quality of amaranths depends on certain conditions such as the nutrient status of the soil, soil type, type of fertilizer applied to the soil and harvesting period.

#### CONCLUSION

Experiments were conducted in Ondo southwestern Nigeria to determine the effect of ratooning on growth parameters, nutrient uptake and nutritional quality of *Amaranthus tricolor.* This research work ascertained that 3 times harvest had the highest cumulative growth and yield of *Amaranthus* tricolor while 2 times harvest recorded the highest crude protein.

#### SIGNIFICANCE STATEMENTS

There are different harvesting methods used in harvesting *Amaranthus* in southwest Nigeria. This study discovered that ratooning can increase the yield of amaranth up to 3 times harvest and that this can be beneficial for farmers who grow *Amaranthus* as means of livelihood. The study also shows that as the time of harvesting of amaranth increased, the level of crude protein and crude fat were reduced, thus, reducing the nutritional quality of the crop. This study will help the researchers to uncover the critical areas of *Amaranthus* that many researchers were not able to explore. Thus a new theory on improving the nutritional quality of *Amaranthus* at 3rd time harvest may be arrived at.

#### REFERENCES

- 1. Prasanna, S.M. and P.S. Pattar, 2014. Successful ratoon management in sugarcane. Res. Rev.: J. Agric. Allied Sci., 3: 39-46.
- Gichunge, C.N., F. Kidwaro, O. Ohiokpehai and E. Oyoo, 2008. Effects of picking frequency on the yield of the grain amaranth (*Amaranthus hypochondriacus*) grown in a greenhouse. Acta Hortic., 806: 181-186.

- Chadha, M.L., M.O. Oluoch and D. Silue, 2007. Promoting indigenous vegetables for health, food security and income generation in Africa. Acta Hortic., 762: 253-262.
- Foidl, N., R.N. Bennett, W.O. Ellis, G.M. Timpo and N.K. Amaglo, 2007. Effect of spacing and harvest frequency on the growth and leaf yield of Moringa (*Moringa oleifera* Lam), a leafy vegetable crop. Ghana J. Hort., 6: 33-40.
- Ayub, M., M.A. Nadeem, M. Tahir, M. Ibrahim and M.N. Aslam, 2009. Effect of nitrogen application and harvesting intervals on forage yield and quality of pearl millet (*Pennisetum americanum* L.). Pak. J. Life Soc. Sci., 7:185-189.
- Adewole, M.B. and O.A. Dedeke, 2012. Growth performance, yield and nutritional quality of *Amaranthus cruentus* L. under repeated applications of poultry manures. Ife J. Sci., 14: 345-355.
- Maboko, M.M. and C.P. du Plooy, 2011. Effect of plant density and harvesting method on yield components of hydroponically grown amaranth. Acta Hortic., 947: 415-421.
- 8. Harpstead, M.I., 1973. The classification of some Nigerian soils. Soil Sci., 116: 437-443.
- AOAC., 1990. Official Methods of the Analysis of the Analytical Chemists. Vol. 2, 15th Edn., AOAC Inc., Virginia, USA., pp: 69-83.
- Agboola, A.A. and O.J. Ayodele, 1985. Prospects and problems of using soil testing for adoption of fertilizer use in ekiti-akoko agricultural development project area. Proceedings of the Workshop on Appropriate Technologies for Farmers in Semi-Arid West Africa, April 2-5, 1985, Purdue University, West Lafeyette, pp: 123-136.
- Rambau, M.D., F. Fushai and J.J. Baloyi, 2016. Productivity, chemical composition and ruminal degradability of irrigated Napier grass leaves harvested at three stages of maturity. S. Afr. J. Anim. Sci., 46: 398-408.
- 12. Bashir, S., M.U. Hassan, N. Fiaz, Z. Khan and Z. Ali, 2013. Ratooning potential of different promising sugarcane genotypes at varying harvesting dates. J. Agric. Biol. Sci., 8:437-440.
- Ayeni, L.S., E.A. Okubena-Dipeolu, D.A. Oladepo and K.J. Oyebamiji, 2018. Growing amaranthus as a means of livelihood among peasant farmers in Africa. Int. J. Plant Soil Sci., 22: 1-5.
- Nouman, W., M.T. Siddiqui, S.M.A. Basra, H. Farooq, M. Zubair and T. Gull, 2013. Biomass production and nutritional quality of *Moringa oleifera* as a field crop. Turk. J. Agric. Forest., 37: 410-419.
- 15. Ojeniyi, S.O. and A.S. Odedina, 2014. Basic Plant Nutrients. Dominion Publishers, Ibadan, Nigeria, pp: 60.

- 16. Conrad, J., 2017. Mobile and immobile nutrients. Rancho Regensis North of Valladolid, Yucatan Newsletter, Mexico.
- Ansah, T., E.L.K. Osafo and H.H. Hansen, 2010. Herbage yield and chemical composition of four varieties of Napier (*Pennisetum purpureum*) grass harvested at three different days after planting. Agric. Biol. J. North Am., 1: 923-930.
- Okubena-Dipeolu, E.A., 2015. Effects of organic, organominral and NPK fertilizer treatments on the sensory evaluation of *Amaranthus cruentus* soup in Ikorodu and LASU. Am. J. Res. Commun., 3: 67-75.
- 19. Mabapa, M.P., K.K. Ayisi and I.K. Mariga, 2017. Effect of planting density and harvest interval on the leaf yield and quality of Moringa (*Moringa oleifera*) under diverse agroecological conditions of northern south Africa. Int. J. Agron., Vol. 2017. 10.1155/2017/2941432.
- Dev, C.M., R.K. Singh, R.N. Meena, A. Kumar and K. Singh, 2012. Production potential and soil fertility status of ratoon sugarcane (*Saccharum officinarum* L.) as influenced by time and level of earthing up and nitrogen levels in north-eastern Uttar Pradesh, India. Sustainable Agric. Res., 2: 143-148.