



Asian Journal of Plant Sciences

ISSN 1682-3974

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Ecological Studies on *Salix* Distribution in Egypt

¹Emad A. Al Sherif, ²Wafaa Amer, ¹Salah Eldin Ali Khodary and ¹Walaa Azmy
¹Department of Botany, Faculty of Science, Beni Suef University, Beni Suef, Egypt
²Department of Botany, Faculty of Science, Cairo University, Egypt

Abstract: The present research studied the ecological factors affecting *Salix* distribution in Egypt. Two species of *Salix* were recorded, *S. tetrasperma* (only male) and *S. mucronata* (both sexes). They were recorded at River Nile canal system and in the Eastern Oasis. Female *S. mucronata* was recorded in the all studied habitats, where the male of same species was recorded only in Fayoum Region. *Salix tetrasperma* neither recorded in Upper Egypt nor Eastern Oasis. Elevation from water surface, soil texture, soil salinity and temperature were the most effective factors affecting the distribution of *Salix* sp.

Key words: *Salix*, distribution, ecology

INTRODUCTION

Salicaceae comprises two main genera, *Salix* (willows) and *Populus* (poplars). It exhibits many characteristics that allow it to grow in habitats subjected to flooding and disturbance through erosion and deposition of sediments (Karrenberg *et al.*, 2002).

Salix is a colonizing floodplain genus characterized by vigorous growth rate and production of a massive root system that can rapidly stabilize stream bank sediments (Grissinger and Bowie, 1984; Hupp, 1992; Van Splunder *et al.*, 1996; Shields *et al.*, 1995; Anonymous, 2005). It also occurs in ditches and on the edges of swamps, lakeshores and other wetland habitats. It is ideal for stream bank restoration because of its easily propagation from cutting, quickly growth and providing rapid soil stabilization (Bentrop and Hoag, 1998; Cronk and Fennessy, 2001). Willow (*Salix*) has been cultivated for an agricultural crop to many purpose, such as for bioenergy (Perttu, 1998), making baskets, made a tea from the bark and some medical uses (Foster and Duke, 1990).

Several environmental factors such as soil moisture and soil texture have been identified as being critical in determining survival and performance of planted willow cutting (Pezeshki *et al.*, 1998; Schaff *et al.*, 2003). Water table decline and the associated water stress have been also, cited as one of the most common causes of mortality in *Salix* species in the field (Mahoney and Rood, 1992; Karrenberg *et al.*, 2002). Inadequate soil moisture leading to plant desiccation was reported as one of the causes of

first year mortality of willow cutting planted in riparian habitats (Shields *et al.*, 1998; Karrenberg *et al.*, 2002). Such unrooted cutting are especially vulnerable to low soil moisture conditions early in the growing season due to transpiration water losses until root systems, with their vascular tissues have developed (Grange and Loach, 1983; Ikeda and Suzuki, 1986).

Genus *Salix* L. in Egypt represented by two wild species, *Salix tetrasperma* Roxb. and *Salix safsaf* Trautv. (latter treated as *Salix mucronata* Thunb.); as recorded by Täckholm (1956, 1974) and Boulos (1999). These species, in addition to its medicinal values, attained a special historic and ethnic importance in Egypt. *Salix* species are distributed in Nile and Faiyum Regions, in addition to Mediterranean coastal strip. Earlier study recorded the presence of both species in western desert Oases (Boulos, 1999). In Egypt, *Salix* sp. are used frequently as a restoration species, while other *Salix* sp. are widely used elsewhere (Svejar *et al.*, 1992; Hoag, 1993, 1995; Anonymous, 2005). *Salix tetrasperma* and *S. mucronata*, as many other *Salix* sp., regenerate from root and shoot fragments and is often planted along stream banks as dormant, unrooted cutting (Bentrop and Hoag, 1998; Cronk and Fennessy, 2001).

Egypt is an arid country which depends almost entirely on the River Nile for its water supply; agriculture is almost totally dependent on this source. It is estimated that the Nile provides 95% of the country's fresh renewable water supply. There is no any earlier study on factors affecting distribution of *Salix* in Egypt. The present study deals with environmental factors affecting the distribution of *Salix* sp. in Egypt.

MATERIALS AND METHODS

Field trips were carried out during the study years 2005-2008. Fifteen trips were carried out covering the River Nile and its water canal system from Upper Egypt (Aswan) to lower (Alexandria) and Baharia Oasis.

Soil sampling and analysis: For each habitat three soil samples were collected from profiles of 10-50 cm depth under *Salix* sp. These samples were then pooled together to form one composite sample, air-dried and thoroughly mixed. Thirty composite samples were obtained representing all *Salix* habitats. Textures were determined by the hydrometer method, providing quantitative data on the percentage of sand, silt and clay. The concentration of soil minerals Na^+ , K^+ , Fe^{3+} , Ca^{2+} and Mg^{2+} in soil were determined using a Perkin 403 atomic absorption spectrophotometer (Perkin-Elmer Corp, Norwalk, Conn., USA) according to Analytical Methods for Atomic Absorption Spectrophotometry, Levinson (1983). Chlorides were quantified by titrating 5 mL of the 1 : 5 soil/distilled water extract against 0.01 N silver nitrate solution using potassium chromate (1%) as indicator. Bicarbonates was determined by titrating 5 mL of the 1.5 soil/distilled water extract against 0.01 N HCl using

phenolphthalein and methyl orange as indicators (Jackson, 1962). pH and conductivity of the soil samples were determined in saturated soil paste extract by pH and conductivity meters, respectively. Temperature measurements were obtained from the Egyptian Meteorological Station. The percentage of organic carbon in the soil was determined by Walkely and Black's rapid titration method according to Ryan *et al.* (1996).

Statistical analysis: Pearson correlation coefficient has been calculated to find out the correlations between *Salix* sp. and soil parameters.

RESULTS

The extensive survey in the present study reported two *Salix* species, *S. tetrasperma* and *S. mucronata* (male of the former species and the two sexes of the other one) (Table 1, Fig. 1). *Salix* species were recorded, in canal banks, ditches, on the edges of swamps and lakeshores, a habitats characterized by high moist content.

Table 2 shows that soil analysis where *Salix* found characterized by electric conductivity (as a salinity indicator) ranging from 0.4 to 2.9 mS cm^{-1} , slightly alkaline (with mean pH 7.4). *Salix* sp. reported in varied soil textures but mainly occurred in a medium soil texture, loamy sandy soil. Female *S. mucronata* reported in




Fig. 1: Distribution of *Salix* sp. in Egypt

Table 1: Distribution of *Salix* sp. in different parts of Egypt

Species	Upper Egypt	Middel Egypt	Lower Egypt	Fayoum	Bahria Oasis
<i>S. tetrasperma</i> (male)	-	+	+	+	-
<i>S. mucronata</i> (male)	+	+	+	+	+
<i>S. mucronata</i> (female)	-	-	-	+	-

+: Present, -: Absent

Table 2: Elevation from water surface and soil parameters of different parts of Egypt

Parameters	Upper Egypt (6)*	Middel Egypt (7)	Lower Egypt (7)	Fayoum (5)	Bahria Oasis (5)
Elevation	1.1±0.1**	0.8±0.1	0.9±0.2	0.5±0.1	0.8±0.2
Sand (%)	85.0±1.4	82.0±0.1	83.2±1.4	81.5±1.7	83.5±1.7
pH	7.3±0.3	7.5±0.4	7.2±0.2	7.4±0.3	7.9±0.4
Organic matter (%)	1.1±0.01	0.3±0.01	1.2±0.1	1.8±0.4	0.1±0.01
EC (mS cm ⁻¹)	0.4±0.02	0.5±0.1	1.3±0.3	0.6±0.1	2.9±0.5
Na (mg kg ⁻¹) soil	289.0±23.1	2500.0±152.6	448.0±53.2	1150.0±47.8	6073.0±140.5
K (mg kg ⁻¹) soil	11.0±1.3	109.0±13.7	97.0±16.5	82.0±52.3	257.0±83.2
Ca (mg kg ⁻¹) soil	572.0±53.4	540.0±42.3	560.0±47.6	365.0±42.1	570.0±74.2
Mg (mg kg ⁻¹) soil	144.0±41.2	180.0±14.3	90.0±12.3	90.0±18.9	185.0±12.6
Fe (mg kg ⁻¹) soil	73.0±13.2	28.0±41.2	35.0±13.2	16.0±2.1	288.0±42.6

*: No. of analyzed soil samples; ** Mean±SE

Table 3: Some soil characters of habitats where *Salix* sp. recorded

Soil factor	<i>S. mucronata</i>	<i>S. tetrasperma</i>
pH	7.35±0.57*	7.630±0.42
EC (mS cm ⁻¹)	3.43±4.53	0.660±0.41
Organic matter (%)	1.28±0.74	0.646±0.533
HCO ₃ (%)	0.21±0.09	0.250±0.08
Cl (%)	0.19±0.1	0.097±0.092
Sand (%)	84.98±8.8	85.800±4.26
Clay (%)	10.10±5.17	9.500±3.11
Silt (%)	4.47±6.6	4.380±3.23
Na (mg kg ⁻¹) soil	289.30±63.2	250.000±74.3
K (mg kg ⁻¹) soil	4.68±0.23	31.000±11.3
Ca (mg kg ⁻¹) soil	370.00±86.3	700.000±85.36
Mg (mg kg ⁻¹) soil	450.00±45.7	950.000±102.3
Fe (mg kg ⁻¹) soil	113.90±32.4	74.100±4.2

*: Mean±SE


different canal banks all over the country, where the male was recorded only in Fayium Region. The male *S. tetrasperma* neither recorded in Upper Egypt nor in Baharia Oasis. It appeared from Beni Suf governorate downward to Alexandria.

With regard to elemental analysis, *S. mucronata* prefers soil with high content of organic matter, Na and Fe than *S. tetrasperma* (Table 3). While, *S. tetrasperma* was recorded in soil with high concentration from Mg and Ca (ranged from 90 to 1850 mg kg⁻¹ soil and from 365 to 900 mg kg⁻¹ soil, respectively), compared with *S. mucronata* existed in soil with low concentration of these elements (from 37.8 to 858 mg kg⁻¹ soil of magnesium and from 118.4 to 810 mg kg⁻¹ soil of calcium) (Table 3).

Temperature recorded a decrease with the direction downward reaching the minimum at Alexandria (Fig. 2). Pearson correlations analysis (Table 4) showed a significant positive correlation of *S. mucronata* with organic matter content and significant negative correlation of *S. tetrasperma* with both soil salinity and

Table 4: Pearson correlations between *Salix* sp. and some soil parameters

Parameters	<i>S. tetrasperma</i> ♂	<i>S. mucronata</i> ♂	<i>S. mucronata</i> ♀
Sand (%)	-0.535	0.241	0.255
EC (mS cm ⁻¹)	-0.436	-0.524	0.213
OM (%)	0.290	+0.498	0.354
Elevation	-0.485	-0.475	-0.542
Na	-0.440	+0.245	+0.235
K	+0.221	+0.248	+0.432
Ca	-0.412	-0.147	-0.257
Mg	-0.231	-0.214	-0.241
Fe	-0.111	-0.368	-0.324

Fig. 2: Average monthly temperatures recorded in *Salix* habitats

sodium content. All species showed a significant negative correlation with elevation from water surface.

DISCUSSION

In the present study *Salix* sp. were recorded only on River Nile water systems and besides wells in Baharia Oasis. The earlier finding because of *Salix* roots prefer growing in the saturated groundwater zone and will rapidly elongate to maintain contact with free water (Amlin and Rood, 2002). Moisture content and soil texture are the most important factors affecting *Salix* growth (Hansen and Phipps, 1983; Phipps *et al.*, 1983; Woods

and Cooper, 2005). Both fine and coarse soil texture components contributed to lower survival rate. Moderate soil texture allowing for an aerated soil condition that also maintained some water-holding capacity. In contrast, fine-textured sediments led to poor plant performance due to compaction and the associated low aeration. Sandy soils, on the other hand, have larger pore spaces but lower water-holding capacity, increasing the risk for periodic drought stress for plants on stream banks. Many studies documented the importance of rapid root growth for the survival and growth of *Salix* trees (Mahoney and Rood, 1992; Segelquist *et al.*, 1993; Van Splunder *et al.*, 1996; Kranjcec *et al.*, 1998; Scott *et al.*, 1999; Horton and Clark, 2001; Amlin and Rood, 2002). The absence of *S. tetrasperma* from Upper Egypt may be due to high temperature. Lammeranner *et al.* (2005) proved the suitability of *S. tetrasperma* for winter plantation at low temperature. Its absence from Oases can be explained according to its intolerant to salinity. Jackson *et al.* (1990) reported the intolerance of *S. gooddingii* to soil salinity.

The significant negative correlation between *Salix* and elevation from water surface because *Salix* species are relatively shallow-rooted, drought-intolerant tree species. (Horton and Clark, 2001; Glenn and Nagler, 2005), they prefer low elevation from water surface. Fine roots of these trees are concentrated in the capillary fringe, just above the water table; they are sensitive to fluctuation in water table depth, particularly on coarse soil with a narrow capillary fringe. The occurrence of *S. tetrasperma* and both male and female *S. mucronata* in Fayoum due to presence of low elevation from water surface. Fayoum is the largest Oases in Egypt, situated 90 km south-west of Cairo, 75 km from Giza Pyramids. It considers a natural depression in the desert, linked to the River Nile by a branch called Bahr Yosuf and characterized by high quantity of water with nearly permanent full canal banks.

CONCLUSION

It is advisable to cultivate *Salix* sp. By cutting in a saturated soil or at canal banks with low elevation from water surface.

REFERENCES

- Amlin, N.M. and S.B. Rood, 2002. Comparative tolerances of riparian willows and cottonwoods to water-table decline. *Wetlands*, 22: 338-346.
- Anonymous, 2005. Streambank Soil Bioengineering. Technical Note 14-I: NRCS Draft Stream Restoration Design Handbook, US Department of Agriculture Natural Resources Conservation Service, Washington DC. ISBN: 0-934213-59-3.
- Bentrup, G. and J.C. Hoag, 1998. The Practical Streambank Bioengineering Guide: Interagency Riparian/Wetland Plant Development Project. SDA-Natural Resources Conservation Service, Plant Materials Center, Aberdeen, ID., ISBN: 0934213593.
- Boulos, L., 1999. Flora of Egypt. Vol. 1, AL-Hadara Publishing, Cairo, Egypt, pp: 419.
- Cronk, J.K. and M.S. Fennessy, 2001. Wetland Plants: Biology and Ecology. CRC Press, LLC., Boca Raton, FL.
- Foster, S.A. and J.A. Duke, 1990. A Field Guide to Medicinal Plants (Eastern/Central). Peterson Field Guide Series, Houghton Mifflin Co., Boston, MA.
- Glenn, E.P. and P.L. Nagler, 2005. Comparative ecophysiology of *Tamarix ramosissima* and native trees in western US riparian zones. *J. Arid Environ.*, 61: 419-446.
- Grange, R.I. and K. Loach, 1983. The water economy of unrooted leafy cutting. *J. Hortic. Sci.*, 58: 9-17.
- Grissinger, E.H. and A.J. Bowie, 1984. Material and site controls of stream bank vegetation. *Trans. Am. Soc. Agric. Eng.*, 27: 1829-1835.
- Hansen, E.A. and H.M. Phipps, 1983. Effect of soil moisture tension and preplant treatments on early growth of hybrid Populus-hardwood cutting. *Can. J. For. Res.*, 13: 458-464.
- Hoag, J.C., 1993. How to Plant Willows and Cottonwoods for Riparian Rehabilitation. USDA Natural Resources Conservation Service, Boise, Idaho.
- Hoag, J.C., 1995. Using dormant pole cutting to revegetate riparian areas. Proceedings of the 5th International Rangeland Congress, Jul. 23-28, Salt Lake City, Utah, pp: 1-2.
- Horton, J.L. and J.L. Clark, 2001. Water table decline alters growth and survival of *Salix gooddingii* and *Tamarix chinensis* seedlings. *For. Ecol. Manage.*, 140: 239-247.
- Hupp, C.R., 1992. Riparian vegetation recovery patterns following stream channelization: A geomorphic perspective. *Ecology*, 73: 1209-1226.
- Ikeda, T. and T. Suzuki, 1986. Influence of hydraulic conductance of xylem on water status in cutting. *Can. J. For. Res.*, 16: 98-102.
- Jackson, M.L., 1962. Soil Chemical Analysis. 1st Edn., Constable and Company Ltd., Prentice-Hall, Inc., New Jersey, USA, London, pp: 934.
- Jackson, J., J.T. Ball and M.R. Rose, 1990. Assessment of the Salinity Tolerance of Eight Sonoran Desert Riparian Trees and Shrubs. Final Report for Bureau of Reclamation, Yuma Projects Office, Yuma, Arizona, pp: 102.
- Karrenberg, S., P.J. Edwards and J. Kollmann, 2002. The life history of Salicaceae living in the active zone of floodplains. *Freshwater Biol.*, 47: 733-748.

- Kranjcec, J., J.M. Mahoney and S.B. Rood, 1998. The responses of three riparian cottonwood species to water table decline. For. Ecol. Manage., 110: 77-87.
- Lammeranner, W., H.P. Rauch and G. Laaha, 2005. Implementation and monitoring of soil bioengineering measures at a landslide in the Middle Mountains of Nepal. Plant Soil, 278: 159-170.
- Levinson, A.A., 1983. Analytical methods for atomic absorption spectrophotometry, Perkin-elmer corp., Norwalk, Connecticut. Geochim. Cosmochimica Acta, 33: 1315-1316.
- Mahoney, J.M. and S.B. Rood, 1992. Response of a hybrid poplar to water table decline in different substrates. For. Ecol. Manage., 54: 141-156.
- Perttu, K., 1998. Environmental justification for short-rotation forestry in Sweden. Biomass Bioenergy, 15: 1-6.
- Pezeshki, S.R., P.H. Anderson and F.D. Shields, 1998. Effects of soil moisture regimes on growth and survival of black willow (*Salix nigra*) posts (cutting). Wetlands, 18: 460-470.
- Phipps, H.M., E.A. Hansen and A.S. Fege, 1983. Pre-Plant Soaking of Dormant Populus hardwood Cutting. U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station, St. Paul, Minnesota.
- Ryan, J., S. Garabet, K. Harmsen and A. Rashid, 1996. A Soil and Plant Analysis Manual Adapted for the West Asia and North Africa Region. International Center for Agricultural Research in the Dry Areas (ICARDA), Aleppo, Syria, ISBN: 92-9127-118-7.
- Schaff, S.D., S.R. Pezeshki and F.D. Shields, 2003. Effects of soil conditions on survival and growth of black willow cutting. Environ. Manage., 31: 748-763.
- Scott, M.L., P.B. Shafroth and G.T. Auble, 1999. Responses of riparian cottonwoods to alluvial water table declines. Environ. Manage., 23: 347-358.
- Segelquist, C.A., M.L. Scott and G.T. Auble, 1993. Establishment of *Populus deltoides* under simulated alluvial groundwater declines. Am. Midland Naturalist, 130: 274-285.
- Shields, F.D., C.M. Cooper and S.S. Knight, 1995. Experiment in stream restoration. J. Hydraulic Eng., 121: 494-502.
- Shields, F.D., S.R. Pezeshki and P.H. Anderson, 1998. Probable Causes for Willow Post Mortality: Engineering Approaches to Ecosystem Restoration. ASCE., Reston, VA.
- Svejcar, T.J., G.M. Riegel, S.D. Conroy and J.D. Trent, 1992. Establishment and growth potential of riparian shrubs in the northern Sierra Nevada. Proceedings of the Symposium on Ecology and Management of Riparian Shrub Communities, 1992, US Department of Agriculture, Forest Service, pp: 1-1.
- Täckholm, V., 1956. Students Flora of Egypt. Anglo-Egyptian Bookshop, Egypt.
- Täckholm, V., 1974. Students Flora of Egypt. 1st Edn., Cairo University, Cooperative Printing Co., Beirut.
- Van Splunder, I., L.A.C.J. Voesenek, H. Coops, X.J.A.D. Vries and C.W.P.M. Blom, 1996. Morphological responses of seedlings of four species of Salicaceae to drought. Can. J. Bot., 74: 1988-1995.
- Woods, S.W. and D.J. Cooper, 2005. Hydrologic factors affecting initial willow seedling establishment along a subalpine stream, Colorado, USA. Arctic Antarctic Alpine Res., 37: 636-643.