



Asian Journal of Scientific Research

ISSN 1992-1454

science
alert
<http://www.scialert.net>

ANSI*net*
an open access publisher
<http://ansinet.com>



Research Article

Nitrogen and Phosphorus Waste Production from Different Fish Species Cultured at Floating Net Cages in Lake Maninjau, Indonesia

¹Hafrijal Syandri, ¹Azrita and ²Ainul Mardiah

¹Department of Aquaculture, Faculty of Fisheries and Marine Science, University of Bung Hatta, Padang, Indonesia

²Department of Aquaculture, Faculty of Marine and Fisheries Science, Nahdlatul Ulama University of West Sumatera, Padang, Indonesia

Abstract

Background and Objective: Aquaculture operations that use floating net cages have become one of the primary mean of intensive fish-culture in Lake Maninjau. The fish-culture species studied were *Cyprinus carpio* (*C. carpio*) (T₁), *Oreochromis niloticus* (*O. niloticus*) (T₂), *Osphronemus goramy* (*O. goramy*) (T₃) and *Clarias gariepinus* (*C. gariepinus*) (T₄). The objective of the research was to estimate the nitrogen (N) and phosphorus (P) loads into Lake Maninjau. **Materials and Methods:** The capacity of floating net cages was approximately 32 m³ (4 × 4 × 2 m) with densities of 32 fish m⁻³ in triplicate groups. Approximately 1,500 kg of feed was used in each cage during the experiment. The difference of N and P loads from different fish species were analyzed using one-way ANOVA (SPSS 16.0) computer software. **Results:** The total N loads into the water bodies from T₁, T₂, T₃ and T₄ were estimated at 37.93 ± 2.59, 49.90 ± 5.17, 45.90 ± 4.18 and 20.35 ± 4.12 kg t⁻¹ of fish production, respectively. The P load was estimated to be 18.30 ± 0.12, 20.01 ± 0.99, 22.60 ± 0.80 and 13.93 ± 1.47 kg t⁻¹ of fish production, respectively. Every ton of feed consumed by each fish species will contribute as much as 38.26 ± 2.55, 35.68 ± 1.69, 32.12 ± 0.39 and 48.99 ± 2.35 kg N load into the water bodies, respectively. The P load was 11.45 ± 2.43, 9.11 ± 0.21, 8.34 ± 0.04 and 12.51 ± 0.30 kg, respectively. **Conclusion:** The *C. gariepinus* species is preferred for aquaculture operations at Lake Maninjau, because it minimizes N and P load releases into water bodies which can maintain sustainable aquaculture operations.

Key words: Aquaculture, floating net cages, N and P mass balance, FCR, environmental factors

Received: August 28, 2017

Accepted: November 16, 2017

Published: March 15, 2018

Citation: Hafrijal Syandri, Azrita and Ainul Mardiah, 2018. Nitrogen and phosphorus waste production from different fish species cultured at floating net cages in Lake Maninjau, Indonesia. Asian J. Sci. Res., 11: 287-294.

Corresponding Author: Hafrijal Syandri, Aquaculture laboratory, Faculty of Fisheries and Marine Science, University of Bung Hatta, Padang, Indonesia Tel +62751-7051678

Copyright: © 2018 Hafrijal Syandri *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

The high levels of nutrients and suspended solids released into water bodies from aquaculture operations is one of the major environmental problems causing water pollution¹⁻⁵. Intensive aquaculture operations can result in the release of dissolved organic and inorganic nutrients, such as nitrogen (N) and phosphorus (P)⁶⁻⁸. N and P levels from intensive aquaculture operations can cause or accelerate eutrophication in natural water systems⁹⁻¹¹. In addition, N and P levels released into water bodies depend on diet composition^{12,13}, type of feed¹⁰, fish species¹⁴, feed conversion ratio¹⁵, stocking densities, feed quality¹⁶, fish mass mortality¹⁷ and the local environment¹⁸. Lazzari and Baldisserotto¹⁹ state that N and P are the primary end products of fish loading, which has had an effect on fish rearing waters and the environment.

Aquaculture around the world has grown at a rapid rate in recent years, including in Indonesia^{20,21}. Medium and large scale freshwater aquaculture operations in Indonesia were conducted in the lake, reservoir and river²¹⁻²³. Fish-cultured species in this location were *Cyprinus carpio*, *Oreochromis niloticus*, *Osphronemus goramy*, *Hemibagrus nemerus*, *Clarias gariepinus*, *Pangasius* spp. and *Leptobarbus hoeveni*^{20,24,25}.

Lake Maninjau is tecto-volcanic with a surface area of 99.5 km², both features that serve very important roles to many Indonesians for aquaculture operations of the *O. niloticus* and *C. carpio* in floating net cages^{8,22}. Total number of floating net cages in the years of 2013, 2014 and 2015 are 16,120, 16,580 and 20,608 U, respectively^{8,22,24}.

In the past decade, the water quality of Lake Maninjau has been decreasing due to the loading of organic matter from aquaculture operations of *C. carpio* and *O. niloticus*^{22,26}. However, upwelling that has occurred at Lake Maninjau every year has caused a lack of oxygen in the water which is a result of mass fish mortality¹⁶. Furthermore, since the year 2014, aquaculture operations of *O. goramy* and *C. gariepinus* have been successful in floating net cages. Both species were resistant to poor water quality and had a wide market in Indonesia, because they are favored by consumers. These species also have high prices and a high demand in the market.

The aim of the study was to estimate the quantitative, values of N and P loads released from each species cultured in floating net cages on Lake Maninjau. The results of this study were used to increase the scientific understanding of the effects of N and P load releases of different fish species into the water at Lake Maninjau.

MATERIALS AND METHODS

Study area: The experiment was conducted in Lake Maninjau of West Sumatera Province, Indonesia. The geographical position is S:00°12'26.63"-S:00°25'02.80" and E:100°07'43.74"-E:100°16'22.48" and it is located at an altitude of 461.50 m above sea level²⁴. Based on the Schmidt Ferguson climate classification, Lake Maninjau has characteristics of climate types A and an annual rainfall of 3,490 mm.

In this study, the *C. carpio*, *O. niloticus*, *O. goramy* and *C. gariepinus* fingerlings were designated T₁, T₂, T₃ and T₄, respectively. All species were collected from a private hatchery in the Luak District, Lima Puluh Kota Regency, West Sumatera Province. The fingerlings were transferred by truck to the Research Center of the Faculty of Fisheries and Marine Science, Bung Hatta University near the Lake Maninjau. Each fish species was treated with a prophylactic formalin bath (100 mg L⁻¹) for 1 h to remove external parasites and acclimatized in a floating net cage (4×4×2 m) for 1 month prior to the experiment. The average initial weight of the T₁, T₂, T₃ and T₄ fingerlings were 56.79±4.45, 53.08±1.60, 55.33±1.14 and 51.18±1.59 g, respectively.

This study was conducted between March, 2017 and June, 2017 (100 days). All species were cultured in the floating net cages. Each floating net cage had a capacity of approximately 32 m³ (4×4×2 m) and was constructed using a 10 mm mesh size sieve. The fingerlings were fed by a commercial feed (pelleted) with drowned type during 100 days of the experiment. The approximate composition of the feed was 12% moisture content, 29% crude protein, 6% crude lipid, 12% crude fiber and 6% crude ash.

The stocking density was 130 fish m⁻³ (4,160 fish/cage). There were 3 replicates for each fish species in each experiment. During the experiment, 1,500 kg of feed was used. The fish were fed daily at a rate of 4% of their biomass at 9:00, 14:00 and 18:00. All fish mortalities were removed and weighed daily. Fish mortality was replaced in each treatment. The amount of feed provided was adjusted according to temporal changes of biomass and growth of fish in the floating net cages.

Measurements parameters: To determine the growth performance of the fish, the following parameters were calculated based on Aryani *et al.*²⁷:

$$\text{Final mean weight, feed conversion ratio (FCR)} = \frac{\text{Total feed fed (g)}}{\text{Total wet weight gain (g)}}$$

and

$$\text{Survival rate (SR\%)} = \frac{\text{Number of fish survived}}{\text{Number of fish stocked}} \times 100$$

Water quality: The water transparency was measured with a Secchi disc. The water samples were collected at a depth of 10 cm from each floating net cage to determine the dissolved oxygen (DO) levels. An oxygen meter (YSI model 52, Yellow Spring Instrument Co., Yellow Springs, OH, USA) was used *in situ* and pH values were determined using a pH meter (Digital Mini-pH Meter, 0-14 pH, IQ Scientific, Chemo-science [Thailand] Co., Ltd, Thailand). Water temperature was measured using a thermometer (Celsius scale). The levels of alkalinity and hardness of the water in each replication were measured according to standard procedures²⁸. The water quality parameters were measured once every month.

Analytical methods: Nitrogen (N) concentrations (as % of dry weight) of feed and fish were determined by standard methods of the Association of Official Analytical Chemists²⁹. The P concentrations were determined using a spectrophotometer (UV 160 A, Japan) and the molybdate-ascorbic acid method indicated by the Association of Official Analytical Chemists²⁹ at the Chemistry Laboratory of the University of Bung Hatta Padang, Indonesia. The results were expressed as absorbance at 400 nm. All samples were performed in triplicate.

Estimation of N and P loads: The levels of N and P loads from fish-culture was estimated according to Ackefors and Enell³⁰. The following parameters were analyzed according to the formulas given below:

$$\text{N load (kg of N)} = [(\text{Feed} \times \text{Feed}_N) - (\text{Fish} \times \text{Fish}_N)]$$

$$\text{P load (kg of P)} = [(\text{Feed} \times \text{Feed}_P) - (\text{Fish} \times \text{Fish}_P)]$$

where, Feed is total feed used during the experiment, Fish is wet weight of fish produced per harvested, Feed_N is N content of feed. Feed_P is P content of feed (expressed as % of dry weight). Fish_N is N content of fish and Fish_P is P content of fish (expressed as % of wet weight).

$$\text{N and P loads from the production of 1 t of fish} = (1 \text{ t feed} \times \text{FCR} \times \text{Feed}_{\text{N and P content}}) - (1 \text{ t fish} \times \text{Fish}_{\text{N and P content}})$$

$$\text{N and P loads from 1 t of feed consumption} = \frac{(1 \text{ t feed} \times \text{N or P content of feed})}{(\text{FCR})}$$

Statistical analysis: The mean values for final weight, feed conversion ratio, mortality parameters of different treatments and monthly variations of water quality parameters, were subjected to a one-way ANOVA test followed by Duncan's new multiple range test³¹. All statistical analyses were performed using SPSS software (version 16.0 for Windows, SPSS Inc., Chicago, IL). The standard deviation of each parameter and treatment was determined and expressed as the Mean ± SD. The treatment effects were considered to be significant at p < 0.05.

RESULTS

The results for certain growth parameters, FCR, mortality and chemical analyses, from each feed and fish species are presented in Table 1. The difference in fish species has a significant (p < 0.05) effect on the final mean weight, FCR and mortality. The N and P content of the feed were 5.52 ± 0.29 and 1.41 ± 0.03%, respectively. The N and P content of each fish species is presented in Table 1. Monthly variations in the water quality parameter in Lake Maninjau are as indicated in Table 2. There was no significant difference in the water transparency, water temperature, dissolved oxygen, pH, alkalinity or hardness values for months of March, April, May and June, of 2017.

Table 1: Growth parameters and chemical analysis of different fish species

Parameters	Species				
	T ₁	T ₂	T ₃	T ₄	T ₅
Initial mean weight (g)	56.79 ± 1.77	53.08 ± 1.60	55.33 ± 1.14		51.18 ± 1.59
Final mean weight (g)	182.45 ± 2.00 ^a	175.10 ± 2.30 ^b	148.55 ± 7.53 ^c		233.30 ± 7.51 ^d
Feed conversion ratio (FCR)	1.44 ± 0.02 ^a	1.55 ± 0.03 ^b	1.69 ± 0.03 ^c		1.13 ± 0.10 ^d
Mortality (%)	12.46 ± 0.93 ^a	10.92 ± 0.36 ^b	8.18 ± 0.28 ^c		5.20 ± 0.30 ^d
N content of fish (%)	4.17 ± 0.05	3.55 ± 0.05	4.59 ± 0.07		4.13 ± 0.04
P content of fish (%)	0.20 ± 0.02	0.18 ± 0.02	0.13 ± 0.03		0.29 ± 0.02
N content of feed (%)	5.52 ± 0.29	5.52 ± 0.29	5.52 ± 0.29		5.52 ± 0.29
P content of feed (%)	1.41 ± 0.03	1.41 ± 0.03	1.41 ± 0.03		1.41 ± 0.03

Values are Mean ± SD, *Values in the same row with a different superscript are significantly different (p < 0.05), T₁: *C. carpio*, T₂: *O. niloticus*, T₃: *O. goramy* and T₄: *C. gariepinus*

Table 2: Monthly variations in physicochemical and water quality parameters

	March, 2017	April, 2017	May, 2017	June, 2017
Water transparency (m)				
Mean	1.96 ^a	1.98 ^a	1.99 ^a	1.91 ^a
Standard deviation	0.14	0.07	0.10	0.10
Median	1.85	1.98	2.01	1.94
Minimum-Maximum	1.80-2.10	1.90-2.08	1.85-2.10	1.90-2.00
Temperature (°C)				
Mean	28.20 ^a	27.20 ^a	27.00 ^a	27.00 ^a
Standard deviation	0.83	0.83	28.00	1.22
Median	28.00	27.00	0.70	27.00
Minimum-Maximum	27.00-29.00	26.00-28.00	27.00-29.00	26.00-29.00
Dissolved oxygen (mg L⁻¹)				
Mean	6.21 ^a	5.78	6.11 ^a	6.00 ^a
Standard deviation	0.12	0.44	0.05	0.04
Median	6.24	5.44	6.13	5.96
Minimum-Maximum	6.00-6.31	5.40-6.24	6.03-6.13	5.96-6.05
pH				
Mean	7.67 ^a	7.45 ^a	7.41 ^a	7.68 ^a
Standard deviation	0.05	0.32	0.35	0.07
Median	7.69	7.65	7.51	7.67
Minimum-Maximum	7.62-7.76	7.08-7.71	6.90-7.68	7.62-7.79
Alkalinity (mg L⁻¹)				
Mean	83.79 ^a	78.30	79.12	79.88 ^a
Standard deviation	4.37	1.48	2.44	5.14
Median	80.70	78.50	77.97	83.74
Minimum-Maximum	80.51-88.90	76.00-80.00	76.68-82.00	76.15-84.34
Hardness (mg L⁻¹)				
Mean	64.84 ^a	65.28 ^a	70.64 ^a	70.20 ^a
Standard deviation	3.31	2.10	5.93	5.95
Median	67.00	66.50	73.71	68.70
Minimum-Maximum	61.64-68.50	62.80-67.01	63.83-74.91	64.59-79.30

*Values in the same row with the same superscript are not significantly different (p>0.05)




Fig. 1: Nitrogen and phosphorus loads from the production of 1 t of fish in Lake Maninjau




Fig. 2: Nitrogen and phosphorus loads released into Lake Maninjau from 1 t of feed consumption

Table 3 presented the summaries of the mass balance of N and P content from four fish species, while Fig. 1 and 2 provide an estimation of N and P loads from the production of 1 t of fish and 1 t of feed consumption.

DISCUSSION

The present study was conducted to report the N and P loads introduced by floating net-cages in Lake Maninjau.

These results indicated that species T₄ had a better growth rate compared to those of T₁, T₂ and T₃. The differences in growth rate might be due to the specific growth rate of each fish species. The specific growth rates (SGR, % day⁻¹) for T₁, T₂, T₃ and T₄ used with the feeding rate of 4% were 1.16, 1.19, 0.98 and 1.52, respectively. Numerous studies elsewhere have shown that the specific growth rate (SGR) of each species of fish is different. The SGRs (% day⁻¹) are

Table 3: Mass balance of nitrogen (N) and phosphorus (P) of different fish species

Species	N from feed (kg)	N retained in fish (kg)	N load (kg)	P from feed (kg)	P retained in fish (kg)	P load (kg)	N and P retained in fish (%)	N and P load (%)
T ₁	60.72±3.19 ^a	31.78±0.83 ^a	28.93±2.36 ^a	15.51±0.33 ^a	1.53±0.17 ^a	13.90±0.12 ^a	52.39±1.42 ^{bN} 9.83±0.88 ^{bP}	47.60±1.42 ^{bN} 90.62±0.59 ^{bP}
T ₂	60.72±3.19 ^a	25.25±0.53 ^b	35.46±3.12 ^b	15.72±0.18 ^b	1.28±0.17 ^b	14.43±0.34 ^b	42.87±0.41 ^{bN} 8.17±1.17 ^{aP}	58.34±2.13 ^{bN} 91.83±1.17 ^{aP}
T ₃	60.72±3.19 ^a	29.87±0.22 ^c	30.85±2.12 ^c	15.50±0.30 ^c	0.84±0.18 ^c	13.67±0.15 ^c	50.06±0.00 ^{cN} 4.87±0.75 ^{cP}	49.94±2.00 ^{cN} 94.60±1.05 ^{cP}
T ₄	60.72±3.19 ^a	40.33±0.84 ^d	20.39±3.27 ^d	15.51±0.33 ^d	2.83±0.17 ^d	12.68±0.17 ^d	66.53±3.62 ^{dN} 18.23±0.78 ^{dP}	33.46±3.62 ^{dN} 81.77±0.78 ^{dP}

Data are presented as the (Mean±SD) of triplicate samples, *The difference between means with different lower case letters in a column and the difference between means with different capitalized letters for each parameter (N, P) are statistically significant (p<0.05), T₁: *C. carpio*, T₂: *O. niloticus*, T₃: *O. gora*my and T₄: *C. gariepinus*

1.63 for *C. carpio*³², 2.14 for *O. niloticus*³³, 2.47 for *C. gariepinus*³⁴ and 1.66 for *O. gora*my²⁷.

In this study, the water quality in each floating net cage during the months of March, April, May and June showed no significant differences. The growth of the fish species depends not only the water quality³⁵⁻³⁸ but also on the fish species^{39,40}. Although each fish species used the drowned feed type with a feeding rate of 4%, the feed conversion ratio (FCR) for each species was significantly different (p<0.05) (Table 1). The FCR is usually used to estimate the efficiency of converting feed into body mass. In this study, the lowest FCR value was observed in T₄ (1.13), while the highest was observed in T₃ (1.69). The differences among the FCR values are caused by differences in fish species (T₁, T₂, T₃ and T₄) and possibly also by food habits. Conversely, a lower FCR value indicated that the efficiency of feed utilization was better. An FCR value that is less than 2.0 or very close to 2.0 is considered "good" in the aquaculture industry⁴¹. In contrast, the FCR for Tilapia fish cages in Lake Malawi are between 2.1 and 3.9 and FCR values tended to be higher in recent production cycles. Production cycles during the study period were on average (±SE), 376±9 days long¹⁴.

Negative environmental impacts of cage aquaculture operations have been reported in many parts of the world^{20,42-46}. In this study, the difference in fish species has a significant effect (p<0.05) on the mass balance of N and P (Table 3). N and P retention (kg) was significantly higher in T₄ compared to that of T₁, followed by T₃ and T₂, while N and P load (kg) was significantly higher for T₂ compared to that of T₁, T₃ and T₄. Although the same feed was applied at an equal ratio, the mass balance of N and P differed significantly among all fish species. The reason for this difference might be due to the differences in FCR for each species and there was less difference in genetic improvement for feed consumption. The N loads in T₁, T₂, T₃ and T₄ were 47.60, 58.34, 49.94 and 33.46%, respectively, while the P loads in T₁, T₂, T₃ and T₄ were 90.62, 91.83, 94.60 and 81.77%, respectively. For *Oreochromis karongae* and *O. shiranus* in Lake Malawi, N loads were 59 and 80%, respectively and P loads were 85 and 92%, respectively¹⁴. In addition, for Rainbow Trout (*Oncorhynchus mykiss*) in the Kesikköprü Dam Lake, N loads were 54.37% and P loads were 70.00%¹⁰. According to Yogev *et al.*⁴⁷ fish only use 20-30% of the N in feed and 50% of P in feed, while the remainder is released into the water.

In this study, total N and P load releases into water bodies were different for each ton of fish production in T₁ (37.93±2.59 and 18.30±0.12), T₂ (49.90±5.17 and 20.01±0.99), T₃ (45.90±4.18 and 22.60±0.80) and T₄ (20.35±4.12 and 13093±1.47) kg t⁻¹, respectively (Fig. 1). These differences could be caused by the FCR and the N and

P content of feed and fish. There was a strong linear relationship between FCR and N ($r^2 = 0.87$) and P ($r^2 = 0.99$) loads for the floating net cages. In comparison other values in the literature, show that N released into water bodies (t^{-1} of live-weight fish) for Tilapia, Black Pacu and Trout were 34.7, 25.8 and 66.1 kg, respectively. Alternatively, the P was 3.0, 9.7 and 9.6 kg, respectively⁴³. Other research found that 56.0 kg of N and 10.66 kg P were released for Trout¹⁰ and 64.0 kg of N and 4.6 kg of P were released for Tilapia⁴⁷. The N loading values also varied considerably with fish species with Rainbow trout having the lowest values of 47.3-124.2 kg t^{-1} , while values given by other fishes ranged from 103.5-320.6 kg t^{-1} ¹⁵.

In this study, total N and P load releases into water bodies were different for each ton of feed consumed by T_1 , T_2 , T_3 and T_4 (Fig. 2). This difference is also caused by FCR and feed composition. The FCRs for T_1 , T_2 , T_3 and T_4 were 1.44 ± 0.02 , 1.55 ± 0.03 , 1.69 ± 0.03 and 1.13 ± 0.10 , respectively. There was a strong linear relationship between FCR and N ($r^2 = 0.99$) and P ($r^2 = 0.87$) loads for the floating net cages. The N and P levels in the feed used and the FCRs in the farms directly affected N and P loads for each ton of pelleted feed used. Every ton of feed consumed by each fish species (T_1 , T_2 , T_3 and T_4) will contribute N loads as much as 38.26 ± 2.55 , 35.68 ± 1.69 , 32.12 ± 0.39 , 48.99 ± 2.35 and P loads as much as 11.45 ± 2.43 , 9.11 ± 0.21 , 8.34 ± 0.04 and 12.51 ± 0.30 into Lake Maninjau. Furthermore, releases of N and P t^{-1} into the Kesikköprü Dam Lake for *Oncorhynchus mykiss* were 44.78 and 8.60 kg, respectively¹⁰. The feed composition and feed conversion of aquaculture operations primarily had a negative effect on the environment. In addition, the aquaculture integrated model, recirculating aquaculture systems, site selection, feeding rate, size of the farm and species of cultivated fish should also be considered important factors^{32,43,45,47,48}.

CONCLUSION

The present study observed clear evidence that different species of fish-cultured in floating net cages can release different levels of total N and P into water bodies of Lake Maninjau. The estimated N and P loads from the production of 1 t of fish were significantly lower for *Clarias gariepinus* compared to those for *O. goramy*, *C. carpio* and *O. niloticus* while the estimated N and P loads from 1 t of feed consumption were significantly lower for *O. goramy* compared to other species. Based on the research, the appropriate species cultured in Lake Maninjau was *C. gariepinus* because this species had low N and P loads into water bodies. Alternatively, N and P loads can be reduced by adjusting the stocking densities and feed regime timing of *C. gariepinus*.

This approach will help to reduce the downstream negative effects on the lake and in turn positively affect the water quality.

SIGNIFICANCE STATEMENTS

This study analyzes the different levels of N and P load releases into water bodies of Maninjau Lake from each fish-culture species. The N and P loads were significantly lower for *Clarias gariepinus* compared to other fish species. Fish-culture of *Clarias gariepinus* in Lake Maninjau is an important consideration for fish farmers and authorities in the future due to the lower N and P load releases into water bodies of Lake Maninjau. This fish species is also resistant to poor water quality, has a higher growth rate and is favored by consumers.

ACKNOWLEDGMENTS

This study was funded by a study grant (*Riset Unggulan Perguruan Tinggi*) from the Directorate of Research and Community Service of the Ministry of Research Technology and Higher Education, Republic of Indonesia (No. SP.DIPA-042.06-0/2017).

REFERENCES

- Rosa, R.D.S., A.C.F. Aguiar, I.G. Boechat and B. Gucker, 2013. Impacts of fish farm pollution on ecosystem structure and function of tropical headwater streams. *Environ. Pollut.*, 174: 204-213.
- Zhu, Z.M., X.T. Lin, J.X. Pan and Z.N. Xu, 2014. Effect of cyclical feeding on compensatory growth, nitrogen and phosphorus budgets in juvenile *Litopenaeus vannamei*. *Aquacult. Res.*, 47: 283-289.
- Maccoux, M.J., A. Dove, S.M. Backus and D.M. Dolan, 2016. Total and soluble reactive phosphorus loadings to Lake Erie: A detailed accounting by year, basin, country and tributary. *J. Great Lakes Res.*, 42: 1151-1165.
- Prathumchai, N., C. Polprasert and A.J. Engle, 2016. Phosphorus leakage from fisheries sector-A case study in Thailand. *Environ. Pollut.*, 219: 967-975.
- Horppila, J., H. Holmroos, J. Niemisto, I. Massa and N. Nygren *et al.*, 2017. Variations of internal phosphorus loading and water quality in a Hypertrophic lake during 40 years of different management efforts. *Ecol. Eng.*, 103: 264-272.
- Qing-Jun, M., F. Qi-Yan, W. Qing-Qing, M. Lei and C. Zhi-Yang, 2009. Distribution characteristics of nitrogen and phosphorus in mining induced subsidence wetland in Panbei coal mine, China. *Proc. Earth Planet. Sci.*, 1: 1237-1241.

7. Kawasaki, N., M.R.M. Kushairi, N. Nagao, F. Yusoff, A. Imai and A. Kohzu, 2016. Release of nitrogen and phosphorus from aquaculture farms to Selangor River, Malaysia. *Int. J. Environ. Sci. Dev.*, 7: 113-116.
8. Syandri, H., Elfiondri, Junaidi and Azrita, 2015. Social status of the fish-farmers of floating-net-cages in lake Maninjau, Indonesia. *J. Aquacult. Res. Dev.*, Vol. 7. 10.4172/2155-9546.100039.
9. Cao, L., W.M. Wang, Y. Yang, C.T. Yang, Z.H. Yuan, S.B. Xiong and J. Diana, 2007. Environmental impact of aquaculture and countermeasures to aquaculture pollution in China. *Environ. Sci. Pollut. Res.*, 14: 452-462.
10. Asir, U. and S. Pulatsu, 2008. Estimation of the nitrogen-phosphorus load caused by rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792) Cage-Culture farms in Kesikkopru Dam Lake: A comparison of pelleted and extruded feed. *Turk. J. Vet. Anim. Sci.*, 32: 417-422.
11. Lepori, F and J.J. Roberts, 2017. Effects of internal phosphorus loadings and food-web structure on the recovery of a deep lake from eutrophication. *J. Great Lakes Res.*, 43: 255-264.
12. Abou, Y., A. Saidou, D. Mama, E.D. Fiogbe and J.C. Micha, 2012. Evaluation of nitrogen and phosphorus wastes produced by Nile tilapia (*Oreochromis niloticus* L.) fed Azolla-diets in earthen ponds. *J. Environ. Protect.*, 3: 502-507.
13. Boyd, C.E., C.S. Tucker and B. Somridhivej, 2016. Alkalinity and hardness: Critical but elusive concepts in aquaculture. *J. World Aquacult. Soc.*, 47: 6-41.
14. Gondwe, M.J.S., S.J. Guildford and R.E. Hecky, 2011. Carbon, nitrogen and phosphorus loadings from tilapia fish cages in Lake Malawi and factors in uencing their magnitude. *J. Great Lakes Res.*, 37: 93-101.
15. Islam, M.S., 2005. Nitrogen and phosphorus budget in coastal and marine cage aquaculture and impacts of effluent loading on ecosystem: Review and analysis towards model development. *Mar. Pollut. Bull.*, 50: 48-61.
16. Herbeck, L.S., D. Unger, Y. Wu and T.C. Jennerjahn, 2013. Ef uent, nutrient and organic matter export from shrimp and fish ponds causing eutrophication in coastal and back-reef waters of NE Hainan, tropical China. *Continental Shelf Res.*, 57: 92-104.
17. Syandri, H., Azrita, Junaidi and A. Mardiah, 2017. Levels of available nitrogen-phosphorus before and after fish mass mortality in Maninjau Lake of Indonesia. *J. Fish. Aquat. Sci.*, 12: 191-196.
18. Carroll, M.L., S. Coahrane, R. Fieler, R. Velvin and P. White, 2003. Organic enrichment of sediments from salmon farming in Norway: Environmental factors, management practices and monitoring techniques. *Aquaculture*, 226: 165-180.
19. Lazzari, R. and B. Baldisserotto, 2008. Nitrogen and phosphorus waste in fish farming. *Bolet. Inst. Pesca*, 34: 591-600.
20. Henriksson, P.J.G., N. Tran, C.V. Mohan, C.Y. Chan and U.P. Rodriguez *et al.*, 2017. Indonesian aquaculture futures-evaluating environmental and socioeconomic potentials and limitations. *J. Cleaner Prod.*, 162: 1482-1490.
21. Tran, N., U.P. Rodriguez, C.Y. Chan, M.J. Phillips and C.V. Mohan *et al.*, 2017. Indonesian aquaculture futures: An analysis of fish supply and demand in Indonesia to 2030 and role of aquaculture using the Asia Fish model. *Mar. Policy*, 79: 25-32.
22. Syandri, H., Azrita and Niagara, 2016. Trophic status and load capacity of water pollution waste fish-culture with floating net cages in Maninjau lake, Indonesia. *Ecol. Environ. Conserv.*, 22: 459-466.
23. Rahman, M., 2013. Revitalization of fish cage aquaculture management in Riam Kanan stream of Kalimantan Selatan Province-Indonesia. *Jurnal Iktiologi Indonesia*, 13: 197-203.
24. Syandri, H., Junaidi, Azrita and T. Yunus, 2014. State of aquatic resources Maninjau lake West Sumatra province, Indonesia. *J. Ecol. Environ. Sci.*, 5: 109-113.
25. Suhenda, N., R. Samsudin and E. Nugroho, 2010. [Growth of green catfish (*Hemibagrus nemurus*) fry in floating net cage feed by artificial food with different protein content]. *Jurnal Iktiologi Indonesia*, 10: 65-71.
26. Junaidi, H. Syandri and Azrita, 2014. Loading and distribution of organic materials in Maninjau lake West Sumatra province-Indonesia. *J. Aquacult. Res. Dev.*, Vol. 5. 10.4172/2155-9546.1000278.
27. Aryani, N., Azrita, A. Mardiah and H. Syandri, 2017. Influence of feeding rate on the growth, feed efficiency and carcass composition of the giant gourami (*Osphronemus goramy*). *Pak. J. Zool.*, 49: 1775-1781.
28. APHA, 1995. Standard Methods for Examination of Water and Wastewater. 19th Edn., American Public Health Association, Washington DC., USA.
29. AOAC., 1990. Official Methods of Analysis. 13th Edn., Association of Official Analytical Chemists (AOAC), Washington, DC., USA.
30. Ackefors, H. and M. Enell, 1990. Discharge of nutrients from Swedish fish farming to adjacent sea areas. *Ambio*, 19: 28-35.
31. Duncan, D.B., 1955. Multiple range and multiple F tests. *Biometrics*, 11: 1-42.
32. Ahmed, A.R., A.J. Moody, A. Fisher and S.J. Davies, 2013. Growth performance and starch utilization in common carp (*Cyprinus carpio* L.) in response to dietary chromium chloride supplementation. *J. Trace Elem. Med. Biol.*, 27: 45-51.
33. Skov, P.V., C.P. Duodu and D. Adjei-Boateng, 2017. The influence of ration size on energetics and nitrogen retention in tilapia (*Oreochromis niloticus*). *Aquaculture*, 473: 121-127.
34. Widyantorow, W., Sarjito and D. Harwanto, 2014. The effect of fasting time on the growth and bloods profile of catfish (*Clarias gariepinus*) in the recirculating system. *J. Aquacult. Manage. Technol.*, 2: 103-108.

35. Mukai, Y. and L.S. Lim, 2011. Larval rearing and feeding behavior of African catfish, *Clarias gariepinus* under dark conditions. J. Fish. Aquat. Sci., 6: 272-278.
36. Masiha, A., E. Ebrahimi, N.M. Soofiani and M. Kadivar, 2015. Effect of dietary canola oil level on the growth performance and fatty acid composition of fingerlings of rainbow trout (*Oncorhynchus mykiss*). Iran. J. Fish. Sci., 14: 336-349.
37. Asuwaju, F.P., V.O. Onyeche, K.E. Ogbuebunu, H.F. Moradun and E.A. Robert, 2014. Effect of feeding frequency on growth and survival rate of *Clarias gariepinus* fingerlings reared in plastic bowls. J. Fish. Aquatic Sci., 9: 425-429.
38. Paray, B.A., M.K. Al-Sadoon and M.A. Haniffa, 2015. Impact of different feeds on growth, survival and feed conversion in stripped snakehead *Channa striatus* (Bloch 1793) larvae. Indian J. Fish., 62: 82-88.
39. Milstein, A., A. Kadir and M.A. Wahab, 2008. The effects of partially substituting Indian carps or adding silver carp on polycultures including small indigenous fish species (SIS). Aquaculture, 279: 92-98.
40. Medeiros, M.V., J. Aubin and A.F. Camargo, 2017. Life cycle assessment of fish and prawn production: Comparison of monoculture and polyculture freshwater systems in Brazil. J. Cleaner Prod., 156: 528-537.
41. Bag, N., S. Moulick and B.C. Mal, 2016. Effect of stocking density on water and soil quality, growth, production and profitability of farming Indian major carps. Indian J. Fish, 63: 39-46.
42. Kassam, L. and A. Dorward, 2017. A comparative assessment of the poverty impacts of pond and cage aquaculture in Ghana. Aquaculture, 470: 110-122.
43. Avadi, A., N. Pelletier, J. Aubin, S. Ralite, J. Nunez and P. Freon, 2015. Comparative environmental performance of artisanal and commercial feed use in Peruvian freshwater aquaculture. Aquaculture, 435: 52-66.
44. David, G.S., E.D.D. Carvalho, D. Lemos, A.N. Silveira and M. Dall'Aglio-Sobrinho, 2015. Ecological carrying capacity for intensive tilapia (*Oreochromis niloticus*) cage aquaculture in a large hydroelectrical reservoir in Southeastern Brazil. Aquacult. Eng., 66: 30-40.
45. Tsagaraki, T.M., G. Petihakis, K. Tsiaras, G. Triantafyllou and M. Tsapakis *et al.*, 2011. Beyond the cage: Ecosystem modelling for impact evaluation in aquaculture. Ecol. Model., 111: 2512-2523.
46. Asche, F., K.H. Roll and R. Tveteras, 2009. Economic inefficiency and environmental impact: An application to aquaculture production. J. Environ. Econ. Manage., 58: 93-105.
47. Yogev, U., K.R. Sowers, N. Mozes and A. Gross, 2017. Nitrogen and carbon balance in a novel near-zero water exchange saline recirculating aquaculture system. Aquaculture, 467: 118-126.
48. Besson, M., J. Aubin, H. Komen, M. Poelman and E. Quillet *et al.*, 2016. Environmental impacts of genetic improvement of growth rate and feed conversion ratio in fish farming under rearing density and nitrogen output limitations. J. Cleaner Prod., 116: 100-109.