

ISSN 1682-296X (Print)

ISSN 1682-2978 (Online)



Bio Technology



ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Chitinolytic and Microsclerostatic Activity of Iranian Strains of *Streptomyces plicatus* and *Frankia* sp. on Olive Isolate of *Verticillium dahliae*

G.H. Shahidi Bonjar and Sonia Aghighi

Department of Plant Pathology, College of Agricultural Sciences, Bahonar University of Kerman, Iran

Abstract: Among soil-borne fungi, cosmopolitan phytopathogen, *Verticillium dahliae* Klebahn is responsible for high yield losses in many plant species. Except for solarization in mediterranean countries and except in the few cases where disease-resistant cultivars are available, control of *Verticillium* wilt in commercial crops has been highly dependent on the application of preplant soil fumigants. Research to develop alternative control measurements should focus on biological approaches aimed at shifting the composition of soil microbial communities to suppress *Verticillium*. The merits of role of actinomycetes in biological control of soil-borne fungal-pathogens are known, however actinomycetes microflora of the Iranian soils has not been very well explored in searching for biofungicide agents. At the present research, *in vitro* studies of some biological effects of two Iranian strains of actinomycetes, *Streptomyces plicatus* strain 101 and *Frankia* sp. strain 103, are presented. Both strains revealed enzymatic activity and inhibited production of microsclerotia in *V. dahliae*. Treating the crude extract with chloroform, denaturated enzymatic activity of both strains. Thermal inactivation point of active phases of *S. plicatus* was 70 and 90°C and in *Frankia* sp. was determined as 60°C. Antifungal active phases of *S. plicatus* tolerate wide range of pH (5-13) but in *Frankia* sp. active phase tolerates pH 7-9. These two strains may be useful candidates for involving in integrated control programs of *Verticillium* vascular wilting.

Key words: *Verticillium dahliae*, Iranian actinomycetes microflora, biofungicide, antifungal, enzymatic activity, olive, thermal inactivation point, *Streptomyces plicatus*, *Frankia* sp.

INTRODUCTION

The genus *Verticillium* Nees (1817) represents one of the world's major pathogens, affecting crop plants mostly in the cool and warm temperate regions, but has also been reported from subtropical and tropical areas. *V. dahliae* is capable of surviving in field soil in several different ways. Various types of mycelia, clusters of hyaline cells and melanized microsclerotia have all been observed to germinate on assay plates used to determine the concentration of *V. dahliae* in naturally infested soils. These various forms of the fungus may be found in soil essentially free from host tissue or embedded in small-to-large fragments of tissue^[1]. Wide host range, long persistence in soil, propagation on rhizosphere of non-host plants and lack of effective control measurements highlights the need for research in evaluation of new methods for control of this pathogen^[1-3]. In future, biological control of soil-borne fungal or bacterial pathogens will be of increasing importance for a more sustainable agriculture. Furthermore, fungicides (biocides) such as methyl bromides will be phased out and thus potential alternatives are needed to control the soil-borne pathogen such as *V. dahliae*. This has prompted the

search for reliable antagonists which show a high degree of competitiveness and are active in the rhizospheres of different crops and in different soil types^[4,5]. Some workers have reported biological control of *Verticillium* by *Streptomyces* spp.^[6]. Chi and Hanson^[7] reported *in vitro* antifungal activity, fungistatic and fungicidal effects of *S. rimosus* against *V. dahliae* and *V. albo-atrum*. Ezrukh^[8] showed metabolites of actinomycetes from a cotton rhizosphere to interact with *V. dahliae*. Today, a commercial biocontrol agent Rhizovit® basis on spores of *Streptomyces rimosus* HR071 (= DSM 12424) is formulated in Germany. It is a phytoprotectant biofungicide and can be used to control several seed-borne and soil-borne fungal pathogens such as *Verticillium*, *Pythium* and *Rhizoctonia* both *in vitro* and *in vivo*^[9]. With extended environmental diversity, however, the actinomycetes microflora of the Iranian soils has not been very well explored with the goal of exploring new means of biocontrols. With the merits of their role in biological control of soil-borne fungal-pathogens, at the present research, *in vitro* studies of some biological effects of two Iranian strains of actinomycetes such as enzymatic activity, effect on melanin biosynthesis and microsclerotia formation on

olive isolate of *V. dahliae* with high pathogenicity in Iran are presented.

MATERIALS AND METHODS

Preparation of *V. dahliae* isolate: A registered phytopathogenic olive isolate of *V. dahliae* was used. It was obtained from Prof. Banihashemi, Mycology Laboratory of the Department of Plant Pathology, College of Agriculture, Shiraz University, Shiraz, Iran. The fungus was isolated from diseased olive trees with vascular wilting in Gorgan Province, Iran.

Culture media: Casein glycerol (or starch) agar (CGA) was used for supporting actinomycetes strains which composed of glycerol 10 g, casein 0.3 g, KNO₃ 2 g, NaCl 2 g, K₂HPO₄ 2 g, MgSO₄·7H₂O 0.05 g, CaCO₃ 0.02 g, FeSO₄·7H₂O 0.01 g and agar 18 g in 1 L of distilled H₂O (pH 7.2)^[10]. In submerged cultures, agar and glycerol was excluded and soluble starch was used (CG medium). Actinomycetes colonies were selected and transferred to CGA slants for further studies. All *V. dahliae* isolates were maintained on potato dextrose agar (PDA) (Difco).

Submerged culture of active strains and preparation of crude extract: The two active strains were grown in submerged cultures of CG medium on rotary shakers under 130 rpm at 30°C. To monitor the activity, aseptically small aliquots of culture media were taken every 24 h for 20 days and the activity was evaluated by well diffusion-method^[10,11]. When the activity reached maximum, the cultures were harvested; spores and mycelia were excluded by filtration through two layers of cheese cloth. The clarified sap was then dried to dark crude under reduced air at 30°C and kept refrigerated before use.

Classification of actinomycete: Actinomycete colonies were characterized morphologically and physiologically following the direction mentioned in the methods manual of international cooperative project for description and deposition of cultures of *Streptomyces* (ISP)^[12]. Identification procedures were done by Saadoun *et al.*^[13,14] Department of Biological Sciences, University of Science and Technology, Irbid, Jordan^[15].

In vitro antifungal bioassays

Agar disk-method: The two active Iranian actinomycete isolates reported previously^[16-18] were smeared on CGA medium as a single streak and after incubation at 28°C for 4-6 days, from well-grown streaks 6 mm agar disks of actinomycete colony masses were prepared by using sterile cork borers. Disks were then aseptically transferred to PDA plates having fresh lawn cultures of *V. dahliae*

isolates. Controls included using plain disks from CGA medium. Plates were incubated at 26°C for 4-6 days and fungicidal property indicative of chitinolytic activity^[19-21] and/or microsclerostatic activity were evaluated by measuring the diameter of inhibition (mm).

Well diffusion method: For evaluation of antifungal activity of aqueous samples, by use of sterile cork borer, wells (6x4 mm, 2 cm apart) were punctured in fresh lawn cultures or at 30 mm distance from plugs of *V. dahliae* isolates. Respective concentrations in dimethyl sulfoxide: methanol (1/1: v/v) solvent (DM solvent) were then administered to fullness in each well. Plates were incubated at 26°C for 4-6 days for lawn cultures and 14 days for dual culture disk-plugs. Plates were incubated at 26°C for 4-6 days and fungicidal property indicative of chitinolytic activity^[19, 21] and/or microsclerostatic activity were evaluated by measuring the diameter of inhibition zones (mm). Each experiment was repeated three times and the mean of inhibitory zones recorded. Controls included use of blank wells and use of DM solvent without test compounds.

Effect of heat and chloroform on bioactivity: To monitor the effect of temperature on bioactivity, small aliquots (10 mg ml⁻¹) of soluble crude were exposed to each of 30, 40, 50, 60, 70, 80 and 90°C for 10 min and cooled on ice afterwards^[22]. Bioactivity of treated samples was evaluated using well diffusion method. Control included incubation of an untreated sample at 26°C. For determination of sensitivity to chloroform, small aliquots of aqueous solutions from 7 day old submerged cultures were vigorously mixed with equal volumes of chloroform for 10 min, centrifuged at 4000 rpm for 10 min, phases were then separated and dried at 40°C under reduced air and placed in a desiccator overnight. All samples were tested by well diffusion method as described earlier.

Effect of pH on antifungal bioactivity: Effect of pH on activity and stability of activity was measured at different pH values by the general standard assay methods. The pH of the reaction mixtures was varied using the buffers described by Covington and Davison^[23]. The pH stability of the active crude sap was evaluated by incubating it for 0.5 h at various pH values at 30°C and evaluating them by agar well diffusion method.

RESULTS AND DISCUSSION

Cultures of active strains: Pure cultures of *Streptomyces plicatus* and *Frankia* sp. in slant CGA medium are indicated in Fig. 1. Streak cultures are indicated in Fig. 2. *In situ* morphology of aerial mycelia in slide cultures of *S. plicatus* under light microscope is indicated in Fig. 3.

Table 1: Bioactivity of *Streptomyces* and *Frankia* sp. on *Verticillium dahliae* determined by agar disk method in mm

Actinomycetes	CZ	MslZ	TI
<i>Streptomyces plicatus</i>	20	21	41
<i>Frankia</i> sp.	0	17	17

CZ: Clear zone of complete inhibition of fungal growth adjacent to tested agar disks of actinomycetes; MslZ: Microsclerostatic or microsclerotial inhibition zone, representing inhibitory effect on microsclerotia formation and development; TI: Total inhibition zone of fungal growth

Table 2: Identification criteria of *Frankia* sp. based on morphological and physiological characterizations

Morphological criteria		Physiological criteria	
ND	Spore chain	+	Arabinose
ND	Spore surface	+	Xylose
W	Aerial mass color	+	Inositol
1	Sporangia formation	+	Mannitol
0	Melanoid pigment	+	Fructose
0	Reverse side pigment	+	Rhamnose
1	Soluble pigment	+	Sucrose
		-	Raffinose

W: White; 1: Positive; 0: Negative; -: No utilization; +: Utilization. Low sporulation on Oatmeal agar medium, diameter of the hyphae is considerably larger than of *Streptomyces*, the mycelium (sporophore) ends to large club shaped sporangium

Table 3: Effect of chloroform on bioactivity of 7-days old submerged cultures and crude extracts of *Streptomyces plicatus* and *Frankia* sp. indicated by diameter of inhibition zones in mm

Actinomycetes	A	B	C	D
<i>Streptomyces plicatus</i>	30 (CZ+ MsS)	0	23 (MsS)	28 (CZ+ MsS)
<i>Frankia</i> sp.	20 (MsS)	0	0	18 (MsS)

A: Control, 7-days old aqueous submerged cultures; B: chloroform phase and aqueous phase © after treated with chloroform; D: Dried crude prepared from 7-days old aqueous submerged cultures. CZ: Clear zone of complete fungal growth inhibition; MsS: Microsclerostatic zone or microsclerotial inhibition zone




Fig. 1: Pure cultures of *Streptomyces plicatus* (Sp) and *Frankia* sp. (Fr) in slant CGA medium




Fig. 2: Streak cultures of *Streptomyces plicatus* (A) and *Frankia* sp. (B) on CGA medium



Fig. 3: *In situ* morphology of aerial mycelia of *Streptomyces plicatus* in slide cultures under light microscope (100 X)




Fig. 4: Clockwise from top, well method bioassay results of *Streptomyces plicatus*, *Frankia* sp. against *Verticillium dahliae* and blank well (control)




Fig. 5: Shake flasks of submerged cultures of *Streptomyces plicatus* (A) and *Frankia* sp. (B) in CG liquid culture media 7 days after inoculation

Bioassays: Well method bioassay results of *S. plicatus* and *Frankia* sp. are indicated in Fig. 4 and Table 1.

Submerged cultures and crude extract: Shake flasks containing submerged cultures of *S. plicatus* and




Fig. 6: Colonies of *Streptomyces plicatus* from 7 days old aqueous submerged cultures of shake flasks under binocular microscope (40 X)




Fig. 7: Temperature effect on antifungal bioactivity of crude extract of *Streptomyces plicatus* and *Frankia* sp.
Sp.: *Streptomyces plicatus*, Fr: *Frankia* sp., MsS: Microsclerostatic zone or microsclerotial inhibition zone




Fig. 8: Effect of pH on antifungal bioactivity of 7 days old aqueous submerged cultures of *Streptomyces plicatus* and *Frankia* sp.
Sp.: *Streptomyces plicatus*; Fr: *Frankia* sp.; MsS: Microsclerostatic zone or microsclerotial inhibition zone

Frankia sp. are indicated in Fig. 5. Colonies from submerged culture under binocular microscope are indicated in Fig. 6. The resulted crude extract composed of dark brown residue which was divided to small portions and preserved under refrigeration conditions at 4°C before use.

Classification of actinomycete: *Streptomyces plicatus* strain 101 was identified and reported as a new record from Iran previously^[6]. Table 2 shows the identification criteria of *Frankia* sp. based on morphological and physiological characterization.

Effect of heat and chloroform on activity: Temperature effect on antifungal activity of *S. plicatus* and *Frankia* sp. are indicated in Fig. 7. Effect of chloroform on bioactivity of both strains is indicated in Table 3.

Effect of pH on activity: Effect of pH on antifungal bioactivity of 7 days old aqueous submerged cultures of *Streptomyces plicatus* and *Frankia* sp. is indicated in Fig. 8.

Since there is high variation in the soil ecology of Iran, the population diversity of the actinomycetes in Iranian soils has to be further explored for new active isolates. Present findings represent the presence of potential antifungal metabolite(s) in both isolates against olive isolate of *V. dahliae*. *Streptomyces plicatus* prohibits mycelial growth and formation of melanin in the pathogen. It can be postulated that lack of melanin

renders the pathogen more susceptible to antagonists and promotes the conditions needed for its biological control in the rhizosphere. To enhance the tolerance and resistance of plants to fungal attack and lower or eliminate the use of environmentally unsafe fungicides, the genes for antifungal metabolites can be engineered into plants. It can be forecasted that "oncoming future" does not tolerate further pollution resulting from vast use of chemical pesticides. Hence, environmentally safe and non chemical-measures have to be developed to control plant maladies. In ideal biological control measures, proper micro organisms are those having well adaptation in soil and rhizosphere by exerting effective antagonistic activity against soil pathogens persistently. In this regard soil-driven actinomycetes do not have adverse effect or alter the biological balance of soils as chemical measures do. They should receive higher attention in research for biological controls worldwide. Conclusively, it may be assumed that the antifungal-metabolite gene from *Streptomyces plicatus* may be a useful candidate for genetic engineering of olive plants for increased tolerance against *V. dahliae*. Findings of this and similar researches may contribute for evolvement of resistant transgenic-plants with recombinant DNA having antifungal genes cloned from biologically active actinomycetes.

ACKNOWLEDGMENTS

Thanks to Head of Research Affairs Office of Bahonar University of Kerman for financial support of the project and helpful information and kind gift of *Verticillium dahliae* isolate by Prof. Banihashemi, Dept. of Plant Protection, College of Agriculture, Shiraz, Iran. We also would like to thank helpful works of Dr. Saadoun and his coworkers for identification of actinomycetes. This research is dedicated to Mr. A. Afzalipour, the founder of Bahonar University in Kerman.

REFERENCES

1. Schnathorst, W.C., 1981. Life Cycle and Epidemiology of *Verticillium*. In: Fungal Wilt Diseases of Plants. (Eds., Mace, M.E., A.A. Bell and C.H. Beckman), Academic Press: New York, pp: 81-111.
2. Schreiber, L.R. and R.J. Green Jr., 1963. Effect of root exudate on germination of conidia and microsclerotia of *Verticillium albo-atrum* inhibited by the soil fungistatic principle. *Phytopathology*, 53: 260-264.
3. Isaac, I., 1967. Speciation in *Verticillium*. *Annual Rev. Phytopathol.*, 5: 201-222.
4. Berg, G., S. Kurze, A. Buchner, E.M.H. Wellington and K. Smalla, 2000. Successful strategy for the selection of new strawberry associated rhizobacteria antagonistic to *Verticillium* wilt. *Canadian J. Microbiol.*, 46: 1128-1137.
5. Smalla, K., G. Wieland, A. Buchner, A. Zock, J. Parzy, S. Kaiser, N. Roskot, H. Heuer and G. Berg, 2001. Bulk and rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis: plant-dependent enrichment and seasonal shifts revealed. *Appl. Environ. Microbiol.*, 67: 4742-4751.
6. Dong, P., Q. Ji, T. Long and G. Ahmat, 1998. Preparation and application of biological control agents to control cotton wilts of *Fusarium* and *Verticillium*. Available on internet at: <http://www.bspp.org.uk/icpp98/5.2/2.html>.
7. Chi, C.C. and E.W. Hanson, 1965. *In vitro* effects of *Streptomyces rimosus* on some soil-inhabiting pathogenic fungi. *Plant Dis. Reporter*, 49: 159-163.
8. Ezrukh, E.N., 1978. Interaction between actinomycetes and fungi of the genus *Verticillium* Wallr. *Mycol. Phytopathol.*, 12: 443-445.
9. Marten, P., S. Bruckner, A. Minkwitz, P. Luth and G. Berg, 2001. Rhizovit®: Impact and Formulation of a New Bacterial Product. In: *Formulation of Microbial Inoculants. Cost Action 830/Microbial Inoculants for Agriculture and Environment*. (Eds., Koch, E. and P. Leinonen), pp: 78-82.
10. Dhingra, O.D. and J.B. Sinclair, 1995. *Basic Plant Pathology Methods*. CRC Press, USA, pp: 287-296, 390-391.
11. Acar, J.F. and F.W. Goldstein, 1996. Disk Susceptibility Test. In: *Antibiotics in Laboratory Medicine (4th Edn.)*, William and Wilkins Co. Baltimore, pp: 1-51.
12. Shirling, E.B. and D. Gottlieb, 1966. Methods for Characterization of *Streptomyces* sp. *Intl. J. Syst. Bacteriol.*, 16: 313-340.
13. Saadoun, I., F. Al-momani, H. Malkaawi and M.J. Mohammad, 1999. Isolation, identification and analysis of antibacterial activity of soil *Streptomyces* isolates from North Jordan. *Microbios*, 100: 41-46.
14. Saadoun, I. and R. Gharaibeh, 2002. The *Streptomyces* flora of Jordan and its potential as a source of antibiotics active against antibiotic-resistant gram-negative bacteria. *World J. Microbiol. Biotechnol.*, 18: 465-470.
15. Shahidi, B.G.H., M.H. Fooladi, M.J. Mahdavi and A. Shahghasi, 2004. Broadspectrum, a novel antibacterial from *Streptomyces* sp. *Biotechnology*, 3: 126-130.

16. Aghighi, S., G.H. Shahidi Bonjar and I. Saadoun, 2004. First report of antifungal properties of a new strain of *Streptomyces plicatus* (strain101) against four Iranian phytopathogenic isolates of *Verticillium dahliae*, a new horizon in biocontrol agents. *Biotechnology*, 3: 90-97.
17. Aghighi, S., G.H. Shahidi Bonjar, I. Saadoun, R. Rawashdeh and S. Batayneh, 2004. First report of antifungal spectra of activity of Iranian actinomycetes strains against *Alternaria solani*, *Alternaria alternata*, *Fusarium solani*, *Phytophthora megasperma*, *Verticillium dahliae* and *Saccharomyces cerevisiae*. *Asian J. Plant Sci.*, 3: 463-471.
18. Aghighi, S., L.S. Bonjar and G.H.S. Bonjar, 2004. *In vitro* anti-phytopathogenic and anti-yeast characterization of ten active strains of soil actinomycetes from Iran and their future prospects in biocontrol strategies. *Biotechnology*, (In Press).
19. Kong, L.D., D.D. Tzeng and C.H. Yang, 2001. Generation of PCR-based DNA fragments for specific detection of *Streptomyces saraceticus* N45. *Proceeding of National Science Council Roc (B)*, 25: 119-127.
20. Robbins, P.W., K. Overby, C. Albright, B. Benfield and J. Pero, 1992. Cloning and high-level expression of chitinase-encoding gene of *Streptomyces plicatus*. *Gene*, 111: 69-76.
21. Ni, X. and J. Westpheling, 1997. Direct repeat sequences in the *Streptomyces plicatus* chitinase-63 promoter direct both glucose repression and chitin induction. *Proceeding of National Academy of Science, USA.*, 94: 13116-13121.
22. Nawani, N.N. and B.P. Kapadnis, 2004. Production dynamics and characterization of the chitinolytic system of the *Streptomyces* sp. NK1057, a well equipped chitin degrader. *World J. Microbiol. Biotechnol.*, (In Press).
23. Covington, A.K. and W. Davison, 2000. Practical pH Measurements on Natural Waters. In: *Hand Book of Chemistry and Physics*. (Ed. Lide, D.R.), CRC Press, USA., pp: 841-845.