

ISSN 1682-296X (Print)

ISSN 1682-2978 (Online)



# Bio Technology



**ANSI***net*

Asian Network for Scientific Information  
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

## Plant Transformation: Needs and Futurity of the Transgenes

<sup>1,2,3</sup>Behrooz Darbani, <sup>4</sup>Safar Farajnia, <sup>2,3</sup>Shahin Noeparvar, <sup>5</sup>C. Neal Stewart Jr.,  
<sup>3</sup>Seyed A. Mohammadi and <sup>1</sup>Saeed Zakerbostanabad

<sup>1</sup>Department of Biotechnology, University of Rabe Rashidi, Tabriz, Iran

<sup>2</sup>Agricultural Biotechnology Research Institute for Northwest and West of Iran, Tabriz, Iran

<sup>3</sup>Department of Agronomy and Plant Breeding, Faculty of Agriculture,  
Tabriz University, Tabriz, Iran

<sup>4</sup>Biotechnology Research Center, Drug Applied Research Center,  
Tabriz University of Medical Science, Tabriz, Iran

<sup>5</sup>Department of Plant Sciences, University of Tennessee, Knoxville, TN, USA

---

**Abstract:** To produce transgenic plants which have various applications in agricultural and non-agricultural fields, a marker gene is necessary to recover a viable transgenic plant. To express or transcribe of transgenes, utilization of promoters is also unavoidable. Analysis of transgenes includes copy number, insertion site, integration stability, expression and its pattern and variability is immensely important in order to develop a successful transgenic event. This review presents the necessities for better recovery of transgenic plants, transcription or expression of transgenes, as well as methods to analyze transgenes.

**Key words:** Expression, integration site, marker gene, promoter, reporter gene

---

### INTRODUCTION

Presently, transgenic crops are increasingly becoming a dominant feature of the agricultural landscapes of developed countries. Since the introduction of Genetic Modified (GM) crops in 1996, the average growth in area has been more than 10% per year. This growth represents one of the highest adoption rates for any crop technology (Dunwell, 2004; James, 2005). Transgenic plants have also great potential impacts in non-agricultural fields, e.g., production of the medically useful products (pharmaceutical farming) or edible vaccine for oral virus immunization (Richter *et al.*, 2000). Introduction of desired genes into the plants consist of several steps. One of the steps of gene delivery to plant is the stably integration of the transgene into the host cell's genome in a manner that will result in proper expression of the gene. To establish an efficient expression from quantity and frequency points, it is unavoidable that up-stream promoting sequences are inserted exactly before the gene without any frame shift. Selection of a promoter depends on some factors, e.g., type of plant, target tissue(s), inducible or non-inducible expression, etc. Insertion region effects strongly the expression. In addition, in order to occur any gene knockout or disruption, it is very important to analysis of

the insertion region. Others factors such copy number and its variation also effect expression. Therefore, quantification of expression for few generations is indispensable. Expression pattern and time are the others which must be analyzed. Marker genes are essential to recover viable transgenic plants because the low transformation efficiency. Then, it must be inserted into the cassette or co-transformed. At present, different methods developed to deletion of the marker due to concerns about widespread occurrence in novel ecosystems, horizontal gene transfer or vertical gene transfer (Darbani *et al.*, 2007; Holger, 2003). Substitution of the new marker genes without any risk after spreading is the other way (Darbani *et al.*, 2007; Mentewab and Stewart, 2005).

In this study, we will discuss about promoters and markers as the necessities to produce transformed plants as well as different points of transgene analysis, e.g. copy number, expression and integration site.

### NEEDS FOR BETTER PLANT TRANSFORMATION

Theoretically, it is possible to produce transgenic plant using a coding region transformation. But it is not applicable due to the low chance of a proper integration

---

**Corresponding Authors:** <sup>1</sup>Behrooz Darbani and <sup>2</sup>Safar Farajnia, <sup>1</sup>Bagmishea, Shahriyar Street, Alley Ostad Gafari, No. 73, P. Box 5158643965, Tabriz, Iran  
<sup>2</sup>Biotechnology Research Center, Drug Applied Research Center, Tabriz University of Medical Science, Tabriz, Iran

(exactly after a promoter) and paucity of the transformation occurrences which caused to need a costly and time consuming process. However, these hindrances could be obviated via promoters and marker genes.

**Promoters:** Expression cassettes are comprised of three segments of DNA that includes up-stream regulatory region (promoters and enhancers), multiple cloning sites with the gene of interest and down-stream regulatory sequences. Up-stream and down-stream regions must be compatible with plants of interest. Therefore, efficient expression of foreign genes in transformed plants requires that they are placed under control of an active promoter in plant cells. Promoters are sequences that bind the RNA polymerase enzymes as cis-acting regulatory region, which is responsible for RNA transcription. Promoters can be generally divided into constitutive promoters, chemically or physically inducible promoters and tissue-specific or development-stage-specific promoters.

The *Cauliflower mosaic virus* (CaMV) 35S promoter is the most widely used constitutive promoter in dicots (Guilley *et al.*, 1982). The 35S promoter has a high level of transcription in nearly all plant tissues (Kay *et al.*, 1987). Other viral promoters have been characterized and cloned for use in transformation experiments (Mor *et al.*, 2003; Carrillo-Tripp *et al.*, 2006) such as the wheat dwarf virus (WDV) promoter. This promoter is particularly effective in monocots (Hofer *et al.*, 1992). Also, Subterranean Clover Stunt Virus (SCSV) promoter has been used either in dicots and monocots with similar pattern to those resulted with the actin (Act-1) or ubiquitin (Ubi) promoters (Petra *et al.*, 2003). Plant Ubi (Christensen and Quail, 1996), Act-1 (McElroy *et al.*, 1990, 1991) and alcohol dehydrogenase 1 (Adh-1) (Dennis *et al.*, 1984; Ellis *et al.*, 1987; Zhang and Wu, 1988) are also some of the monocot constitutive promoters that are widely used in plant expression cassettes.

Inducible expression is importance not only because of transgene properties, for example, if they are developmentally lethal, but it is often advantageous to obtain conditional expression to address research goals. Some inducible promoters include those induced by tetracycline, ecdysteroid agonist and glucocorticoid (Reynolds, 1999), as well as heat-shock promoters such as Gmhsp17.5-E gene promoter (Ainley and Key, 1990), wound based inducible promoters, stress induced promoters (Reynolds, 1999) and a light induced RUBISCO small subunit gene (rbcS) promoter (Schardl *et al.*, 1987).

Also, gene expression in a tissue-dependent manner and according to the developmental stage of the plant is achievable using tissue specific promoters such as root specific promoter of tobacco TobRB7 gene

(Yamamoto *et al.*, 1991), tuber specific promoter of potato patatin B33 gene (Rocha-Sosa *et al.*, 1989), flower specific promoter of the petunia chsA gene (Koes *et al.*, 1989, 1990), endosperm-specific promoter from multigene family of monocot glutenin subunits (Roberts *et al.*, 1989) and seed specific napin promoter of *Brassica napus* (DeLisle and Crouch, 1989; Stayton *et al.*, 1991; Stalberg *et al.*, 1993). In addition, it was demonstrated the ability of T7 promoter to promote expression in plants even with tissue specificity and inducibility expression which it can be useful in both dicot and monocot plants (Nguyen *et al.*, 2004; Dunn *et al.*, 1988; Gatz *et al.*, 1992).

**Marker genes:** A selectable marker gene is used to recover transformants which permits the preferential growth of transformed cells. Also, the stable introduction of genes into plants through genetic engineering normally necessitates the use of a selectable marker, especially when the transformation frequency is low,  $10^{-3}$  to  $10^{-6}$  (Flavell *et al.*, 1992).

Several types of selectable marker genes exist, which could be separated into five groups including antibiotic-resistance, herbicide-resistance and metabolic pathway-based and positive and negative selectable marker genes. Some of well-known antibiotic selectable marker genes are: neomycin phosphotransferase gene (nptII) from *Escherichia coli* transposon Tn5 as the first used selectable marker for plant transformation (Herrera-Estrella *et al.*, 1983; Bevan *et al.*, 1983), hygromycin phosphotransferase gene (hpt or aphIV) from *E. coli* (Waldron *et al.*, 1985; Van den Elzen *et al.*, 1985; Lulsdorf *et al.*, 1991; Walters *et al.*, 1992), streptomycin resistance gene (spt) from transposon Tn5 (Jones *et al.*, 1987), aminoglycoside-3'-adenyltransferase (aadA) (Svab *et al.*, 1990), dihydrofolate reductase (dhfr) mouse gene (Hauptmann *et al.*, 1988; Eichholtz *et al.*, 1987) and bacterial kanamycin-resistant nptII gene and an *Arabidopsis thaliana* ATP Binding Cassette (ABC) transporter (Atwbc19) gene (Mentewab and Stewart, 2005). It is worth mentioning that while most marker genes are from bacterial origin, there is a renewed emphasis on using marker genes from plants, which might have higher biosafety. Herbicide resistance markers include a bialaphos resistance gene (bar) from *Streptomyces hygroscopicus* (Thompson *et al.*, 1987; Somers *et al.*, 1992; Gordon-Kamm *et al.*, 1990; Schroeder *et al.*, 1993), chlorsulfuron resistance gene (csr1-1) from *Arabidopsis* (Haughn *et al.*, 1988; Fromm *et al.*, 1990), sulfonamide resistance gene from plasmid R46 (Guerineau and Mullineaux, 1989) and a mutant petunia epsps gene (Shah *et al.*, 1986).

There still remain biosafety concerns about marker genes in transgenic plants. Application of the metabolic pathway based markers, such as bacterial dihydrodipicolinate synthase (dhps) (Perl *et al.*, 1993) as aspartate family metabolic pathway based marker, betaine aldehyde dehydrogenase (badh) gene from spinach (Daniell *et al.*, 2001), *E. coli* derived phosphomannose isomerase (PMI) (Wright *et al.*, 2001), xylose isomerase (xylA) gene of *Streptomyces rubiginosus* (Jaiwal *et al.*, 2002) and trehalose-6-phosphate synthase, AtTPS1, (Leyman *et al.*, 2004), without any concerns about horizontal or vertical gene transfer is one of the ways to reduce the marker gene effects from horizontal or vertical transfer points. By the way, insertion of any additional sequence into the host genome is unfavorable. Therefore, expelling of the marker gene could be the ideal which applicable via different approaches, e.g., co-transformation especially using P-DNA, site-specific recombinase-mediated marker deletion (e.g., Cre/loxP, FLP/FRT and R/Rs site-specific recombination systems), transposon-based expelling systems (e.g., Ac transposon) and intrachromosomal recombination based excision (Holger, 2003; Scutt *et al.*, 2002; Darbani *et al.*, 2007).

Cyanamide hydratase (Cah) gene (Weeks, 2000) as positive selectable markers and nitrate reductase gene (nia) (Nussaume *et al.*, 1991), cytosine deaminase (codA) (Perera *et al.*, 1993) and human herpes simplex virus thymidine kinase type 1 gene (HSVtk) (Czako and Marton, 1994) as negative selectable marker genes and dao1, encoding D-amino acid oxidase as either positive and/or negative marker gene (Erikson *et al.*, 2004) are some of the other markers.

The mentioned segregation or expelling based marker gene deletion approaches could be more efficient using a positive-negative selection system (Holger, 2003; Scutt *et al.*, 2002; Darbani *et al.*, 2007).

### TRANSGENE ANALYSIS

The development of efficient crop transformation systems has necessitated the development of efficient methods for detailed molecular characterization of putative transgenic events. Transgene analysis has different aspects such as transgene quantification, position and expression assays will be discussed in this section.

**Copy number and integration site:** Standard Polymerase Chain Reaction (PCR) methods could be utilized to detect the presence of foreign DNA in transformed plants, but this presence/absence assay is problematic because of the chance of trace DNA contamination or residual

*Agrobacterium*. It is more useful to determine of the number of inserted copies of transgenes. Traditionally, researchers have applied nucleic acid blotting techniques to provide an estimate of transgene copy number and organization (Southern, 1975; Sabelli and Shewry, 1991, 1995a). Although Southern blot analysis is the gold-standard for determining loci number and intactness of the insert, this method is unwieldy to apply in high-throughput screening of putative transformants. Recently, the development of quantitative PCR (Q-PCR) for determining transgene copy number has overcome the limitations of standard PCR and Southern blot analysis (Ingham *et al.*, 2001; Mason *et al.*, 2002) which is high-throughput approach using small amounts of sample with capacity of selection as soon as at tissue culture phase, is faster and less expensive than southern blot analysis (Mason *et al.*, 2002). It is important to use proper technical and statistical techniques for robust estimations of transgene copy number.

Chromosomal site of transgene can be detected via radioactive based *in situ* hybridization (Gall and Pardue, 1969) or via fluorescence *in situ* hybridization (FISH) methods (Ten Hoopen *et al.*, 1996; Snowdon *et al.*, 2001; Choi *et al.*, 2002; Bourdon *et al.*, 2002). In order to precisely studying of integration site of transgenes, thermal asymmetric interlaced polymerase chain reaction (TAIL-PCR) is usable. TAIL-PCR and high-throughput TAILPCR (HT-TAILPCR) are fast and efficient methods to amplify unknown sequences adjacent to known insertion sites (Liu and Whittier, 1995; Budziszewski *et al.*, 2001; Parinov *et al.*, 1999; Tissier *et al.*, 1999; Singer and Burke, 2003).

**Variation of transgene expression:** Various factors are thought to influence transgene expression in plants, including transgene copy number, transgene insertion site and RNA silencing in the host plant. Also, regulatory sequences could be effect on expression level and its variation. For example the mannopine synthase promoter yields less variable but also lower transgene expression than that of 35S (De Bolle *et al.*, 2003). It is determined that the transgene copy number may depends on the transformation method (Butaye *et al.*, 2005). Gene targeting into a predefined chromosomal site could be achieved by homologous recombination. To date, however, the moss *Physcomitrella patens* is the only (lower) plant that has shown a high rate of homologous recombination. Integration of transgenic DNA through homologous recombination is so infrequent in higher eukaryotes and therefore, targeting transgenes to a specific position is nearly infeasible (Kumar and Fladung, 2001). A more promising method to generate single-copy

transgenic plants is the recombinase-mediated resolution of multiple insertions, which allows site-specific integration of the transgene (Ow, 2002). However, this method has only been partially successful and efficiency is poor. Also, the implementation of Matrix Attachment Regions (MARs) in transgene constructs might be considered when low variation of transgene expression is desired (Allen *et al.*, 2000). MARs can increase the expression of transgenes in whole organisms or cells in culture and is necessary for the proper formation, maintenance, or regulation of chromatin structure. MARs have been isolated from diverse range of eukaryotes, including mammals, birds, insects and plants varying from 50-200 kb in length (Schöffl *et al.*, 1993; Allen *et al.*, 1993, 1996; Vain *et al.*, 1999; Chinn and Comai, 1996; Mlynarova *et al.*, 1994; Liu and Tabe, 1998; Van der Geest *et al.*, 1994; Holmes-Davis and Comai, 1998). Experimental data indicated that MARs enhance transgene expression from 2- to 60-fold in different plant systems (Holmes-Davis and Comai, 1998; Allen *et al.*, 2000; James *et al.*, 2002; Nowak *et al.*, 2001; Ülker *et al.*, 1999; Halweg *et al.*, 2005). It has been recently shown that the RB7 MAR, a 1.2 kb fragment isolated from the 3' flanking region of the RB7 root-specific gene of tobacco, caused a significant increase in the activity of highly active promoters (CaMV 35S, nopaline synthase promoter and octopine synthase promoter) as compared to controls. However, the presence of RB7 MAR did not significantly increase the activity of weak promoters. Importantly, most transgenes flanked by RB7 MAR showed a large reduction in the number of low-expressing transformants relative to control constructs without MARs, suggesting that these elements can reduce the frequency of gene silencing in primary transformants (Mankin *et al.*, 2003; Halfhill *et al.*, 2003). Also, MARs play a role in reducing variability of that expression between independent transformants (Liu and Tabe, 1998; Mlynarova *et al.*, 1995, 1996).

**Expression assays:** The study of transgene expression is of vital importance whenever transgenic plants are produced. Many techniques exist to analyze of transgene expression such as western blot analysis and Enzyme-Linked Immunosorbent Assay (ELISA) at the protein level (Towbin *et al.*, 1979; Burnette, 1981; Yalow and Berson, 1959; Engvall and Perlman, 1971). *In situ* hybridization is a method to assay patterns of protein expression (Angerer *et al.*, 1987; Long and Rebagliati, 2002). Gene expression can also be assayed at the transcript level. Several methods include northern blotting (Alwine *et al.*, 1977; Kumar and Lindberg, 1972; Sabelli and Shewry, 1995b), nuclear run-on (or run-off) transcription assays

(Kafatos, 1972; Ganguly and Skoultchi, 1985; Gatehouse and Thompson, 1995), RNase A/T1 mapping (Brown and Waugh, 1989; Simpson and Brown, 1995a), primer extension assays (Hershberger and Culp, 1990; Simpson and Brown, 1995b), reverse transcriptase-polymerase chain reaction (RT-PCR) and quantitative or real-time RT-PCR (Robinson and Simon, 1991; Ferre, 1992; Freeman *et al.*, 1999; Gause and Adamovicz, 1994; Page and Minocha, 2004) have been developed for expression quantification at RNA level which real-time RT-PCR allows a highly sensitive quantification of transcriptional levels of the gene of interest in a few hours with minimal handling of the samples (Kammula *et al.*, 2000; Mandigers *et al.*, 1998; Tichopad *et al.*, 2003; Valasek and Repa, 2005). RT-PCR can detect low-abundant mRNAs, but it cannot quantify them, which is achievable using real-time RT-PCR (Deprez *et al.*, 2002; Bustin, 2000). Some advantages of real-time PCR quantification include a large dynamic range, tremendous sensitivity, highly sequence-specificity, minimum post-amplification processing, which make it a good technique also for large scale screening (Wong and Medrano, 2005).

**Reporter genes:** Many studies on plant promoters and on the regulation of gene expression have been made possible using reporter genes as well as monitoring transgenes under field conditions. Reporter genes allows for direct visual detection of transformed tissue, thus avoiding the need for more time-consuming biochemical characterization. Widely used reporter genes include the B-glucuronidase gene (*uidA* or *gus*), which originates from *E. coli* (Jefferson *et al.*, 1986), firefly luciferase gene (*luc*) (De Wet *et al.*, 1985; Ow *et al.*, 1986; Mullineaux *et al.*, 1990; Riggs and Chrispeels, 1987; Bolivar, 1978; Alton and Vapnek, 1979), chloramphenicol acetyltransferase gene (*cat*) originates from transposon Tn9 (Alton and Vapnek, 1979; Davis *et al.*, 1992), *lacZ* gene encoding B-galactosidase (B-GAL) in *E. coli* (Alton and Vapnek, 1979) and the green fluorescent protein (GFP) gene from jellyfish (Prasher *et al.*, 1992). GFP is especially useful as an *in vivo* whole-plant marker for field-level applications (Stewart, 1996; Richards *et al.*, 2003). Also, the R genes of *Zea mays* are useful as a scorable marker in mature and differentiated cells which stimulate endogenous anthocyanin accumulation in the vacuoles of plant tissues (Kloti *et al.*, 1993). There are additional fluorescent protein markers that have recently been described that will enable multi-color labeling in transgenic plants (Stewart, 2006).

Most researchers are familiar with GUS assays that have good sensitivity that an enzymatic-based system can deliver. GUS requires, however, the destructive

harvest of the tissues to be assayed. Therefore, researchers have increasingly adopted GFP and other fluorescent proteins to in vivo image and quantify gene expression in plants (Stewart, 2001, 2006).

### CONCLUSIONS

Even though plant transformation can no longer be described as a new technology (it is over 20 years-old), it continues to evolve. In addition, tools such as marker genes continue to expand at an astonishing rate, leading to more efficient transformation. Old tools are also applied to new plants, which opens up the number of traits for crop improvement. Despite controversies, worldwide adoption of transgenic plants is expanding every year. Technological advances, such as those provided by plant biotechnology are needed to better feed the world's people.

### REFERENCES

- Ainley, W.M. and J.L. Key, 1990. Development of a heat shock inducible expression cassette for plants characterization of parameters for its use in transient expression assays. *Plant Mol. Biol.*, 14 (6): 949-967.
- Allen, G.C., G.E. Hall, L.C. Childs, A.K. Weissinger, S. Spiker and W.F. Thompson, 1993. Scaffold attachment regions increase reporter gene expression in stably transformed plant cells. *Plant Cell*, 5 (6): 603-613.
- Allen, G.C., G. Hall, S. Michalowski, W. Newman, S. Spiker, A.K. Weissinger and W.F. Thompson, 1996. High-level transgene expression in plant cells: Effects of a strong scaffold attachment region from tobacco. *Plant Cell*, 8 (55): 899-913.
- Allen, G., S. Spiker and W. Thompson, 2000. Use of Matrix Attachment Regions (MARs) to minimize transgene silencing. *Plant Mol. Biol.*, 43 (2-3): 361-376.
- Alton, N.K. and D. Vapnek, 1979. Nucleotide sequence analysis of the chloramphenicol resistance transposon *tn9*. *Nature*, 282 (5741): 864-869.
- Alwine, J.C., D.J. Kemp and G.R. Stark, 1977. Method for detection of specific RNAs in agarose gels by transfer to diazobenzyloxymethyl-paper and hybridization with DNA probes. *Proc. Natl. Acad. Sci.*, 74 (12): 5350-5354.
- Angerer, L.M., K.H. Cox and R.C. Angerer, 1987. Demonstration of tissue specific gene expression by in situ hybridization. *Meth. Enzymol.*, 152: 649-661.
- Bevan, M.W., R.B. Flavell and M.D. Chilton, 1983. A chimeric antibiotic resistance gene as a selectable marker for plant cell transformation. *Nature*, 304 (5897): 184-187.
- Bolivar, F., 1978. Construction and characterization of new cloning vehicles. III. Derivatives of plasmid pBR322 carrying unique EcoRI sites for selection of EcoRI generated recombinant DNA molecules. *Gene*, 4 (2): 121-136.
- Bourdon, V., Z. Ladbroke, A. Wickham, D. Lonsdale and W.A. Harwood, 2002. Homozygous transgenic wheat plants with increased luciferase activity do not maintain their high level of expression in the next generation. *Plant Sci.*, 163 (2): 297-305.
- Brown, J.W.S. and R. Waugh, 1989. Maize U2 snRNAs gene sequence and expression. *Nucleic Acids Res.*, 17 (22): 8991-9001.
- Budziszewski, G.J., S.P. Lewis, L.W. Glover, J. Reineke, G. Jones, L.S. Ziemnik, J. Lonowski, B. Nyfeler, G. Aux, Q. Zhou, J. McElver, D.A. Patton, R. Martienssen, U. Grossniklaus, H. Ma, M. Law and J.Z. Levin, 2001. Arabidopsis genes essential for seedling viability: Isolation of insertional mutants and molecular cloning. *Genetics*, 159 (4): 1765-1778.
- Burnette, W.N., 1981. Western blotting, electrophoretic transfer of proteins from sodium dodecylsulphate-polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein. *A. Anal. Biochem.*, 112 (2): 195-203.
- Bustin, S.A., 2000. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J. Mol. Endocrinol.*, 25 (2): 169-193.
- Butaye, K., B. Cammue, S. Delaure and M. De Bolle, 2005. Approaches to minimize variation of transgene expression in plants. *Mol. Breed.*, 16 (1): 79-91.
- Carrillo-Tripp, J., H. Shimada-Beltran and R. Rivera-Bustamante, 2006. Use of geminiviral vectors for functional genomics. *Curr. Opin. Plant Biol.*, 9 (2): 209-215.
- Chinn, A.M. and L. Comai, 1996. The heat shock cognate 80 gene of tomato is flanked by matrix attachment regions. *Plant Mol. Biol.*, 32 (5): 959-968.
- Choi, H.W., P.G. Lemaux and M.J. Cho, 2002. Use of fluorescence *in situ* hybridisation for gross mapping of transgenes and screening for homozygous plants in transgenic barley (*Hordeum vulgare* L.). *Theor. Applied Genet.*, 106 (1): 92-100.
- Christensen, A.H. and P.H. Quail, 1996. Ubiquitin promoter-based vectors for high-level expression of selectable and/or screenable marker genes in monocotyledonous plants. *Transgenic Res.*, 5 (3): 213-218.
- Czako, M. and L. Marton, 1994. The herpes simplex virus thymidine kinase gene as a conditional negative-selection marker gene in *Arabidopsis thaliana*. *Plant Physiol.*, 104 (3): 1067-1071.

- Daniell, H., B. Muthukumar and S.B. Lee, 2001. Marker free transgenic plants: Engineering the chloroplast genome without the use of antibiotic selection. *Curr. Genet.*, 39 (2): 109-116.
- Darbani, B., A. Eimanifar, C.N. Stewart and W.N. Camargo, 2007. Methods to produce marker-free transgenic plants. *Biotechnol. J.*, 2 (1): 83-90.
- Davis, A.S., M.R. Davey, R.C. Clothier and E.C. Cocking, 1992. Quantification and comparison of chloramphenicol acetyltransferase activity in transformed plant protoplasts using high-performance liquid chromatography and radioisotope based assays. *Anal. Biochem.*, 201 (1): 87-93.
- De Bolle, M., K. Butaye, W. Coucke, I. Goderis, P. Wouters, N. Van Boxel, W. Broekaert and B. Cammue, 2003. Analysis of the influence of promoter elements and a matrix attachment region on the inter-individual variation of transgene expression in populations of *Arabidopsis thaliana*. *Plant Sci.*, 165 (1): 169-179.
- De Lisle, A.J. and M.L. Crouch, 1989. Seed storage protein transcription and mRNA levels in *Brassica napus* during development and in response to exogenous abscisic acid. *Plant Physiol.*, 91 (2): 617-623.
- De Wet, J.R., K.V. Wood, D.R. Helinski and M. DeLuca, 1985. Cloning of firefly luciferase cDNA and the expression of active luciferase in *Escherichia coli*. *Proc. Natl. Acad. Sci.*, 82 (23): 7870-7873.
- Dennis, E.S., W.L. Gerlach, A.J. Pryor, J.L. Bernetzen, A. Inglis, D. Llewellyn, M.M. Sachs, R.J. Ferl and W.J. Peacock, 1984. Molecular analysis of the alcohol dehydrogenase (Adh1) gene of maize. *Nucleic Acids Res.*, 12 (9): 3983-4000.
- Deprez, R.H.L., A.C. Fijnvandraat, J.M. Ruijter and A.F.M. Moorman, 2002. Sensitivity and accuracy of quantitative real-time polymerase chain reaction using SYBR green I depends on cDNA synthesis conditions. *Anal. Biochem.*, 307 (1): 63-69.
- Dunn, J.J., B. Kripl, K.E. Bernstein, H. Westphal and F.W. Studier, 1988. Targeting bacteriophage T7 RNA polymerase to the mammalian cell nucleus. *Gene*, 68 (2): 259-266.
- Dunwell, J.M., 2004. Transgenic Crops: The Current and Next Generations. In: *Methods in Molecular Biology. Transgenic Plants: Methods and Protocols*, Peña, L. (Ed.). Vol. 286, Humana Press Inc: Totowa, NJ, pp: 377-397.
- Eichholtz, D.A., S.G. Rogers, R.B. Horsch, H.J. Klee, M. Hayford, N.L. Hoffmann, S.B. Bradford, C. Fink, J. Flick, K.M. O'Connell and R.T. Fraley, 1987. Expression of mouse dihydrofolate reductase gene confers methotrexate resistance in transgenic petunia plants. *Somatic Cell Mol. Genet.*, 13 (1): 67-76.
- Ellis, J.G., D.J. Llewellyn, E.S. Dennis and W.J. Peacock, 1987. Maize Adh-1 promoter sequences control anaerobic regulation: Addition of upstream promoter elements from constitutive genes is necessary for expression in tobacco. *EMBO. J.*, 6 (1): 11-16.
- Engvall, E. and P. Perlman, 1971. Enzyme-Linked Immunosorbent Assay (ELISA) quantitative assay of immunoglobulin G. *Immunochemistry*, 8 (9): 871-874.
- Erikson, O., M. Hertzberg and T. Nasholm, 2004. A conditional marker gene allowing both positive and negative selection in plants. *Nat. Biotechnol.*, 22 (4): 455-458.
- Ferre, I., 1992. Quantitative or semi-quantitative PCR: Reality versus myth PCR. *Meth. Applied*, 2 (1): 1-9.
- Flavell, R.B., E. Dart, R.L. Fuchs and R.T. Fraley, 1992. Selectable marker genes: Safe for plants? *Biotechnology*, 10 (2): 141-144.
- Freeman, W.M., S.J. Walker and K.E. Vrana, 1999. Quantitative RT-PCR: Pitfalls and potential. *Biotechniques*, 26 (1): 112-125.
- Fromm, M.E., F. Morrish, C. Armstrong, R. Williams, J. Thomas and T.M. Klein, 1990. Inheritance and expression of chimeric genes in the progeny of transgenic maize plants. *Biotechnology*, 8 (9): 833-839.
- Gall, J. and M.L. Pardue, 1969. Formation and detection of RNA-DNA hybrid molecules in cytological preparations. *Proc. Natl. Acad. Sci.*, 63 (3): 378-383.
- Ganguly, S. and A.I. Skoultchi, 1985. Absolute rates of globin gene transcription and mRNA formation during differentiation of cultured mouse erythroleukaemia cells. *J. Biol. Chem.*, 260 (12): 167-173.
- Gatehouse, J. and A.J. Thompson, 1995. Nuclear Run-on Transcription Assays. In: *Methods in Molecular Biology*, Jones, H. (Ed.). Vol. 49, Gene Transfer and Expression Protocols. Humana Press Inc., Totowa, NJ. pp: 229-238.
- Gatz, C., C. Froberg and R. Wendenburg, 1992. Stringent repression and homogeneous de-repression by tetracycline of a modified CaMV 35S promoter in intact transgenic tobacco plants. *Plant J.*, 2 (3): 397-404.
- Gause, W.C. and J. Adamovicz, 1994. The use of PCR to quantitate gene-expression. *PCR Methods Appl.*, 3 (6): S123-S135.
- Gordon-Kamm, W.J., T.M. Spencer, M.L. Mangano, T.R. Adams, R.J. Daines, W.G. Start, J.V. O'Brien, S.A. Chambers, W.R. Adams, N.G. Willetts, T.B. Rice, C.J. Mackey, R.W. Krueger, A.P. Kausch and P.G. Lemaux, 1990. Transformation of maize cells and regeneration of fertile transgenic plants. *Plant Cell*, 2 (7): 603-618.

- Guerineau, F. and P. Mullineaux, 1989. Nucleotide sequence of the sulfonamide resistance gene from plasmid R46. *Nucleic Acids Res.*, 17 (11): 43-70.
- Guilley, H., R.K. Dudley, G. Jonard, E. Balazs and K.E. Richards, 1982. Transcription of cauliflower mosaic virus DNA: Detection of promoter sequences and characterization of transcription. *Cell*, 30 (3): 763-773.
- Halfhill, M.D., R.J. Millwood, A.K. Weissinger, S.L. Warwick and C.N. Stewart, 2003. Additive transgene expression and genetic introgression in multiple green-fluorescent protein transgenic crop-weed hybrid generations. *Theor. Applied Genet.*, 107 (8): 1533-1540.
- Halweg, C., W.F. Thompson and W. Spikera, 2005. The Rb7 matrix attachment region increases the likelihood and magnitude of transgene expression in tobacco cells: A flow cytometric study. *Plant Cell*, 17 (2): 418-429.
- Hauptmann, R.M., V. Vasil, P. Ozias-Akins, Z. Tabaeizadeh, S.G. Rogers, R.T. Fraley, R.B. Horsch and I.K. Vasil, 1988. Evaluation of selectable markers for obtaining stable transformants in the Gramineae. *Plant Physiol.*, 86 (2): 602-606.
- Haughn, G.W., J. Smith, B. Mazur and C. Somerville, 1988. Transformation with a mutant *Arabidopsis* acetolactate synthase gene renders tobacco resistant to sulfonylurea herbicides. *Mol. Gen. Genet.*, 211 (2): 266-271.
- Herrera-Estrella, L., M. De Block, E. Messens, J.P. Hernalsteens, M. Van Montagu and J. Schell, 1983. Chimeric genes as dominant selectable markers in plant cells. *EMBO. J.*, 2: 2987-2995.
- Hershberger, R.P. and L.A. Culp, 1990. Cell-type-specific expression of alternatively spliced human fibronectin III CS mRNAs. *Mol. Cell Biol.*, 10 (2): 662-671.
- Hofer, J.M.I., E.L. Dekker, H.V. Reynolds, C.J. Woolston, B.S. Cox and P.M. Mullineaux, 1992. Coordinate regulation of replication and virion sense gene expression in wheat dwarf virus. *Plant Cell*, 4 (2): 213-223.
- Holger, P., 2003. Marker-free transgenic plants. *Plant Cell Tissue Organ Cult.*, 74 (2): 123-134.
- Holmes-Davis, R. and L. Comai, 1998. Nuclear matrix attachment regions and plant gene expression. *Trends Plant Sci.*, 3 (3): 91-97.
- Ingham, D.J., S. Beer, S. Money and G. Hansen, 2001. Quantitative real-time PCR assay for determining transgene copy number in transformed plants. *Biotechniques*, 31 (1): 132-140.
- Jaiwal, P.K., L. Sahoo, N.D. Singh and R.P. Singh, 2002. Strategies to deal with the concern about marker genes in transgenic plants: Some environment friendly approaches. *Curr. Sci.*, 83 (2): 128-136.
- James, C., 2005. Executive Summary of Global Status of Commercialized Biotech/GM Crops, SAAA Briefs No. 34. ISAAA: Ithaca, N.Y.
- James, V.A., C. Avart, B. Worland, J.W. Snape and P. Vain, 2002. The relationship between homozygous and hemizygous transgene expression levels over generations in populations of transgenic rice. *Theor. Applied Genet.*, 104 (4): 553-561.
- Jefferson, R.A., S.M. Burgess and D. Hush, 1986. P-Glucuronidase from *Escherichia coli* as a gene-fusion marker. *Proc. Natl. Acad. Sci.*, 83 (22): 8447-8451.
- Jones, J.D.G., Z. Svab, E.C. Harper, C.D. Hurwitz and P. Maliga, 1987. A dominant nuclear streptomycin resistance marker for plant cell transformation. *Mol. Gen. Genet.*, 210 (1): 86-91.
- Kafatos, F.C., 1972. mRNA stability and cellular differentiation. *Acta Endocrinol. Suppl. (Copenh.)*, 168: 319-345.
- Kammula, U.S., F.M. Marincola and S.A. Rosenberg, 2000. Real-time quantitative polymerase chain reaction assessment of immune reactivity in melanoma patients after tumor peptide vaccination. *J. Natl. Cancer Inst.*, 92 (16): 1336-1344.
- Kay, R., A. Chan, M. Daly and J. McPherson, 1987. Duplication of CaMV 35S promoter sequences creates a strong enhancer for plant genes. *Science*, 236 (4806): 1299-1302.
- Kloti, A., V.A. Iglesias, J. Wijnm, P.K. Burkhardt, S.K. Datta and I. Potrykus, 1993. Gene transfer by electroporation into intact scutellum cells of wheat embryos. *Plant Cell Rep.*, 12 (12): 671-675.
- Koes, R.E., C.E. Spelt and J.N.M. Mel, 1989. The chalcone synthase multigene family, of *Petunia hybrida* (V30): Differential, light-regulated expression during flower development and UV light induction. *Plant Mol. Biol.*, 12 (2): 213-225.
- Koes, R.E., R. Van Blokland, F. Quattrocchio, A.J. Van Tunen and J. Mol, 1990. Chalcone synthase promoters in petunia are active in pigmented and unpigmented cell types. *Plant Cell*, 2 (5): 379-392.
- Kumar, A. and U. Lindberg, 1972. Characterization of messenger ribonucleoprotein and messenger RNA from KB cells. *Proc. Natl. Acad. Sci.*, 69 (3): 681-685.
- Kumar, S. and M. Fladung, 2001. Controlling transgene integration in plants. *Trends Plant Sci.*, 6 (4): 155-159.



- Leyman, B., N. Avonce, M. Ramon, P. Van, Dijk, J.M. Thevelein and G. Iturriaga, 2004. New Selection Marker for Plant Transformation. In: Methods in Molecular Biology, Reviews and Protocols, Balbas, P. and A. Lorence (Eds.). Vol. 267, Humana Press Inc: Totowa, NJ. pp: 385-396.
- Liu, Y.G. and R.F. Whittier, 1995. Thermal asymmetric interlaced PCR: Automatable amplification and sequencing of insert end fragments from P1 and YAC clones for chromosome walking. *Genomics*, 25 (3): 674-681.
- Liu, J.W. and L.M. Tabe, 1998. The influences of two plant nuclear Matrix Attachment Regions (MARs) on gene expression in transgenic plants. *Plant Cell Physiol.*, 39 (1): 115-123.
- Long, S. and M. Rebagliati, 2002. Sensitive two-color whole-mount *in situ* hybridizations using digoxigenin- and dinitrophenol-labeled RNA probes. *Biotechniques*, 32 (3): 494-500.
- Lulsdorf, M.M., H. Rempel, J.A. Jackson, D.S. Baliski and S.L.A. Hobbs, 1991. Optimizing the production of transformed pea (*Pisum sativum* L.) callus using disarmed *Agrobacterium tumefaciens* strains. *Plant Cell Rep.*, 9 (9): 479-483.
- Mandigers, C., J. Meijerink and J. Raemaekers, 1998. Graft versus lymphoma effect of donor leucocyte infusion shown by real time quantitative PCR analysis of (14;18). *Lancet*, 352 (9139): 1522-1523.
- Mankin, S.L., G.C. Allen, T. Phelan, S. Spiker and W.F. Thompson, 2003. Elevation of transgene expression level by flanking Matrix Attachment Regions (MAR) is promoter dependent: A study of the interactions of six promoters with the RB7 3' MAR. *Transgenic Res.*, 12 (1): 3-12.
- Mason, G., P. Provero, A.M. Vaira and G.P. Accotto, 2002. Estimating the number of integrations in transformed plants by quantitative real-time PCR. *BMC Biotechnol.*, 2 (1): 20.
- McElroy, D., E. Zhang, J. Cao and R. Wu, 1990. Isolation of an efficient actin promoter for use in rice transformation. *Plant Cell*, 2 (2): 163-171.
- McElroy, D., A.D. Blowers, B. Jenes and R. Wu, 1991. Construction of expression vectors based on the rice actin 1 (Act1) 5' region for use in monocot transformation. *Mol. Gen. Genet.*, 231 (1): 150-160.
- Mentewab, A. and C.N. Stewart, 2005. Over expression of an arabidopsis thaliana abc transporter confers kanamycin resistance to transgenic plants. *Nat. Biotechnol.*, 23 (9): 1177-1180.
- Mlynarova, L., A. Loonen, J. Heldens, R.C. Jansen, P. Keizer, W.J. Stiekema and J.P. Nap, 1994. Reduced position effect in mature transgenic plants conferred by the chicken lysozyme matrix-associated region. *Plant Cell*, 6 (3): 417-426.
- Mlynarova, L., R.C. Jansen, A.J. Conner, W.J. Stiekema and J.P. Nap, 1995. The MAR mediated reduction in position effect can be uncoupled from copy number-dependent expression in transgenic plants. *Plant Cell*, 7 (5): 599-609.
- Mlynarova, L., L.C.P. Keizer, W.J. Stiekema and J.P. Nap, 1996. Approaching the lower limits of transgene variability. *Plant Cell*, 8 (9): 1589-1599.
- Mor, T.S., Y.S. Moon, K.E. Palmer and H.S. Mason, 2003. Geminivirus vectors for high-level expression of foreign proteins in plant cells. *Biotechnol. Bioeng.*, 81 (4): 430-437.
- Mullineaux, P.M., F. Guerineau and G.P. Accotto, 1990. Processing of complementary sense RNAs of *Digitaria streak* virus in its host and in transgenic tobacco. *Nucleic Acids Res.*, 18 (24): 7259-7265.
- Nguyen, H.T., S. Leelavathi and V.S. Reddy, 2004. Bacteriophage T7 RNA polymerase directed inducible and tissue-specific over-expression of foreign genes in transgenic plants. *Plant Biotechnol. J.*, 2 (4): 301-310.
- Nowak, W., M. Gawlowska, A. Jarmolowski and J. Augustyniak, 2001. Effect of nuclear matrix attachment regions on transgene expression in tobacco plants. *Acta Biochem. Polonica*, 48 (3): 637-646.
- Nussaume, L., M. Vincentz and M. Caboche, 1991. Constitutive nitrate reductase: A dominant conditional marker for plant genetics. *Plant J.*, 1 (2): 267-274.
- Ow, D., 2002. Recombinase-directed plant transformation for the post-genomic era. *Plant Mol. Biol.*, 48 (1-2): 183-200.
- Ow, D.W., K.V. Wood, M. DeLuca, J.R. De Wet, D.R. Helinski and S.H. Howell, 1986. Transient and stable expression of the firefly luciferase gene in plant cells and transgenic plants. *Science*, 234 (4778): 856-859.
- Page, A.F. and S.C. Minocha, 2004. Analysis of Gene Expression in Transgenic Plants. In: Methods in Molecular Biology. Transgenic Plants: Methods and Protocols, Pena, L. (Ed.). Vol. 286, Humana Press Inc., Totowa, NJ., pp: 291-311.
- Parinov, S., M. Sevugan, D. Ye, W.C. Yang, M. Kumaran and V. Sundaresan. 1999. Analysis of flanking sequences from dissociation insertion lines. A database for reverse genetics in Arabidopsis. *Plant Cell*, 11 (12): 2263-2270.
- Perera, R.J., C.G. Linard and E.R. Signer, 1993. Cytosine deaminase as a negative selective marker for Arabidopsis. *Plant Mol. Biol.*, 23 (4): 793-799.

- Perl, A., S. Galili, O. Shaul, I. Ben-Tzvi and G. Galili, 1993. Bacterial dihydrodipicolinate synthase and desensitized aspartate kinase: Two novel selectable markers for plant transformation. *Biotechnology*, 11 (6): 715-718.
- Petra, H.D., S.B. Surin and P.M. Waterhouse, 2003. A suite of novel promoters and terminators for plant biotechnology. II. The pPLEX series for use in monocots. *Funct. Plant Biol.*, 30 (4): 453-460.
- Prasher, D.C., V.K. Eckenrode, W.W. Ward, F.G. Prendergast and M.J. Cormier, 1992. Primary structure of the *Aequorea victoria* green fluorescent protein. *Gene*, 111 (2): 229-233.
- Reynolds, P.H.S., 1999. Inducible Gene Expression in Plants. CABI Publishing: Oxon, New York.
- Richards, H.A., M.D. Halfhill, R.J. Millwood and C.N. Stewart, 2003. Quantitative GFP fluorescence as an indicator of recombinant protein synthesis in transgenic plants. *Plant Cell Rep.*, 22 (2): 117-121.
- Richter, L.J., Y. Thanavala, C.J. Arntzen and H. S. Mason, 2000. Production of hepatitis B surface antigen in transgenic plants for oral immunization. *Natl. Biotechnol.*, 18 (11): 1167-1171.
- Riggs, C.D. and M.J. Chrispeels, 1987. Luciferase reporter gene cassettes for plant gene expression studies. *Nucleic Acids Res.*, 15 (19): 8115.
- Roberts, L.S., R.D. Thompson and R.B. Flavell, 1989. Tissue-specific expression of a wheat high molecular weight glutenin gene in transgenic tobacco. *Plant Cell*, 1 (6): 569-578.
- Robinson, M.O. and M.I. Simon, 1991. Determining transcript number using the polymerase chain reaction. P<sub>gk</sub>-2, mP2 and PGK-2 transgene mRNA levels during spermatogenesis. *Nucleic Acids Res.*, 19 (7): 1557-1562.
- Rocha-Sosa, M., U. Sonnewald, W. Frommer, M. Stratmann, J. Schell and L. Willmitzer, 1989. Both developmental and metabolic signals activate the promoter of a class I patatin gene. *EMBO. J.*, 8: 23-29.
- Sabelli, P.A. and P.R. Shewry, 1991. Characterization and organization of gene families at the Gli-1 loci of bread and durum wheats by restriction fragment analysis. *Theor. Applied Genet.*, 83 (2): 209-216.
- Sabelli, P.A. and P.R. Shewry, 1995a. Gene Characterization by Southern Analysis. In: *Methods in Molecular Biology. Plant Gene Transfer and Expression Protocols*, Jones, H. (Ed.). Vol. 49, Humana Press Inc: Totowa, NJ., pp: 161-180.
- Sabelli, P.A. and P.R. Shewry, 1995b. Northern Analysis and Nucleic Acid Probes. In: *Methods in Molecular Biology. Plant Gene Transfer and Expression Protocols*, Jones, H. (Ed.). Vol. 49, Humana Press Inc: Totowa, NJ. pp: 213-228.
- Schardl, C.L., A.D. Byrd, G. Benzion, M.A. Altschuler, D.F. Hildebrand and A.G. Hunt, 1987. Design and construction of a versatile system for the expression of foreign genes in plants. *Gene*, 61 (1): 1-11.
- Schöffl, F., G. Schroder, M. Kliem, M. Rieping, 1993. An SAR-sequence containing 395 bp-DNA fragment mediates enhanced, genedose-correlated expression of a chimeric heat shock gene in transgenic tobacco plants. *Transgenic Res.*, 2 (2): 93-100.
- Schroeder, H.E., A.H. Schotz, T. Wardley-Richardson, D. Spencer and T.J.V. Higgins, 1993. Transformation and regeneration of two cultivars of pea (*Pisum sativum* L.). *Plant Physiol.*, 101 (3): 751-757.
- Scutt, C.P., E. Zubko and P. Meyer, 2002. Techniques for the removal of marker genes from transgenic plants. *Biochimie*, 84 (11): 1119-1126.
- Shah, D.M., R.B. Horsch, H.J. Klee, G.M. Kishore, J.A. Winter, N.E. Turner, C.M. Hironaka, P.R. Sanders, C.S. Gasser, S. Aykent, N.R. Siegel, S.G. Rogers and R.T. Fraley, 1986. Engineering herbicide tolerance in transgenic plants. *Science*, 233 (4762): 478-481.
- Simpson, C.G. and J.W.S. Brown, 1995a. RNase A/T1 Protection Assay. In: *Methods in Molecular Biology. Gene Transfer and Expression Protocols*, Jones, H. (Ed.). Vol. 49, Humana Press Inc., Totowa, NJ. pp: 239-247.
- Simpson, C.G. and J.W.S. Brown, 1995b. Primer Extension Assay. In: *Methods in Molecular Biology. Gene Transfer and Expression Protocols*, Jones, H. (Ed.). Vol. 49, Humana Press Inc., Totowa, NJ. pp: 249-256.
- Singer, T. and E. Burke, 2003. High-Throughput TAIL-PCR as a Tool to Identify DNA Flanking Insertions. In: *Methods in Molecular Biology. Methods and Protocols*, Grotewold, E. Vol. 236. Humana Press, Inc., Totowa, NJ., pp: 241-271.
- Snowdon, R.J., P. Böttinger, T. Pickardt, W. Köhler and W. Friedt, 2001. Physical localization of transgenes on *Vicia faba* chromosomes. *Chromosome Res.*, 9 (7): 607-610.
- Somers, D.A., H.W. Rines, W. Gu, H.F. Kaeppler and W.R. Bushnell, 1992. Fertile, transgenic oat plants. *Biotechnology*, 10 (12): 1589-1594.
- Southern, E.M., 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.*, 98 (3): 503-517.
- Stalberg, K., M. Ellerstrom, L.G. Josefsson and L. Rask, 1993. Deletion analysis of a 2S seed storage protein promoter of *Brassica napus* in transgenic tobacco. *Plant Mol. Biol.*, 23 (4): 671-683.

- Stayton, M., M. Harpster, P. Brosio and P. Dunsmuir, 1991. High-level, seed-specific expression of foreign coding sequences in *Brassica napus*. Aust. J. Plant Physiol., 18 (5): 507-517.
- Stewart, C.N., 1996. Monitoring transgenic plants using in vivo markers. Nat. Biotechnol., 14 (6): 682.
- Stewart, C.N., 2001. The utility of green fluorescent protein in transgenic plants. Plant Cell Rep., 20 (5): 376-382.
- Stewart, C.N., 2006. Go with the glow: Fluorescent proteins to light transgenic organisms. Trends Biotechnol., 24 (4): 155-162.
- Svab, I., E.C. Harper, J.D.G. Jones and P. Maliga, 1990. Aminoglycoside-3'-denyltransferase confers resistance to spectinomycin and streptomycin in *Nicotiana tabaccum*. Plant Mol. Biol., 14 (2): 197-205.
- Ten Hoopen, R., T.R. Robbins, P.F. Fransz, B.M. Montijn, J.L. Oud, A.G.M. Gerats and N. Nanninga, 1996. Localization of T-DNA insertions in petunia by fluorescence in situ hybridization: Physical evidence for suppression of recombination. Plant Cell, 8 (5): 823-830.
- Thompson, C.J., N.R. Movva, R. Tizard, R. Cramer, J.E. Davies, M. Lauwereys and J. Botterman, 1987. Characterization of the herbicide-resistance gene bar from *Streptomyces hygrosopicus*. EMBO. J., 6 (9): 2519-2523.
- Tichopad, A., M.W. Pfaffl and A. Didier, 2003. Tissue-specific expression pattern of bovine prion gene: Quantification using real-time RT-PCR. Mol. Cell Probes, 17 (1): 5-10.
- Tissier, A.F., S. Marillonnet, V. Klimyuk, K. Patel, M.A. Torres, G. Murphy and J.D.G. Jones, 1999. Multiple independent defective suppressor-mutator transposon insertions in Arabidopsis: A tool for functional genomics. Plant Cell, 11 (10): 1841-1852.
- Towbin, H., T. Staehelin and J. Gordon, 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose. Procedure and some applications. Proc. Natl. Acad. Sci., 76 (9): 4350-4354.
- Ülker, B., G.C. Allen, W.F. Thompson, S. Spiker and A.K. Weissinger, 1999. A tobacco matrix attachment region reduces the loss of transgene expression in the progeny of transgenic tobacco plants. Plant J., 18 (3): 253-263.
- Vain, P., B. Worland, A. Kohli, J.W. Snape, P. Christou, G.C. Allen and W.F. Thompson, 1999. Matrix attachment regions increase transgene expression levels and stability in transgenic rice plants and their progeny. Plant J., 18 (3): 233-242.
- Valasek, M.A. and J.J. Repa, 2005. The power of real-time PCR. Adv. Physiol. Educ., 29 (3): 151-159.
- Van den Elzen, P.J.M., J. Townsend, K.Y. Lee and J.R. Bedbrook, 1985. A chimeric hygromycin resistance gene as a selectable marker in plant cells. Plant Mol. Biol., 5 (5): 299-302.
- Van der Geest, A.H.M., G.E. Hall, S. Spiker and T.C. Hall, 1994. The beta-phaseolin gene is flanked by matrix attachment regions. Plant J., 6 (3): 413-423.
- Waldron, C., E.B. Murphy, J.L. Roberts, G.D. Gustafson, S.L. Armour and S.K. Malcolm, 1985. Resistance to hygromycin B. A new marker for plant transformation studies. Plant Mol. Biol., 5 (2): 103-108.
- Walters, D.A., C.S. Vetsch, D.E. Potts and R.C. Lundquist, 1992. Transformation and inheritance of a hygromycin phosphotransferase gene in maize plants. Plant Mol. Biol., 18 (2): 189-200.
- Weeks, T., 2000. Transformation of wheat with the cyanamide hydratase gene. United State Patent 09/518,988. Available on world wide web, <http://ott.arsusda.gov/>.
- Wong, M.L. and J.F. Medrano, 2005. Real-time PCR for mRNA quantitation. Biotechniques, 39 (1): 1-11.
- Wright, M., J. Dawson, E. D under, J. Suttie, J. Reed, C. Kramer, Y. Chang, R. Novitzky, H. Wang and L. Artim-Moore, 2001. Efficient biolistic transformation of maize (*Zea mays* L.) using the phosphomannose isomerase gene, *pmi*, as the selectable marker. Plant Cell Rep., 20 (5): 429-436.
- Yalow, R.S. and S.A. Berson, 1959. Assay of plasma insulin in human subjects by immunological methods. Nature, 184 (4699): 1648-1649.
- Yamamoto, Y.T., C.G. Taylor, G.N. Acedo, C.L. Cheng and M.A. Conkling, 1991. Characterization of cis-acting sequences regulating root-specific gene expression in tobacco. Plant Cell, 3 (4): 371-382.
- Zhang, W. and R. Wu, 1988. Efficient regeneration of transgenic rice plants from rice protoplasts and correctly regulated expression of foreign genes in the plants. Theor. Applied Genet., 76 (6): 835-840.