

ISSN 1682-296X (Print)

ISSN 1682-2978 (Online)



Bio Technology



ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Increased Gum Arabic Production after Infestation of *Acacia senegal* with *Aspergillus flavus* and *Pseudomonas pseudoalcaligenes* Transmitted by *Agrilus nubeculosus*

¹H.A.El Atta, ¹I.M. Aref and ²S.A. Khalil

¹Department of Plant Production, King Saud University, Postal Code 11451,
P.O. Box 2460, Riyadh, Saudi Arabia

²Forest National Corporation, Postal Code 123543, P.O. Box 331, Khartoum, Sudan

Abstract: The aim of this study was to investigate the correlation between the beetle *A. nubeculosus* and gum arabic production by *A. senegal*. Some trees were tapped and left open to facilitate infestation by *A. nubeculosus* and others were covered with wire mesh as control. Gum yield, physical and chemical properties of gum were determined for infested and control trees. *A. senegal* infested by *A. nubeculosus* produced significantly more gum than control trees. Infestation also caused significant changes in some physical properties of gum (Colour, shape, size, moisture content and optical rotation) and chemical properties (Ferrous, calcium, magnesium and nitrogen contents), whereas, no significant difference was recorded in phosphorus and manganese contents. *Aspergillus flavus* Link. and *Pseudomonas pseudoalcaligenes* Monias were isolated from the mouth parts of *A. nubeculosus*. Inoculation of the tapped branches of *Acacia senegal* with a suspension of *A. flavus* alone or in combination with *P. pseudoalcaligenes*, resulted in highly significant gum yield as compared to control. It seems that *A. flavus* and *P. pseudoalcaligenes* acted as elicitors that have stimulated the synthesis of gum arabic. *A. nubeculosus* transmitted *A. flavus* and *P. pseudoalcaligenes* to the tapped areas. Hence, the beetle was significantly associated with gum production.

Key words: *Acacia senegal*, gum arabic, *Aspergillus flavus*, *Pseudomonas pseudoalcaligenes*, *Agrilus nubeculosus*

INTRODUCTION

Plant secondary metabolites are important sources of many food ingredients and disease-preventive phytochemicals. In recent years, the demand for these products has increased dramatically. The supply of the source plants, however, is often limited because of disease, changes in climate, changes in the economical development, or other problems in the growing regions (Fu, 1998). Gum arabic (INS 414) is the dried gummy exudate from the stems and branches of *Acacia senegal* (L) Willd. or of other related species of Acacia (Family: Leguminosae) (Dondain and Phillips, 1999). It is defined by the FAO/WHO Joint Expert Committee for Food Additives (JECFA) as 'a dried exudation obtained from the stems of *A. senegal* or closely related species of Acacia (family Leguminosae)' FAO/WHO, 1999). Although there are many species of Acacia trees botanically, only two species, namely *A. senegal* and *A. seyal* are acceptable to the Codex Alimentarius Commission (Al-Assaf *et al.*, 2003; Awouda, 2000;

FAO/WHO, 1999). Gum arabic has wide industrial uses as a stabilizer, thickening agent and emulsifier, mainly in the food industry (e.g., in soft drinks syrup, gummy candies and marshmallows) but also in the textile, pottery, lithography, cosmetics and pharmaceutical industries (Verbeken *et al.*, 2003). It has been approved for use as food additives by the US Food and Drug Administration and is on the list of substances that is a Generally Recognized as Safe (GRAS) with specific limitations. Supplementing diet of the laying hens with gum arabic resulted in 15% increase ($p < 0.05$) in feed intake and egg shell thickness by 5.98 and 31.58%, respectively as compared to the control (El Kheir *et al.*, 2009). In folk medicine, gum arabic has been reported to be used internally for the treatment of inflammation of the intestinal mucosa and externally to cover inflamed surfaces (Gamal El-din *et al.*, 2003). It is an edible, dried, gummy exudate that is rich in non-viscous soluble fiber (Phillips and Williams, 2000). Clinically, it has been tried in patients with chronic renal failure and it was claimed that it helps reduce urea and creatinine plasma concentrations

and reduces the need for dialysis from 3 to 2 times per week (Suliman *et al.*, 2000). Despite the fact that gum arabic is widely used as a vehicle for drugs in experimental physiological and pharmacological experiments and is assumed to be an “inert” substance, some recent reports have claimed that it possesses anti-oxidant, nephroprotectant and other effects (Ali *et al.*, 2008). Pharmacologically, gum arabic has been claimed to act as an anti-oxidant and to protect against experimental hepatic-, renal- and cardiac toxicities in rats (Ali *et al.*, 2009). Analysis of gum arabic has indicated that it consists of three distinct components. Fraction 1 which represents 88.4% of the total, is an arabinogalactan with molecular mass 2.79×10^5 and is deficient in protein. Fraction 2 which represents 10.4% of the total, is an arabinogalactan-protein complex with a molecular mass of 1.45×10^6 , containing ~50% of the total protein. It is envisaged that on average each molecule of fraction 2 consists of five carbohydrate blocks of molecular mass $\sim 2.8 \times 10^5$ covalently linked through a chain of amino acid residues. Fraction 3 represents only 1.24% of the total gum but contains ~25% of the total protein and has been shown to consist of one or more glycoproteins. Whereas the proteinaceous components of fractions 1 and 2 contain predominantly hydroxyproline and serine, this is not the case for fraction 3 (Randall *et al.*, 1989). Gum arabic is a branched-chain, complex polysaccharide, either neutral or slightly acidic, found as a mixed calcium, magnesium and potassium salt of a polysaccharidic acid. The backbone is composed of 1, 3-linked β -D-galactopyranosyl units. The side chains are composed of two to five 1, 3-linked β -D-galactopyranosyl units, joined to the main chain by 1, 6-linkages. Only a few plant species are cultivated at present to obtain gums used in the food industry as additives; most of them belong to the Leguminosae family. Some examples are: *Acacia senegal*, source of acacia or arabic gum; *Astragalus* spp., source of tragacanth; *Cyamopsis tetragonolobus*, source of guar gum; *Ceratonia siliqua*, source of locust bean gum (Ibanez and Ferrero, 2003). The most commonly recognized is arabic gum but a wide range of other tree exudates is used for variety of uses in their countries of origin, such as mesquite gum (Anderson and Farquhar, 1982; Anderson and Wang, 1990; Vernon-Carter *et al.*, 2000). *A. senegal* trees grow widely across in Sahelian countries of Africa, especially in Sudan and gum arabic, as a food additive, has been an important item of commerce since ancient times (Glicksman, 1969). The gum belt in Sudan provides a natural buffer zone between the desert in the North and the more fertile agricultural lands in the

South. Deforestation within the gum belt has led to an increase in desert encroachment and threatens agricultural production (Keddeman, 1994; Olsson and Ardo, 2002). Following the Sahel drought of the 1970s and 1980s a southward shift in the tapping of gum has been reported (IIED and IES, 1990) as people moved from the more fragile environment in the northern parts of the gum belt to the less fragile and better environment of the south. Over the last three to four decades the land use practices have moved from a rotation with long fallow periods (15-20 years) of gum cultivation interspersed with short period of cultivation (4-6 years) towards a more or less continuous cultivation (Barbier, 2000). Gum arabic agriculture plays an important role as a cash crop produced in the traditional rain-fed areas of North Kordofan in western Sudan (El-Dukheri, 1997). *A. senegal* trees are managed in the Sudan in an agroforestry system known as the bush-fallow system. However, the recent disruption of this traditional agroforestry system due to the misuse of land, drought and desertification is considered to be among the main factors that have led to fluctuations in gum arabic yield and the consequent instability of supply (Seif El Din, 1995). Apart from drought, desertification and mismanagement, gum arabic production also varies as a result of complex factors in the physical, biological and socio-economic environments. The impact of all or some of these factors on gum arabic production has been reported by Abdel Rahman (2001). There is still an information gap regarding the factors that control gum arabic yield. The International Institute for Environment and Development and the Institute of Environmental Studies (IIED and IES, 1989) reported that rainfall and its distribution pattern together with the minimum temperature during tapping and gum picking and the relative humidity, are the main factors affecting gum arabic yield. The decision to tap or not to tap trees is based on the production of gum per unit area (Mohammed and Rohle, 2011). However, the relationship between gum arabic yield and rainfall is complex and the available information is sparse and imprecise (IIED and IES, 1989). Large-scale planting programs with the help of local communities have been implemented since the early 1980s to restock the gum arabic belt in order to curb desertification and to improve the gum arabic yield and production in western Sudan (Rahim *et al.*, 2007). Gummosis is widespread in plant kingdom and is known to be produced by stress conditions such as heat, drought and wounding. Gums form a barrier at lesions hindering the invasion of microorganisms. Fungal and bacterial infections have been linked with the synthesis process, although this

has not been proved (Greenwood and Morey, 1979; Ghosh and Purkayastha, 1962; Luckner, 1990).

Natives inhabiting the gum producing areas in the Sudan, believe that high gum yield by *A. senegal* is directly correlated with the abundance of certain beetles locally known as Al Garraha (Injector) i.e., in years when the occurrence of these beetles was high, gum production was maximum (Osman, 1993). Several gum producers interviewed in Kordofan State (western Sudan), firmly believe that *Al Garraha* (Injector) pierces holes into the tapped branches of *A. senegal* (El Khalifa *et al.*, 1989).

This study aimed at testing the hypothesis that gum arabic production is correlated with certain beetles that infest the freshly tapped branches of *A. senegal*. The study is the first attempt in this direction.

MATERIALS AND METHODS

The study area: The study was carried out in January-December (2009) in El Himaiyra forest plantation (13°19'05 N-13°09'58 E, 545 m above sea level) 12 km north to El Obeid (Capital of Northern Kordofan State). Rainfall varies between 250-450 mm from May to October. The soil is uniform deep reddish sand with little textural differentiation in the profile. The climate is long hot season, rainy season with scattered irregular rains and short winter.

Identification of Al Garraha (Injector): Samples of the adult *Al Garraha* (Injector) were collected from the tapped branches of *A. senegal* and sent to the British Museum of Natural History for identification. The insect was identified as *Agrilus nubeculosus* Fairm (Coleoptera, Buprestidae).

The effect of *A. nubeculosus* on gum arabic yield: A total of 120 healthy *A. senegal* trees (5-9, 10-14 and 15-20 years old) with mean diameters of 24.75, 33.2 and 39.9 cm, respectively and 2.7, 3.4 and 4.3 m mean height, respectively were selected randomly from El Himaiyra forest. Each tree class comprised 40 trees. In each tree, two branches were selected randomly and tapped with a sharp knife locally known as sonky (20 cm long and 5 cm wide with a 3 m long wooden handle). Tapping removes only the outer bark without injuring the cambium. Tapping was carried out at two intervals, the first tapping immediately after the rainy season (November) and the second tapping in summer (February). Half of the tapped trees in each class was left open (Group 1), whereas the remainder (Group 2) were covered with a wire mesh to prevent insect infestation and served as control. The produced gum arabic was collected from trees after 40

days from tapping, weighed and kept in a refrigerator at 10°C for further analysis.

Physical properties of gum arabic: Physical properties of gum arabic produced by both groups of trees were determined as follows:

Viscosity: Viscosity of gum collected from all trees was determined using a viscometer (Brook-Field model DV-1+1). One gram of gum was dissolved in 100 ml distilled water in a 500 mL conical flask to make 1% solution in reference to 4% Na OH. This was replicated three times for each group of trees (1 and 2). Viscosity of gum was measured in cps (centipoises) and calculated as follows:

$$\text{Reduced Viscosity} = V - V_0 / V_0 * C$$

$$\text{Intrinsic Viscosity} = (V - V_0) * C_0 / V_0 * C$$

Where:

- V = Viscosity of solvent
- V₀ = Viscosity of gum solution
- C = Concentration of gum solution
- C₀ = Concentration of solvent

Moisture content: Two gram samples of gum from each group of trees (1 and 2) were placed on crucibles and oven dried in an oven at 105°C for 5 h until constant weights were obtained and by subtraction the moisture content % was calculated (FAO, 1995). Samples from each group of trees were tri-replicated.

Ash value: *A. crucible* was heated at 55°C, cooled in a dissector and weighed. Two grams of each gum sample were placed in a crucible and ignited at 550°C in a Heraeus electronic muffle until free from carbon, cooled in a dissector and weighed. This was replicated three times for gum from each group of trees. The ash value was then calculated according to FAO (1995).

Specific optical rotation: Optical rotation of gum from each group of trees was determined using ADP 220 Polarimeter, The Specific Optical Rotation (SOR) was determined for 1% solution (w/v) of gum in distilled water on dry weight basis using an optical activity Polarimeter fitted with a sodium lamp and a cell path length of 20 cm. The solutions were passed through a No. 42 filter paper before carrying out the measurements at room temperature (30.2-33.4°C). Readings were taken 5 times. Tri-replicates of gum samples were used. The specific optical rotation was calculated as follows:

$$\text{SOR} = \alpha * 100 / (C * L)$$

Where:

- α = Observed optical rotation
C = Concentration of the solution
L = Length of the Polarimeter tube

Chemical properties of gum arabic: Gum arabic produced by trees of each group was chemically analyzed.

Nitrogen content: The nitrogen content of gum was determined according to the Micro-Kjeldahl method (AOAC, 1984).

Cationic composition: The cationic composition of gum obtained from group 1 and 2 trees was determined (FAO, 1995).

Extraction of material from *A. nubeculosus*: Twenty adults of *A. nubeculosus* were randomly selected, rinsed thrice in sterile distilled water. A small loop was inserted into the mouth of *A. nubeculosus* to collect some of the liquid contents. The mouth contents were then streaked on petridishes containing sterile PDA and Nutrient Agar and incubated at $30 \pm 2^\circ\text{C}$ in an attempt to isolate any microorganism (s). Gram staining and oxidase test were performed to assist identification of bacteria (Collins *et al.*, 1995).

Identification of cultures: On PDA, a fungus was isolated, purified and identified. As for the bacterium isolated on Nutrient Agar, the following tests for identification were carried out (Collins *et al.*, 1995).

Gram staining: A loop-full of bacterial colony from the slanted stock culture was mixed with a drop of sterile distilled water at the centre of a clean glass slide. A thin film was obtained and left to dry. It was then fixed by passing over a flame 5 times. The dry smear was then flooded with crystal violet-ammonium oxalate prepared by dissolving 2 g of crystal violet in 20 mL of 95% ethanol and 0.8 g ammonium oxalate in 80 mL of distilled water. The excess dye was washed off with gently running tap water. The smear was decolorized with 95% ethanol. It was then stained with safranin (0.25 g safranin+10 mL ethanol and completed to 100 mL by the addition of water) for 30 sec. Excess safranin was washed off with running tap water and the smears were blotted dry and finally examined under the microscope. Bacterial cells which were violet coloured were recorded as gram-positive and those which were red are gram-negative.

Oxidase test: Two milliliter of 10% of aqueous methyl phenylenediamine dihydrochloride (w/v) were added to a filter paper in a sterile petridish. Using a platinum loop, a heavy amount of growth from a 24 h old culture on Nutrient Agar was added to the filter paper and quickly spread. Development of a purple colour within 10 sec was recorded as a positive reaction indicating the presence of cytochrome oxidase, otherwise the reaction was considered negative (Collins *et al.*, 1995).

Description and identification of cultures: The cultures prepared were dominated by a fungus and a bacterium and were identified. As for the fungus:

The colony diameter was 4.0-4.5 cm, Yellowish-green becoming green with age on Obverse view and Creamy-yellow on reverse view. The head: Radiating, becoming loosely columnar with age. The stipe: Long, verrucose and hyaline. The Vesicle: Dome-shaped. Medullae: Present and small. Phialidae: small ampullifprm. Conidia: Globose to sub-globose, usually rough yellowish green. Accordingly, the fungus was identified as *Aspergillus flavus* Link.

The bacterium colony was negative for gram stain and oxidase test and identified as *Pseudomonas pseudoalcaligenes* Monias.

The effect of *A. flavus* and *P. pseudoalcaligenes* on gum production: Stock cultures of *A. flavus* on PDB (potato dextrose broth) and *P. pseudoalcaligenes* on NB (nutrient broth) were prepared and incubated at $30 \pm 1^\circ\text{C}$ for 3 weeks. A total of fifty healthy 5-10 years old *A. senegal* trees in the study area were selected randomly and 2 branches/tree were tapped and immediately treated as follows:

- Inoculated with *A. flavus* (A)
- Inoculated with *P. pseudoalcaligenes* (P)
- Inoculated with both *A. flavus* and *P. pseudoalcaligenes* (AP)
- Not inoculated and left open (N)
- Not inoculated and covered with a wire mesh and served as control trees (group C)

Inoculation was carried out by taking 10 mL of each liquid culture by using a syringe and sprayed on the surface of the freshly tapped branch. Ten trees were used for each treatment.

Statistical analysis was carried out using ANOVA, Tukey's test for mean separation and T-test by using SAS statistical package (Cody and Smith, 1987). Significance tests of means were determined at the 5% level.

RESULTS

Activity of *A. nubeculosus*: Two to eight hours after tapping, the insect pierces holes into the tapped branches of *A. senegal* by pushing a sharp posterior organ into the freshly tapped surface. The beetle secretes a dirty greenish substance over the wounded area. Usually 1-3 insects were found on the tapped branch. The insect took flight from the tapped area of the branch after 1-4 h leaving behind a more or less rounded dirty green spots.

The effect of *A. nubeculosus* on gum yield: Gum production was considerably higher in group 1 trees (infested by *A. nubeculosus*) as compared to group 2 trees (control) in all of the three age classes studied 5-9 (Table 1), 10-14 (Table 2) and 15-20 years old (Table 3) after both first and second tapping. These differences were highly significant ($p = 0.0001$).

Physical properties of gum arabic: Table 4 summarizes description of colour, shape and size of gum arabic obtained from trees in groups 1 and 2. Colour and shape of the produced gum were different between the two groups of trees. Gum produced by infested trees was more or less spherical or rounded and red in colour, whereas that from control trees was irregular in shape and white creamy. Moisture content of gum was significantly ($p = 0.01$) higher in the control trees as compared to those infested. The specific optical rotation of gum was significantly ($p = 0.001$) greater in infested than in control trees. However, there was no significant difference in viscosity and ash contents of gum (Table 4).

Chemical properties of gum Arabic: Infestation of *A. senegal* with *A. nubeculosus* had affected the chemical

Table 1: Mean gum arabic yield/tree (g) of 5-9 years old *A. senegal*

Tree group	Mean/tree	t-value	p-value
First tapping			
1 (infested)	102.4	-60.80	≤ 0.0001
2 (control)	3.5		
Second tapping			
1 (infested)	99.2	-60.80	≤ 0.0001
2 (control)	2.1		

Means were computed for 20 replicates in each treatment

Table 2: Mean gum arabic yield/tree (g) of 10-14 years old *A. senegal*

Tree group	Mean/tree	t-value	p-value
First tapping			
1 (infested)	123.3	-24.47	≤ 0.0001
2 (control)	2.5		
Second tapping			
1 (infested)	110.6	-63.21	≤ 0.0001
2 (control)	2.7		

Means were computed for 20 replicates in each treatment

Table 3: Mean gum arabic yield/tree (g) of 15-20 years old *A. senegal*

Tree group	Mean/tree	t-value	p-value
First tapping			
1 (infested)	146.2	-13.11	≤ 0.0001
2 (control)	2.3		
Second tapping			
1 (infested)	111.6	-65.15	≤ 0.0001
2 (control)	1.3		

Means were computed for 20 replicates in each treatment

Table 4: Physical properties of gum arabic from infested and control *A. senegal*

Properties	Moisture (%)	t-value	p-value
Tree group			
1	7.66	7.11	≤ 0.02
2	8.70		
Specific Optical Rotation (°)			
1	-33.23	31.90	≤ 0.001
2	-29.40		
Viscosity (cps)			
1	1.56	-1.67	$> 0.05^*$
2	1.30		
Ash %			
1	3.10	-2.60	$> 0.05^*$
2	2.80		

1: Infested, 2: Control, * Not significant

Table 5: Chemical properties of gum arabic from infested and control *A. senegal*

Tree group	Moisture (%)	t-value	p-value
Ferrous %			
1	1.66×10^{-3}	-5.80	≤ 0.02
2	1.56×10^{-3}		
Calcium %			
1	0.50	-10.97	≤ 0.05
2	0.41		
Magnesium%			
1	4.2×10^{-2}	37.00	≤ 0.001
2	7.9×10^{-2}		
Nitrogen%			
1	0.30	8.66	≤ 0.01
2	0.35		
Phosphorus%			
1	0.24	-1.99	$> 0.05^*$
2	0.23		
Manganese%			
1	6.13×10^{-5}	-3.63	$> 0.05^*$
2	8.17×10^{-5}		

1: infested, 2: control, *Not significant

Table 6: Effect of inoculation on gum production by *A. senegal* (ANOVA)

Source	d.f.	SS	MS	F value	p-value
Model	4	355867.16	88966.79	7509.87	0.0001
Error	45	533.09	11.84		

$R^2 = 0.99$

Table 7: Gum produced by inoculated and control *A. senegal*

Treatment	Mean gum (g)
A	249.66 ^a
P	144.33 ^b
AP	136.19 ^c
N	60.53 ^d
C	3.4 ^e

Means were computed for 10 replicates in each treatment. Means followed by the same letter are not significantly different according to Tukey's test A: Inoculated with *A. flavus* S: Inoculated with *P. pseudoalcaligenes* AS: Inoculated with both *A. flavus* and *P. pseudoalcaligenes* N: Not inoculated and left open C: Not inoculated and covered with a wire mesh and served as control (group C)

constitution of gum arabic. Ferrous, calcium, magnesium and nitrogen contents were significantly affected, whereas, no significant effect was recorded in phosphorus and manganese contents (Table 5).

The effect of *A. flavus* and *P. pseudoalcaligenes* on gum production: Inoculation of the tapped branches of *A. senegal* with *A. flavus* and *P. pseudoalcaligenes* alone or in combination significantly ($p = 0.0001$) increased gum production as compared to control trees (Table 6). Gum production was significantly more from trees inoculated with *A. flavus* (249.6 g) than from those inoculated with *P. pseudoalcaligenes* (136.2 g), both *A. flavus* and *P. pseudoalcaligenes* (144.3 g), not inoculated and left open (60.5 g) and not inoculated and covered with a wire mesh (3.4 g). These differences were highly significant (Table 7).

DISCUSSION

A. senegal trees infested by *A. nubeculosus*, produced significantly more gum arabic than control trees. Most physical and chemical properties of gum arabic were different when infested trees were compared with control ones. Gum from normally tapped *A. senegal* was different from trees infested by wood-borers in two important aspects concerning some of the peripheral end group positions involving rhamnose and glucuronic acid residues (Anderson and Dea, 1967). This was substantiated by the fact that inoculation of the tapped areas of *A. senegal* with *A. flavus* alone or in combination with *P. pseudoalcaligenes*, resulted in a highly significant increase in gum arabic production as compared to control. Both *A. flavus* and *P. pseudoalcaligenes* were isolated from the mouth of the adult *A. nubeculosus*. Hence, the beetle is basically a vector that mechanically transmitted both microorganisms. These microorganisms might have exerted stress on *A. senegal* trees that probably modified some physiological processes as part of the defensive mechanism of trees against infection. It has been suggested that gum cysts development was preceded by various changes in the xylem and phloem induced by physiological disturbances in the tree due to some pathological conditions caused by bacterial activity (Ghosh and Purkayastha, 1962). In almond plants, gum formation was attributed to the diffusion of fungal toxins (Frisullo and Graniti, 1985). However, non of these studies have reported a particular microorganism which might be directly or indirectly involved in the process of gum production. It is probable that *A. flavus* and *P. pseudoalcaligenes* acted as elicitors. Elicitors are

compounds that stimulate the synthesis of phytoalexins and various secondary metabolites in plants. Elicitors such as fungal wall materials, plant and microbial polysaccharides and some chemicals increase secondary metabolite production in various plant cell and tissue culture systems. The results of the present study are in agreement with Figueredo (1990) and Johnson *et al.* (1991) who showed that elicitation increased the production of many food ingredients in various types of plant cell and tissue culture. Biotic elicitors are classified as complex or defined depending on their origin and molecular structure. Elicitors with defined composition include carbohydrates, oligosaccharides, peptides and elicitors with complex composition include yeast cell wall, mycelia cell wall and fungal spores (Radman *et al.*, 2003) which is in agreement with the findings of the present study. The greatest gum production by *Prosopis laevigata* nodal explants cultures occurred when the culture medium was inoculated with the fungal mycelium of *A. nidulans* and bacterial biomass of *P. pseudoalcaligenes* in combination with an incubation temperature of 35°C. These treatments were non-significantly different among themselves but were significantly different from the rest of the treatments (Orozco-Villafuerte *et al.*, 2005). The present study shed the light on the link between *A. nubeculosus* and gum arabic production. The beetle acted as a vector that transmitted a fungus (*A. flavus*) and a bacterium (*P. pseudoalcaligenes*) which both resulted in a considerable increase in gum production and changes in some physical and chemical properties of gum arabic. The exact process behind quantitative as well as qualitative effect of these microorganisms on gum arabic needs further research. Also there is a need for this technology to be processed in a suitable form and transferred to the farmers in the gum arabic belt.

CONCLUSION

Forest resources are extremely important sources of subsistence for rural communities (Tolunay *et al.*, 2008). Microorganisms played an important role in gum arabic yield and its physical as well as chemical properties. Inoculation of tapped *A. senegal* trees with *A. flavus* alone produced the maximum gum yield followed by inoculation with *P. pseudoalcaligenes* and a comparatively less gum yield was produced but significantly more than in control, when trees were inoculated simultaneously with both fungus and bacterium. This biotechnology can be applied and it is of high economic value since many poor countries rely on exports of gum arabic.

ACKNOWLEDGMENTS

The authors would like to thank King Abdul Aziz City for Science and Technology for supporting this study.

REFERENCES

- AOAC, 1984. Official Methods of Analysis. 14th Edn., Association of Official Analytical Chemists, Washington, DC.
- Abdel Rahman, H.M., 2001. Effect of some growth parameters and age on gum and seed yield of *Acacia senegal* plantations and natural stands in North Kordofan. M.Sc. Thesis, Faculty of Forestry, University of Khartoum, Sudan.
- Al-Assaf, S., T. Katayama, G.O. Phillips, Y. Sasaki and P.A. Williams, 2003. Quality control of gum arabic. *Foods Ingredients J. Jpn.*, 208: 771-780.
- Ali, A.A., K.E. Ali, A. Fadlalla and K.E. Khalid, 2008. The effects of G.A. oral treatment on the metabolic profile of chronic renal failure patients under regular haemodialysis in Central Sudan. *Nat. Prod. Res.*, 22: 12-21.
- Ali, B.H., A. Ziada and G. Blunden, 2009. Biological effects of gum arabic: A review of some recent research. *Food. Chem. Toxicol.*, 47: 1-8.
- Anderson, D.M.W. and I.C.M. Dea, 1967. Studies on Uronic Acid Materials. Part XXV. Some unusual forms of the gum from *Acacia senegal* Willd. *Carbohydr. Res.*, 6: 104-110.
- Anderson, D.M.W. and J.G.K. Farquhar, 1982. Gum exudates from the genus *Prosopis*. *Int. Tree Crops J.*, 2: 15-24.
- Anderson, D.M.W. and W.P. Wang, 1990. Composition of the gum from *Combretumpaniculatum* and four other gums which are not permitted food additives. *Phytochemistry*, 29: 1193-1195.
- Awouda, E.H.M., 2000. Production and commercialisation of gum Arabic. Proceedings of the Workshop on Management of Trees for Farmland Rehabilitation and Development, (MTFRD'00), Khartoum, Sudan, pp: 15-15.
- Barbier, E.B., 2000. The economic linkages between rural poverty and land degradation: Some evidence from Africa. *Agric. Ecosyst. Environ.*, 82: 355-370.
- Cody, R.P. and J.K. Smith, 1987. Applied Statistics and the SAS Programming Language. 2nd Edn., Elsevier Science Publishing Co. Inc., New York.
- Collins, C.H., P.M. Lyne and J.M. Grange, 1995. Collins and Lyne's Microbiological Methods. 7th Edn., A Hodder Arnold Publication, Linacre House, Jordan Hill, Oxford, ISBN-13: 978-0750606530, pp: 512.
- Dondain, G. and G.O. Phillips, 1999. The regulatory journey of gum arabic. *Food Foods Ingredients J. Jpn.*, 179: 38-56.
- EL-Khier, M.K.S., K.E.A. Ishag, A.E.G.A. Yagoub and A.A.A. Baker, 2009. Supplementing laying hen diet with gum Arabic (*Acacia senegal*): Effect on egg production, shell thickness and yolk content of cholesterol, calcium and phosphorus. *Asian J. Poultry Sci.*, 3: 9-14.
- El Khalifa, M.D., M.O.E. Samani and H.O. Abdel Nour, 1989. Gum arabic rehabilitation in The Republic of the Sudan. Report No. 1 (2), Institute of Environmental Studies, University of Khartoum, Sudan.
- El-Dukheri, I.A., 1997. Past changes and future prospects of traditional rainfed farming in North Kordofan, Sudan. Ph.D. Thesis, University of Munchen, Germany.
- FAO, 1995. Compendium of food additive specifications. Addendum 3 FAO Food and Nutrition Paper No. 52. Add.3. Rome: FAO.
- FAO/WHO, 1999. Compendium of food additives. Food and Nutrition Paper 52 Addendum 7, Rome.
- Figuereido, A.A., 1990. Mesquite: History, composition and food uses. *Food Technol.*, 44: 118-128.
- Frisullo, S. and A. Graniti, 1985. Gum formation in Almond plants infected by *Fusicum anygadali* Del. and possible consequences on the diffusion of fungal toxins. *Phytopathol. Meter.*, 24: 284-288.
- Fu, T.J., 1998. Safety considerations for food ingredients produced by plant cell and tissue culture. *Chemtech*, 28: 40-46.
- Gamal El-Din, A.M., A.M. Mostafa, O.A. Al-Shabanah, A.M. Al-Bekairi and M.N. Nagi, 2003. Protective effect of arabic gum against acetaminophen-induced hepatotoxicity in mice. *Pharmacol. Res.*, 48: 631-635.
- Ghosh, S.S. and S.K. Purkayastha, 1962. Anatomical studies of wood and bark of *Acacia senegal* Willd. trees with special reference to gum exudation. *Indian For.*, 88: 92-99.
- Glicksman, M., 1969. Natural Plant Exudates. Academic Press Inc., New York, USA..
- Greenwood, C. and P. Morey, 1979. Gummosis in honey mesquite. *Bot. Gazette*, 140: 32-38.
- IIED and IES, 1989. Gum arabic rehabilitation in the Republic of the Sudan: Stage 1 Report Volume 1. International Institute for Environment and Development (IIED) and Institute of Environmental Studies (IES), London.

- IIED and IES, 1990. Gum arabic belt rehabilitation in the Republic of the Sudan: Stage 1 Report Volume 2. Main Report and Recommendations for Action for the Project Preparation Unit of the Ministry of Finance and Economic Planning (Planning), The Republic of the Sudan. (IIED), England, London. <http://scialert.net/fulltext/?doi=rjf.2011.17.26> and org=10.
- Ibanez, M.C. and C. Ferrero, 2003. Extraction and characterization of the hydrocolloid from *Prosopis flexuosa* DC seeds. *Food Res. Int.*, 36: 455-460.
- Johnson, T.S., G.A. Ravishankar and L.V. Venkataraman, 1991. Elicitation of capsaicin production in freely suspended cells and immobilized cell cultures of *Capsicum frutescens* mill. *Food Biotechnol.*, 5: 197-205.
- Keddeman, W., 1994. Restocking of the gum belt for desertification control in Kordofan, Phase III. Report of the Evaluation Mission (UNSO/SUD/89/X05). Netherlands Economic Institute, Agricultural and Rural Development Department, Rotterdam, The Netherlands.
- Luckner, M., 1990. Secondary Metabolism Microorganisms, Plants and Animals. 3rd Edn., Springer-Verlag, Berlin, Germany, ISBN-13: 9783540502876, pp: 563.
- Mohammed, M.H. and H. Rohle, 2011. Gum Talha from *Acacia seyal* Del. variety Seyal in South Kordofan, Sudan. *Res. J. For.*, 5: 17-26.
- Olsson, L. and J. Ardo, 2002. Soil carbon sequestration in the degraded semiarid agro-ecosystems-perils and potentials. *Ambio*, 31: 471-477.
- Orozco-Villafuerte, J., L. Buendia-González, F. Cruz-Sosa and E. Vernon-Carter, 2005. Increased mesquite gum formation in nodal explants cultures after treatment with a microbial biomass preparation. *Plant Physiol. Biochem.*, 43: 802-807.
- Osman, M.E., 1993. Characterization and fractionation of the gum from *Acacia senegal*. Ph.D. Thesis, Salford University, United Kingdom.
- Phillips, G.O. and P.A. Williams, 2000. Gum Arabic. In: *Handbook of Hydrocolloids*, Phillips, G.O. and P.A. Williams (Eds.). CRC Press, Boca Raton, FL. USA., pp: 155-168.
- Radman, R., T. Saez, C. Bucke and T. Keshavarz, 2003. Elicitation of plants and microbial cell systems. *Biotechnol. Applied Biochem.*, 37: 91-102.
- Rahim, A.H., E.C. van Ierland and J. Wesseler, 2007. Economic incentives for abandoning or expanding gum arabic production in Sudan. *For. Policy Econ.*, 10: 36-47.
- Randall, R.C., G.O. Phillips and P.A. Williams, 1989. Fractionation and characterization of gum from *Acacia senegal*. *Food Hydrocolloids*, 3: 65-75.
- Seif El Din, A., 1995. Gum arabic production in Sudan. Proceedings of the Agricultural Insurance Workshop in Sudan, Sept. 27-29, Sheikan Insurance and Reinsurance Co. Ltd., Sudan, pp: 15-15.
- Suliman, S.M., A.M. Hamdouk and M.B. Elfaki, 2000. G.A. fiber as a supplement to low protein diet in chronic renal failure patients. Proceedings of the 17th Conference of Sudan Association of Physicians, March 21-23, Friendship Hall, Khartoum, Sudan.
- Tolunay, A., A. Akyol and M. Ozcan, 2008. Usage of trees and forest resources at household level: A case study of Asagi Yumrutas village from the West Mediterranean region of Turkey. *Res. J. Forestry*, 2: 1-14.
- Verbeken, D., S. Dierckx and K. Dewettinck, 2003. Exudate gums: Occurrence, production and applications. *Applied Microbiol. Biotechnol.*, 63: 10-21.
- Vernon-Carter, E.J., C.I. Beristain and R. Pedroza-Islas, 2000. Mesquite gum (*Prosopis* gum). *Dev. Food Sci.*, 41: 217-238.