



International Journal of
Dairy Science

ISSN 1811-9743



Academic
Journals Inc.

www.academicjournals.com

Identification of Egyptian Goat α S1-casein Alleles Using PCR-RFLP Method

Sahar Ahmed

Department of Cell Biology, National Research Center, Dokki, Giza, Egypt

Abstract: This study aimed to detect the allelic frequency and identification of the genetic variants of α S1-casein gene in Egyptian goat using PCR-RFLP method. Forty-four animals used for detecting the presence of 10 polymorphisms of goat α S1-casein. The PCR products were digested by restriction enzyme *XmnI* and resolved using gel electrophoresis. Present results showed that Egyptian goat breeds carry low percentage of homozygous strong alleles A (4.5%) and B (6.8%) of milk casein used for cheese industry. The mild allele D recorded very low percentage 2.3%. This allele associated with medium level of milk protein favorable for allergic nutritional purposes. In conclusion, the allele frequencies and identification for animals α S1-casein are important for identifying animals carrying strong and mild alleles needed for improving a given trait depend on the frequency of favorable allele. The more will gained by selecting for strong and mild alleles of α S1-casein. So that, the breeding strategies should be changed toward milk protein alleles required for economic milk industry.

Key words: Goat, polymorphism, PCR-RFLP, α S1-casein

Introduction

The fluid milk and other dairy products have been essential components of the human diet in a variety of cultures, mainly because the high quality of the protein contained in milk and the variety of minerals and vitamins. It is known that differences in the primary structure of milk proteins may effect, directly or indirectly, milk properties (Grosclaude, 1988). Little attention has been paid to the possibility of changing the manufacturing properties of milk through genetic selection, particularly when compared with what has been achieved in increased genetic potential for milk yield.

The relative amounts of the four casein fractions (α S1, β , α S2 and K) affect the physicochemical, nutritional and technological properties of milk of domestic ruminants (Ramunno *et al.*, 2000). The first description of polymorphism in milk protein was reported for two variants β -lactoglobulin in cow's milk (Aschaffenburg and Drewery, 1955). Since then, many other polymorphic have been characterized, both in caseins and milk whey protein, especially over the last decade, owing to the improvement of both biochemical and molecular techniques.

The molecular analysis of the goat genome is currently carried out using specific goat sequences as well as heterologous ones isolated from cattle and sheep (Vaiman *et al.*, 1996). Deoxyribonucleic Acid (DNA) based markers have a number of favorable characteristics which represent ideal tool for studying the genetic quantitative traits (Ajmone-Marsan *et al.*, 2001).

The goat α S1-casein (CSN1S1) locus is characterized by at least 13 alleles associated with four different levels synthesis: high, medium, low and null (Martin *et al.*, 1999). The goat homozygous for alleles associated with a high content of α S1-casein produce milk characterized by a minor diameter of micelles, significantly higher percentage of protein, fat, total calcium and better parameters for curd firming time, curd firmness and cheese yield compared with goats homozygous for alleles associated with low or intermediate content (Martin *et al.*, 1999; Remeuf *et al.*, 1989; Manfredi *et al.*, 1993; Remeuf *et al.*, 1993; Grosclaude *et al.*, 1994; Pirisi *et al.*, 1994; Remeuf *et al.*, 1995; Lamberet *et al.*, 1996).

This study aimed to detect the allelic frequency and identification of the genetic variants of α S1-casein gene in Egyptian goat using PCR-RFLP method.

Materials and Methods

Genomic DNA Extraction

Genomic DNA was extracted from whole blood by phenol-chloroform method with minor modification (John *et al.*, 1991). Ten milliliter of blood taken on EDTA was mixed with 25 mL cold sucrose-triton and the volume was completed to 50 mL by autoclaved double distilled water. The solution was mixed well and the nuclear pellet was obtained by spin and discarding the supernatant. The nuclear pellet was suspended in lysis buffer (10 mM Tris base, 400 mM NaCl and 2 mM sodium EDTA) pH 8.2, with 20% Sodium Dodecyl Sulfate (SDS) and proteinase K (10 mg mL⁻¹) and incubated overnight in a shaking water-bath at 37°C.

Nucleic acids were extracted once with phenol, saturated with Tris-EDTA (TE) buffer (10 mM Tris, 10 mM NaCl and 1mM EDTA), followed by extraction with phenol-chloroform-isoamyl alcohol (25:24:1) until there was no protein at the interface. This was followed by extraction with chloroform-isoamyl alcohol (24:1).

To each extraction, equal volumes of the solvent were added, followed by thorough mixing and centrifugation for 10 min at 2000 rpm. The top layer was carefully transferred to another Falcon tube for the next extraction. To the final aqueous phase, 0.1 volume of 2.5 M Na acetate and 2.5 volume of cold 95% ethanol were added. The tubes were agitated gently to mix the liquids and a fluffy white ball of DNA was formed. The DNA was picked up with a heat-sealed Pasteur pipette and washed briefly in 70% ethanol.

The DNA was finally dissolved in an appropriate volume of 1X TE buffer. DNA concentrations were determined and diluted to the working concentration of 50 ng µL⁻¹, which is suitable for polymerase chain reaction.

Polymerase Chain Reaction (PCR)

The primer used in this study was synthesized from published nucleotide sequence of goat α S1-casein gene (forward: 5' TTCTAAAAGTCTCAGAGGCAG 3'), (reverse: 5' GGGTTGATAGCCTTGTATGT 3') (Ramunno *et al.*, 2000). Its PCR product enable us to detect 10 genetic variants of α S1-casein after digestion with a specific restriction enzyme.

A PCR cocktail consists of 1.0 (µM upper and lower primers and 0.2 mM dNTPs, 10 mM Tris (pH 9), 50 mM KCl, 1.5 mM MgCl₂, 0.01% gelatin (w/v), 0.1% Triton X-100 and 1.25 units of Taq polymerase. The cocktail was aliquot into tubes with 100 ng DNA of goat. The reaction mixture was overlaid with sterile mineral oil and is run in a Coy Temp Cycler II (Coy Corporation, MI, USA) the amplification protocol consisted of 30 cycles of 94°C for 1 min, 60°C for 1 min and 72°C for 2 min.

RFLP and Agarose Gel Electrophoresis

Twenty microliter of PCR product were digested with 10 units of restriction enzyme *XmnI* used in a final reaction volume 25. The reaction mixture was incubated at 37°C in water bath over night. After restriction digestion, the restricted fragments were analyzed by electrophoresis on 2.5% agarose/1X TBE gel stained with ethidium bromide. The 100 bp ladder was used as molecular size marker. The bands were visualized under UV light and photographed with yellow filter on black and white film.

Results and Discussion

Forty four animals used for detecting the presence of 10 polymorphisms of goat α S1-casein. The PCR products were digested by restriction enzyme *XmnI* and resolved using gel electrophoresis. The restriction fragment length polymorphism determined by *XmnI* allowed to identify the genotypes a/a at 150 bp, b/b at 161 bp, c/c at 212 bp, d/d at 223 bp. The presence of a/b should be illustrated

Table 1: Genotype frequencies of α S1-casein after XmnI digestion of fragment obtained by PCR products of Egyptian goat DNA

Genotype	Frequency (%)	RFLP (bp)	Genotype	Frequency (%)	RFLP (bp)
a/a	4.5	150	a/c	40.9	150+212
b/b	6.8	161	a/d	-----	150+223
c/c	-----	212	b/c	-----	161+212
d/d	2.3	223	b/d	45.5	161+223
a/b	-----	150+161	c/d	-----	212+223

Table 2: Genotype frequencies of α S1-casein among different Egyptian goat breeds

Breed	No. of animals	a/a (%)	b/b (%)	d/d (%)	a/c (%)	b/d (%)
Damascus	12	16.7	16.7	8.3	33.3	25.0
Baladi	11	---	---	---	90.9	9.1
Sahrawi	11	---	---	---	27.3	72.3
Zarabi	10	---	10	---	10	80.0




Fig. 1: DNA electrophoretic pattern obtained after digestion of PCR amplified of goat casein α 1 (CSN1S1) products with XmnI, M: 100-bp ladder marker, Lane 1: homozygous a/a genotype, Lane 2: homozygous b/b genotype, Lane 3: heterozygous b/d genotype, Lane 4: heterozygous a/c genotype




Fig. 2: DNA electrophoretic pattern obtained after digestion of PCR amplified of goat casein α 1 (CSN1S1) products with XmnI, M: 100-bp ladder marker, Lane 1: heterozygous a/c genotype, Lane 2: heterozygous b/d genotype, Lane 3: homozygous d/d genotype

by two fragments at 150 and 161 bp, a/c by two fragments at 150 and 212 bp, a/d by two fragments at 150 and 223 bp, b/c by two fragments at 161 and 212 bp, b/d by two fragments at 161 and 223 bp, c/d by two fragments at 212 and 223 bp (Table 1).

Among different breeds (Table 2), the results of genetic variants showed that the Damascus breed has five different genotypes a/a, b/b, d/d, a/c, b/d (Fig. 1 and 2), the Sahrawi breed has a/c with high frequency 90.9%, while the Baladi breed carries two different genotypes with different frequencies (Table 2). In Zarabi breed, the genotype b/d is the most common by 80%.

In caprine breeds, four types of alpha S1-casein alleles are identified and associated with various amounts of protein in milk. The A, B and C are strong alleles associated with a high level of α S1-casein in milk (3.5 g L^{-1}). The D and E alleles with a medium level (1.1 g L^{-1}), the F and G alleles with a low level (0.45 g L^{-1}). The contribution of null allele is very low or non existent of α S1-casein (Ramunno *et al.*, 2000). The quantity of total caseins in caprine milk is positively correlated with the amount of α S1-casein. Therefore, milk from animals possessing strong alleles contain significantly more total caseins than milk from animals without those alleles. This is important because animals with mild alleles can be employed to produce milk for allergic subjects, while the other animals can be used to produce milk for the dairy industry (Martin *et al.*, 1999; Leroux *et al.*, 1990; Roncada *et al.*, 2002).

Present results revealed that Egyptian goat breeds showed low percentage of homozygous strong alleles A (4.5%) and B (6.8%) of milk casein used for cheese industry. The mild allele D recorded very low percentage 2.3%. This allele associated with medium level 1.1 g L^{-1} of milk protein favorable for allergic subject (nutritional purposes) especially the infant diet where the goat milk with low casein is reported less allergenic than cow's milk (Roncada *et al.*, 2002). These results indicated that alleles require for economic milk industry present with low percentage in Egyptian goat breeds.

In conclusion, the alleles frequencies and identification for animals α S1-casein are important for identifying animals carrying the strong and mild alleles needed for improving a given trait depend on the frequency of favorable allele. The more will gained by selecting for strong and mild alleles of α S1-casein. So that, the breeding strategies should be changed toward milk protein alleles required for economic milk industry.

Acknowledgments

Especial thanks to Prof. Dr. Soheir M. El Nahas and Dr. Othman E. Othman for their support. This research was conducted as a part of the project Genetic polymorphism of some genes related to quantitative traits in different breeds of goat in Egypt (2002-2005), which was supported from the National Research Center (Egypt).

References

- Ajmone-Marsan, P., R. Negrini, P. Crepaldi, E. Milanese, C. Gorni, A. Valentini and M. Cicogna, 2001. Assessing genetic diversity in Italian goat populations using AFLP markers. *Anim. Gen.*, 32: 281-288.
- Aschaffenburg, R. and J. Drewery, 1955. Occurrence of different β -lactoglobulins in cow's milk. *Nature*, 176: 218-219.
- Grosclaude, F., 1988. Le polymorphisme genetique des principales lactoproteins bovines. *INRA Prod. Anim.*, 1: 5-17.
- Grosclaude, F., G. Ricordeau, P. Martin, F. Remeuf, L. Vassal and J. Bouillon, 1994. Du gene au fromage: Le polymorphisme de la caseine α S1 caprine, ses effects, son evolution. *INRA Prod. Anim.*, 7: 3-9.
- John, S., G. Weitzner, R. Rozen and C. Scriver, 1991. A rapid procedure for extracting genomeic DNA from leukocytes. *Nucleic Acid Res.*, 19: 408-412.
- Lamberet, G., C. Degas, A. Delacroix-Buchet and L. Vassal, 1996. Effect of characters linked to A and F caprine α S1-casein alleles on goat flavours: Cheese making with protein-fat exchange. *Lait*, 76: 349-61.
- Leroux, C., P. Martin, M. Mahe, H. Leveziel and J. Mercier, 1990. Restriction fragment length polymorphism identification of goat alpha S1-casein alleles: A potential tool in selection of individuals carrying alleles associated with a high level protein synthesis. *Anim. Gen.*, 21: 341-51.

- Manfredi, E., M. Barbieri, J. Bouillon, A. Piacere, M. Mahe, F. Grosclaude and G. Ricordeau, 1993. Effects des variants de la caseine α S1 le performances Laitieres de chevres. *Lait*, 73: 567-72.
- Martin, P., M. Ollivier-Bousquet and F. Grosclaude, 1999. Genetic polymorphism of casein: A tool to investigate casein micelle organization. *Intl. Dairy J.*, 9: 163-71.
- Pirisi, A., P. Colin, F. Laurent, J. Scher and M. Parmentier, 1994. Comparison of milk composition, cheese making properties and textural characteristics of the cheese from two groups of goats with a high or low rate of α S1-casein synthesis. *Intl. Dairy J.*, 4: 329-45.
- Ramunno, L., G. Cosenza, M. Pappalardo, N. Pastore, D. Gallo, P. Di Gregorio and P. Masina, 2000. Identification of the goat CSN1S1 allele by means of PCR-RFLP method. *Anim. Gen.*, 31: 333-346.
- Remeuf, F., 1993. Influence du polymorphisme genetique de la caseins α S1 caprine sur les caracteristiques physico-chimiques et technologiques du lait. *Lait*, 73: 549-57.
- Remeuf, F., J. Lenoir and C. Duby, 1989. Etude des relations entre les caractéristiques physico-chimiques des laits de chvre et leur aptitude à la coagulation par la presure. *Lait*, 69: 499-518.
- Remeuf, F., I. Verdalet-Guzman and J. Lenoir, 1995. Technological adjustment of goat milk containing low synthesis-rate α S1-casein variants. *Intl. Dairy J.*, 5: 381-92.
- Roncada, P., A. Gaviraghi, S. Liberatori, B. Canas, L. Bini and G. Greppi, 2002. Identification of caseins in goat milk. *Proteomics*, 2: 723-726.
- Vaiman, D., L. Schibler, F. Bourgeois, A. Oustry, Y. Amigues and E.P. Cribiu, 1996. A genetic linkage map of the male goat genome. *Genetics*, 144: 279-305.