



International Journal of
Dairy Science

ISSN 1811-9743



Academic
Journals Inc.

www.academicjournals.com

Assessment of Freeze-Dried Hydrodistilled Extracts from Clove; Caraway and Coriander Herbs as Natural Preservatives for Butter Oil

Hanaa F.M. Ali

Department of Chemistry, Faculty of Science, Taif University,
P.O. Box 888, Kingdom of Saudi Arabia

Abstract: Antioxidant activities of Freeze-Dried Hydrodistilled (FDH) extract of three herbs; clove, (*Syzygium aromaticum* L.); caraway (*Carum carvil* L.) and coriander (*Coriandrum sativum* L.) was evaluated *in vitro* by iron reduction; iron chelation and inhibition of lipid peroxidation methods. Also, total phenols content and the extraction yield was determined. Furthermore, simple model of butter oil was designed for evaluate the produced extracts as natural preservatives. Acid value, peroxide value and thiobarbituric acid (TBA) test were determined for oxidation system of butter oil. Antioxidant activity of clove FDH extract was significantly higher than caraway and coriander extracts when evaluated in iron reduction; iron chelation and inhibition of lipid peroxidation methods. Based on acid value, peroxide value and thiobarbituric acid test, different FDH extracts exhibited antioxidant effect specially at rate of 400 ppm from individual extracts with order of clove>coriander >caraway. Also, different correlation relationships were recorded in present study.

Key words: Antioxidant, freeze-dried, hydrodistilled, caraway, clove, coriander, natural preservatives, butter oil

INTRODUCTION

Food quality decreasing occurs during processing and storage is related to oxidative processes which cause degradation affects on lipids, carbohydrates and protein these are often catalyzed by e.g., ferrous or copper (Halliwell, 1997). Synthetic antioxidants such as BHA (butylated hydroxyl anisole) or BHT (butylated hydroxyl toluene) usually used to decelerate these processes. These antioxidants suffer from the disadvantage that they are volatile and easily decompose at high temperatures. Additionally, it is still unclear whether chronic consumption can lead to health risks (Martinez-Tome *et al.*, 2001). Plant-derived food additives, especially polyphenolic compounds, have also been ascribed health-promoting properties, as for example in terms of prevention of chronic cardiovascular diseases (Harborne and Williams, 2000; Singh *et al.*, 2008).

Such food additives are required to be odour free and tasteless. Many herbs and spices, usually used as an excellent source of phenolic compounds which have been reported to serve as natural food preservatives due to their good antioxidant activity (Rice-Evans *et al.*, 1996; Zheng and Wang, 2001; Justescn and Knuthsen, 2001). However, herbs and spices usually contain essential oil which show antioxidant activity but also carry flavour. Thus, the preparation of hydrodistilled extracts is using to remove the intrinsic flavour from the plant material. Furthermore, use of an aqueous solvent prevent solubility problems and this avoids harmful residues from organic solvents. Also, hydrodistilled extracts may use in the functionalization of foods and beverages as source of phenolic compounds which has a health-promoting properties (Harborne and Williams, 2000; Hinneburg *et al.*, 2006; Ruberto *et al.*, 2000; Teissedre and Waterhouse, 2000).

Corresponding Author: Hanna F.M. Ali, Department of Biochemistry, Faculty of Agriculture, Cairo University,
P.O. Box 12613, Cairo, Egypt

Some studies concerned for evaluate antioxidant activity of FDH extracts of different herbs (Dorman *et al.*, 2003; Hinneburg *et al.*, 2006; El-Ghorab *et al.*, 2008), but there is no study about clove, caraway and coriander FDH extracts. In present study, FDH extracts were prepared from three aromatic plants; clove, (*Syzygium aromaticum* L.); caraway (*Carum carvil* L.) and coriander (*Coriandum satvium* L.). Also, total phenols content and extraction yields were determined and *in vitro* antioxidant activity (iron reduction; iron chelation and inhibition of lipid peroxidation) was studied in producing extracts. Furthermore, simple model of butter oil was designed for evaluate the produced extracts as natural preservatives.

MATERIALS AND METHODS

Materials

Dried plant materials were obtained from experimental station of Faculty of Pharmacy, Cairo University during 2008. Ultrapure water was prepared using a Millipore Milli-RO 12 plus system (Millipore Corp., Bedford, MA). All reagents and solvents were either of analytical or HPLC grade and were obtained from Sigma Chemical Co., (St. Louis, MO).

Preparation of Freeze-Dried Hydrodistilled Extracts (FDH Extract)

Hundred gram of ground plant material were suspended in 500 mL ultrapure water and hydrodistilled for 3 h in a hydrodistillation glass apparatus. This process was repeated for 2 h with fresh solvent and the combined aqueous extracts were then filtered, reduced in volume *in vacuo* (-45°C), freeze-dried and stored at -4°C. The extracts were dissolved in ultrapure water prior to use.

Determination of Total Phenols Content

The total phenols were estimated according to the Folin-Ciocalteu method (Singleton *et al.*, 1999). To 100 μ L sample were added 250 μ L of undiluted Folin-Ciocalteu-reagent. After 1 min, 750 μ L of 20% (w/v) aqueous Na₂CO₃ were added and the volume was made up to 5.0 mL with H₂O. The control contained all the reaction reagents except the extract. After 2 h of incubation at 25°C, the absorbance was measured at 760 nm and compared to a gallic acid calibration curve. Total phenols were determined as gallic acid equivalents (mg gallic acid/g extract) and the values are presented as means of triplicate analysis.

Measurement of Iron (III) to Iron (II)-Reducing Activity

The ability of the extracts to reduce iron (III) was assessed by the method of Benjakul *et al.* (2005). Three milliliter aliquot of each extract, was dissolved in water, was mixed with 5 mL of phosphate buffer (0.2 M, pH 6.6) and 5 mL of a 1% potassium hexacyanoferrate [K₃Fe(CN)₆] solution. After a 30 min incubation at 50°C, 5 mL of 10% trichloroacetic acid were added and the mixture was centrifuged for 10 min. Five milliliter aliquot of the upper layer was mixed with 5 mL of water and 1 mL of 0.1% aqueous FeCl₃ and the absorbance was recorded at 700 nm. Iron (III) reducing activity was determined as ascorbic acid equivalents (mmol ascorbic acid/g extract). The values are presented as the means of triplicate analysis.

Study the Inhibition of Linoleic Acid Peroxidation

The antioxidant activity was determined as the degree of inhibition on the haemoglobin-catalysed peroxidation of linoleic acid according to Kuo *et al.* (1999). Sample (20 μ L) were added to 0.5 mL of 0.05 M phosphate buffer (pH 7.0), containing 0.14% Tween 20 and 4 mM linoleic acid and then equilibrated at 37°C for 3 min. The peroxidation of linoleic acid in the above reaction mixture was initiated by adding 30 μ L of 0.035% haemoglobin (in water), followed by incubation at the same temperature for 10 min and stopped by adding 5 mL of 0.6% HCl (in ethanol). The hydroperoxide

formed was assayed according to a ferric thiocyanate method with mixing, in order, of 30% ammonium thiocyanate (0.2 mL) and 0.02 M ferrous chloride (0.1 mL). The absorbance (A_s) was measured at 480 nm after 5 min. The absorbance blank (A_0) was obtained without adding haemoglobin to the above reaction mixture; the absorbance of the control (A_{100}) was obtained by replacing the sample by buffer.

$$AA(\%) = \frac{1 - (A_s - A_0)}{A_{100} - A_0} \times 100 \quad (1)$$

The antioxidant activities of the samples were calculated as trolox equivalents (mg trolox/g extract) according to Eq. 1. The values are presented as the means of triplicate analysis.

Measurement of Iron (II) Chelation Activity

The chelation of iron (II) ions by the different extracts was carried out as described by Carter (1971). Two hundred microlitres of each extract were added to 100 μ L of 2.0 mM aqueous $FeCl_2$ and 900 μ L methanol. The controls contained all the reaction reagents except the extract or positive control substance. After a 5 min incubation, the reaction was initiated by added 400 μ L of 5.0 mM ferrozine. After a 10 min equilibrium period, absorbance at 562 nm was recorded. The iron chelation activities were calculated from the absorbance of the control (A_c) and of the sample (A_s) using Eq. 2 and expressed as Na_2EDTA equivalents (mg Na_2EDTA/g extract). The values are presented as the means of triplicate analysis.

$$\text{Inhibition (\%)} = \frac{A_c - A_s}{A_c} \times 100 \quad (2)$$

Preparation of Butter Oil

Fresh buffalo milk was pasteurized and separated into cream and skimmed milk using an Alfa Laval separator (AESC, Sweden). The cream was churned to obtain unsalted butter. The resultant butter was melted using moderate heat (55°C) and the clear butter oil layer was carefully decanted.

Chemical Analysis

Acid and peroxide values were determined according to the methods of the Association of Official Analytical Chemists (1980). The thiobarbituric acid (TBA) test was carried out according to the method described by Patton (1973).

Design of Oxidation System

The Butter Oil (BO) was distributed into eight portions (200 g each) in sterilized glass bottles. The first portion served as a control. BHT at rate of 200 ppm was added to the second portion. To the 3rd and 4th portions, FDH extract of clove was added at a rate of 200 and 400 ppm, respectively. To the 5th and 6th, FDH extract of caraway was added at a rate of 200 and 400 ppm, respectively. In the 7th and 8th, FDH extract of coriander was added at a rate of 200 and 400 ppm, respectively. Control and treated samples were placed in an incubator at $60 \pm 1^\circ C$ to accelerate the auto-oxidation of butter oil, (Thompson, 1960). The experimental period was terminated when an objectional odour and high lipid rancidity values were obtained with the samples after 28 days.

Statistical Analysis

All statistical analysis were carried out using iner-STAT-a (Vargas, Mexico) as add-in for Microsoft Excel 2007. Analysis of Variance (ANOVA) was followed by Tukeys' pairwise comparison test at a level of $p < 0.05$ for the determination of significant differences between treatments and control.

RESULTS AND DISCUSSION

Extraction Yield and Total Phenols

Table 1 shows the extraction yields and total phenols, for the FDH extract of selected herbs. The extraction yields was 245; 210 and 130 mg g⁻¹ for clove; coriander and caraway, respectively. Significant association found between the extraction yields and total phenols and the results from the different antioxidant assays. Phenolic substances which investigated by the Folin-Ciocalteu method (mg gallic acid/g extract) have been shown to be responsible for the antioxidant activity of plant materials. Phenolic substances were found in the order clove>coriander>caraway (p<0.05). The content of total phenolic shows a good correlation with the iron reduction assay (R² = 0.8871, p<0.05) and inhibition of lipid peroxidation assay (R² = 0.7327, p<0.05), while no correlation with the iron (III) to iron (II)-reducing assay or iron (II) chelation assay. The total phenols/extractable compounds ratio was 33.7% for clove; 28.6% for coriander and 18.4% for caraway.

Iron (III) to Iron (II)-Reducing Activity

Fe (III) reduction is often used as an indicator of electron-donating activity, which is an important mechanism of phenolic antioxidant action (Yildirim *et al.*, 2001). The iron (III) to iron (II)-reducing activity is expressed as ascorbic acid equivalents (mmol ascorbic acid/g sample). The ascorbic acid equivalent in clove extract was significantly higher than for the other extracts (p<0.05). Ascorbic acid equivalents was 0.78 mmol ascorbic acid/g for clove extract; 0.32 mmol ascorbic acid/g for coriander extract and 0.13 mmol ascorbic acid/g for caraway extract.

Peroxidation of Linoleic Acid

In the haemoglobin-catalysed peroxidation of linoleic acid assay, linoleic acid served as a lipid model. Peroxidation was induced by haemoglobin and the damage was assayed following the thiocyanate and Trolox was used as reference substance (Kuo *et al.*, 1999). Results indicated that clove was significantly better inhibitors of lipid peroxidation than the other extracts (p<0.05), Trolox equivalents ranked from 443±12.11 mg trolox/g extract for clove to 93±4.21 mg trolox/g extract for caraway. The antioxidant activity in this assay correlated well only with iron reduction (R² = 0.8958, p<0.05).

Iron (II) Chelation

Foods contaminated with transition metal ions which may be introduced by processing methods, these ions play an important role as catalysts of oxidative processes, leading to the formation of hydroxyl radicals and hydroperoxide decomposition reactions via., Fenton chemistry (Halliwell, 1997). These processes can be delayed by iron chelation and deactivation. Therefore, the ability of the extracts to chelate iron (II) ions was evaluated and expressed as mg Na₂EDTA/g extract. Clove show the significantly (p<0.05) best iron chelation (234±5.21 mg Na₂EDTA/g extract), followed by coriander (143±2.71 mg Na₂EDTA/g extract) and (87±1.43 mg Na₂EDTA/g extract) for caraway. No significant correlation was found between the iron chelation ability of the extracts and the other antioxidant activities.

Effect of FDH Extract of Different Plants on Lipid Rancidity

In the present study the simple model system consists of butter oil mixed with FDH extract of selected herbs were designed to study oxidative process during storage. BHT and BHA are added as

Table 1: Extraction yields and total phenols for hydrodistilled extracts

Herbs	Extraction yield (mg g ⁻¹)	Total phenols (mg GA g ⁻¹)
Clove	245	85.60±9.50
Coriander	210	60.21±6.40
Caraway	130	35.32±8.10

Table 2: Acid values for Butter Oil (BO) mixed with BHT and different FDH extract during storage

Treatments	Storage period (days)									Mean
	0	3	6	10	14	18	22	24	28	
Control	0.45	0.51	0.53	0.54	0.52	0.64	0.71	0.75	0.74	0.610
BO+BHT at 200 ppm	0.44	0.54	0.55	0.57	0.53	0.50	0.53	0.56	0.55	0.541
BO+clove FDH 200 ppm	0.48	0.50	0.54	0.51	0.52	0.54	0.51	0.53	0.50	0.526
BO+clove FDH 400 ppm	0.42	0.54	0.57	0.57	0.57	0.54	0.51	0.54	0.49	0.530
BO+coriander FDH 200 ppm	0.43	0.53	0.58	0.57	0.54	0.56	0.69	0.68	0.64	0.590
BO+coriander FDH 400 ppm	0.44	0.56	0.58	0.54	0.64	0.60	0.59	0.58	0.57	0.570
BO+caraway FDH 200 ppm	0.41	0.52	0.58	0.55	0.60	0.62	0.59	0.58	0.60	0.570
BO+caraway FDH 400 ppm	0.47	0.50	0.54	0.56	0.55	0.58	0.61	0.62	0.60	0.580

Acid value is expressed as mg KOH /1 g butter oil




Fig. 1: Effect of different FDH extract on Butter Oil (BO) oxidative rancidity during storage

common effective synthetic antioxidants to food products in the range of 100-400 ppm (Allen and Hamilton, 1999). Experiments were conducted using BHT at 200 ppm in order to compare the anti-oxidative activity of different FDH extract towards butter oil rancidity. A high storage temperature (60±1°C) was chosen to accelerate of BO oxidation due to a low degree of unsaturation of butter oil (Thompson, 1960). The experimental period was terminated when an objectional odour and high lipid rancidity values were obtained with the samples after 28 days.

Butter Oil System

Acid Values (AV) of butter oil (control) and mixed with different FDH extract at different rate and BHT during storage is shown in Table 2 and Fig. 1. The AV results indicate that different FDH extract exhibited antioxidant effect specially at rate of 400 ppm from different extracts with order of clove > coriander > caraway. Changes in the PV of butter oil mixed with different FDH extract at different rate and BHT during storage are shown in Fig. 1. The PV values of BO (control) and mixed with different extracts were very low at zero time and gradually increased with storage time. BHT produced significantly lower PV of butter oil when compared with other treatments. The addition of different extracts was effective as antioxidant when compared with the control experiment and in the same order; clove > coriander > caraway. The changes in the secondary oxidation products of BO are shown in Fig. 2, a result of the addition of different FDH extract based on the data of TBA values. Addition of BHT at 200 ppm to butter oil caused very low increase in TBA values over time. On the other hand, the systems containing different FDH extract at different rate significantly linear reduce the formation of secondary oxidation products in comparison with the control experiment. The inhibitory effect of antioxidants against oxidation process in butter oil has been attributed to their donation of electrons or hydrogen atoms from phenolic hydroxyl groups to butter oil containing free




Fig. 2: Effect of different FDH extract on Butter Oil (BO) secondary oxidation products

radicals and to the formation of stable free radicals which do not initiate or propagate further oxidation of oils (Farag *et al.*, 1993; Wollgast and Anklam, 2000). The good correlation between the results from total phenols analysis and the antioxidative assays has been earlier reported by Zheng and Wang (2001) and Dorman *et al.* (2003, 2004).

In complex systems, such as food and food preparations, various different mechanisms may contribute to oxidative processes, such as in Fenton reactions, where transition metal ions play a vital role, different reactive oxygen species might be generated and various target structures such as lipids, proteins and carbohydrates, can be affected. Therefore, it is important to characterize the extracts by a variety of antioxidant assays (Halliwell, 1997; El-Ghorab *et al.*, 2008; Ollanketo *et al.*, 2002).

CONCLUSION

The result of the present study showed that the different FDH extracts which contain highest amount of phenolic compounds, exhibited the greatest antioxidant activity. All of the extracts in this research exhibited different extent of antioxidant activity. The catalytic effectiveness of the added materials can be ranked according to inhibition power as BHT (200 ppm) > clove > coriander > caraway > control. Plant extracts might substitute synthetic food antioxidants, which may influence human health when consumed chronically (Martinez-Tome *et al.*, 2001).

ACKNOWLEDGMENT

This study funded by Taif University, Faculty of Science, Chemistry Department, Kingdom of Saudi Arabia.

REFERENCES

- Allen, C. and J. Hamilton, 1999. Rancidity in Foods. 3rd Edn., Applied Science Publishers, London.
- Association of Official Analytical Chemists, 1980. Official Methods of Analysis. 15th Edn., Association of Official Analytical Chemists, W. Horowitz, Washington DC., ISBN: 0935584145.
- Benjakul, S., W. Lertittikul and F. Bauer, 2005. Antioxidant activity of Maillard reaction products from a porcine plasma protein-sugar model system. *Food Chem.*, 93: 189-196.
- Carter, P., 1971. Spectrophotometric determination of serum iron at the submicrogram level with a new reagent (ferrozine). *Anal. Biochem.*, 40: 450-458.

- Dorman, H.J.D., M. Kosar, K. Kahlos, Y. Holm and R. Hiltunen, 2003. Antioxidant properties and composition of aqueous extracts from *Mentha* species, hybrids, varieties and cultivars. *J. Agric. Food Chem.*, 51: 4563-4569.
- Dorman, H.J., O. Bachmayer, M. Kosar and R. Hiltunen, 2004. Antioxidant properties of aqueous extracts from selected Lamiaceae species grown in Turkey. *J. Agric. Food Chem.*, 52: 762-770.
- El-Ghorab, H., A. Mansour and K. El-massry, 2008. Effect of extraction methods on the chemical composition and antioxidant activity of Egyptian marjoram (*Majorana hortensis* Moench). *Flav. Fragr. J.*, 19: 54-61.
- Farag, S., A. Hassan and M. Ali, 1993. Bees wax and its unsaponifiables as natural preservative for butter and cottonseed oils. *Grasas Y Aceites*, 44: 183-189.
- Halliwell, B., 1997. Antioxidants: The basics-what they are and how to evaluate them. *Adv. Pharm.*, 38: 3-20.
- Harborne, J.B. and C.A. Williams, 2000. Advances in flavonoid research since 1992. *Phytochemistry*, 55: 481-504.
- Hirneburg, I., H.J.D. Dorman and R. Hiltunen, 2006. Antioxidant activities of extracts from selected culinary herbs and spices. *Food Chem.*, 97: 122-129.
- Justesen, U. and P. Knuthsen, 2001. Composition of flavonoids in fresh herbs and calculation of flavonoid intake by use of herbs in traditional Danish dishes. *Food Chem.*, 73: 245-250.
- Kuo, J., D. Yeh and B. Sun Pan, 1999. Rapid photometric assay evaluating antioxidative activity in edible plant material. *J. Agric. Food Chem.*, 47: 3206-3209.
- Martinez-Tome, M., A. Jimenez, S. Ruggieri, N. Frega, R. Strabbioli and M. Murcia, 2001. Antioxidant properties of Mediterranean spices compared with common food additives. *J. Food Prot.*, 64: 1412-1419.
- Ollanketo, M., A. Peltoketo, K. Hartonen, R. Hiltunen and M. Riekkola, 2002. Extraction of sage (*Salvia officinalis* L.) by pressurized hot water and conventional methods: Antioxidant activity of the extracts. *Eur. Food Res. Technol.*, 215: 158-163.
- Patton, S., 1973. Malonaldehyde, lipid oxidation and the thiobarbituric acid test. *J. Am. Oil Chem. Soc.*, 51: 114-120.
- Rice-Evans, C., N.J. Miller and G. Paganga, 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.*, 20: 933-956.
- Ruberto, G., M.T. Baratta, S.G. Deans and H.J. Dorman, 2000. Antioxidant and antimicrobial activity of *Foeniculum vulgare* and *Crithmum maritimum* essential oils. *Planta Med.*, 66: 687-693.
- Singh, G., S. Kiran, P. Marimuthu, M.P. de Lampasona, C.S. de Heluani and C.A.N. Catalán, 2008. Chemistry, biocidal and antioxidant activities of essential oil and oleoresins from *Piper cubeba* (seed). *Int. J. Essent. Oil Therap.*, 2: 50-59.
- Singleton, V., R. Orthofer and R. Lamuela-Raventó, 1999. Analysis of Total Phenols and Other Oxidation Substrates and Antioxidants by Means of Folin-Ciocalteu Reagent. In: *Oxidants and Antioxidants*, Packer, L. (Ed.). *Methods in Enzymology*, Academic Press, New York, pp: 152-178.
- Teissedre, P. and A. Waterhouse, 2000. Inhibition of oxidation of human low-density lipoproteins by phenolic substances in different essential oils varieties. *J. Agric. Food Chem.*, 48: 3801-3805.
- Thompson, W., 1960. Survey of Accelerated Test for Determining the Stability of Oils and Fats. In: *Laboratory Handbook for Oil and Fat Analysis*, Cooks, L.V. and C. Van Rede (Eds.). Academic Press, London, ISBN: 2906941.
- Wollgast, J. and E.E. Anklam, 2000. Review on polyphenols in theobroma cacao: Changes in composition during the manufacture of chocolate and methodology for identification and quantification. *Food Res. Int.*, 33: 423-447.
- Yildirim, A., A. Mavi and A. Kara, 2001. Determination of antioxidant and antimicrobial activities of *Rumex crispus* L. extracts. *J. Agric. Food Chem.*, 49: 4083-4089.
- Zheng, W. and S.Y. Wang, 2001. Antioxidant activity and phenolic compounds in selected herbs. *J. Agric. Food Chem.*, 49: 5165-5170.