



International Journal of
Dairy Science

ISSN 1811-9743



Academic
Journals Inc.

www.academicjournals.com

Characterisation of Yeasts Isolated from Artisanal Turkish Dairy Products*

G. Kavas, O. Kinik, H. Uysal, S. Kilic, N. Celikel and N. Akbulut
Department of Dairy Technology, Faculty of Agriculture,
Ege University, 35040, Izmir, Turkey

Abstract: Economic and sensorial losses of dairy products due to spoilage by yeast have been increasing in Turkey because of poor hygienic conditions during processing and to shorten the anticipated shelf life of products and/or also reduced use of preservatives that do not strictly control the growth of these organisms. This study reports the results of a survey of yeast species in samples of cow's, ewe's and goat's milk products collected in some regional bazaar of Izmir, Western Turkey. Yeasts were isolated from white pickled cheese, tulum cheese and kashkaval cheese, yoghurt and strained yoghurt respectively. The most frequently occurring yeast species on dairy products were *Candida* sp., *Saccharomyces* sp., *Kluyveromyces* sp. and *Trichosporon* sp. respectively. Even though yeasts are considered as environmental contaminants, the occurrence of some of them in dairy products at high levels could represent a potential risk for human health especially for immuno compromised patients.

Key words: Yeasts, identification spoilage, dairy products

Introduction

The main group of microorganisms generally associated with dairy products is covered by lactic acid bacteria although recently, it is well recognized that yeasts isolated from especially cheese and functional dairy products play a significant role in their ripening and developing some aroma components (Jacobsen and Narvhus, 1996; Pereira-Dias *et al.*, 2000; Fadda *et al.*, 2001; Minervini *et al.*, 2001). Yeasts in dairy products may interact with other microorganism in three different ways:

- They may inhibit or eliminate microorganisms which are undesired because they cause quality defects or potentially pathogenic characters
- They may inhibit the starter culture activity or
- They may positively associate with the fermentation or maturation process by supporting the function of the starter culture (Jacobsen and Narvhus, 1996; Loureiro and Querol, 1999).

On the contrary, yeast can have a negative effect as spoilage organisms in fermented milk products and cheeses (Pereira-Dias *et al.*, 2000; Minervini *et al.*, 2001). The main defects caused by spoilage yeasts are fruity, bitter or yeasty off-flavours, gas production, discoloration changes and a softening of product texture (Corbo *et al.*, 2001). Whereas, the use of yeasts in dairy industry could determine potential advantages covering production of flavour components and acceleration of ripening process of some cheese species by means of its lipolytic and proteolytic properties, fermentation ability of

Corresponding Author: Dr. G. Kavas, Department of Dairy Technology, Faculty of Agriculture, Ege University, 35040, Izmir, Turkey Tel: + 90 232 388 40 00/1420 Fax: +90 232 388 18 64

*Originally Published in *International Journal of Dairy Science*, 2006

lactose, assimilation of lactate and positively interactions with the starter culture (Jacobsen and Narvhus, 1996; Fleet and Mian, 1987; Fleet, 1990; Roostita and Fleet, 1996).

In this study, we were undertaken to investigate the frequency of occurrence and types of yeasts associated with different animal origin dairy products located in some regions of Izmir.

Materials and Methods

The study was carried out on 106 samples of dairy products, representing manufactured from usually cow's milk and cow's, ewe's and goat's milk mixture. The samples were collected from grocery stores and bazaar in Izmir and Aydin two regions of western Turkey. Samples originated from different animal origin were distributed as follows according to type of dairy products:

- Forty nine different time ripened products; 15 white pickled cheese, 15 kashkaval cheese, 19 tulum cheese canned and strained
- Fifty seven unripened dairy products (no ripening required) 30 strained yoghurt, 27 set type yoghurt. Also quide by date coding we chose for analysis only yoghurt and stained yoghurt samples that had not been manufactured more than 5 days previously.

The samples were transported in an ice box to the laboratory where they were immediately analyzed (Suriyarachi and Fleet, 1981).

Identification of Yeasts

Yeasts isolation was performed according to ISO 8261. From each sample, 20 g of the different dairy products including yoghurt, strained yoghurt and cheese species were diluted in 180 mL peptone-water and homogenized in Stomacher Lab-Blender 400. Ten fold dilutions were prepared in quarter strength ringer's solution and suitable aliquots were plated on yeast glucose agar (Merck 0500 KgaA Darmsadt) with chloramphenicol (100 mg L^{-1}) added and plates were incubated at $25 \pm 1 \text{ C}^\circ$ for 3-5 days. For the identification, at least three colonies of each different morphological type were selected from primary cultures. After purification on yeast glucose agar, the yeasts were grown on the same media and kept at 4 C° until they were identified.

The colonie isolates were characterized to (Vander and Yarrow, 1984). Identification was carried out by comparing the test datas with (Lodder, 1970). Microscopic observations of cell morphology and the production of ascospores, arthrospores and true hyphae were used to supplement of biochemical tests (Deak, 1986; Deak and Beuchat, 1987).

Results and Discussion

Yeasts in fermented dairy products and especially cheese varieties are assumed as insignificant at the earlier stages of storage life, but are important component of the microflora of many, if not all, cheese varieties and some fermented products in later stages, being present as natural contaminants in the curd and the end product during maturation and storage (Corbo *et al.*, 2001; Welthagen and Viljoen, 1998). The environment during ripening of the cheese and storage of fermented products consist of a low pH, moisture content, elevated salt concentration and low temperature which are often very favorable and selective for yeast growth (Vanden and Jakobsen, 1998; Viljoen and Greyling, 1995; Wojtatowicz *et al.*, 2001). The increase in the number of yeasts corresponds with the depletion in lactose content and the simultaneous stabilization of lactic acid bacteria numbers used as a starter (Jacobsen and Narvhus, 1996; Fleet, 1990; Gadaga *et al.*, 2000).

Table 2: Properties of those yeast species that exhibited growth in yoghurt strained yoghurt samples

Species identified	No. of species	Fermentation					Assimilation					Morphologic characteristics								
		Gl	Ga	L	M	Ra	Ga	Tr	In	L.A	Me	S	Ga	L.A	KNO ₃	Urea	P	A	G.in.B.	Gr at 37°C
<i>Saccharomyces cerevisiae</i>																				
<i>biovar I</i>	17	+	+	V	+	+	V	-	-	V	+	+	+	-	-	-	-	-	Films+Turbidity	-
<i>biovar II</i>	5	+	-	-	+	-	-	-	-	-	+	+	+	-	-	+	V	-	Turbidity	-
<i>biovar III</i>	1	+	+	-	-	+	+	-	-	-	V	V	+	+	-	-	V	-	Films	+
<i>P. farinosa</i>	3	+	-	-	-	+	-	-	-	-	-	+	+	-	-	+	-	-	Films+Turbidity	-
<i>P. anomala</i>	1	+	-	-	-	+	+	-	V	V	V	+	+	-	-	+	+	+	Films+Turbidity	+
<i>C. blankii</i>	1	-	-	-	+	+	+	-	V	-	-	+	+	+	-	-	+	-	Films+Slight turbidity	-
<i>C. lipolytica</i>	3	V	-	V	+	+	+	-	V	-	+	+	+	-	-	-	V	-	Films	-
<i>C. tropicalis</i>	2	+	V	V	+	V	V	-	V	-	+	V	+	-	+	+	+	V	Films	-
<i>C. kefir</i>	1	+	+	V	+	+	V	-	-	-	+	+	+	+	-	+	-	-	Films+Turbidity	-
<i>K. marxianus</i>	5	+	+	+	-	+	+	-	-	+	-	+	+	+	-	-	+	-	Turbidity	+
<i>var. lactis</i>																				
<i>K.marxianus</i>	3	+	+	+	V	+	+	-	-	+	V	+	+	+	-	-	+	-	Films+Turbidity	V
<i>var. marxianus</i>																				
<i>G. candidum</i>	7	+	-	-	-	-	+	V	-	-	-	+	+	+	-	-	V	+	Turbidity+White films	+
<i>T. cutaneum</i>	9	-	-	-	+	-	+	-	-	-	+	+	+	+	-	-	+	+	Turbidity+Sediment	-
<i>T.brassicae</i>	4	-	-	-	-	+	-	V	-	+	+	+	+	-	-	V	+	+	Turbidity+Sediment	-
<i>D.hansenii</i>	7	-	-	-	+	+	+	-	-	+	+	+	+	-	-	-	-	-	Films+Turbidity	+

*V: Variability, Gl: Glucose, Ga: Galactose, L: Lactose, M: Maltose, Ra: Rannose, Tr: Trehalose, In: Inocitol, L.A: Lactic Acide, Me: Mellibiose, S: Saccharose, P: Pseudomicelyum, A: Ascospore, G.in.B.: Growing in Broth, Gr at 37°C: Growth at 37°C

species, other fermented and unfermented dairy products (Fleet, 1990; Welthagen and Viljoen, 1998). In addition, the occurrence of *Kluyveromyces lactis* during ripening is probably related to the ability of these species to metabolize milk protein and fat and in some cases lactic acid (Fleet and Mian, 1987).

The occurrence of *Rhodotorula glutinis*, *Sporobolomyces roseus* in cheese has been reported previously (Fadda *et al.*, 2001; Fleet and Mian, 1987; Vanden and Jakobsen, 1998; Viljoen and Greyling, 1995). These yeast strains are considered to be contributing as common air or natural contaminants in the cheese before they were stored, although *Rhodotorula* sp. are able to grow at sub-zero temperatures (Deak, 1986; Welthagen and Viljoen, 1998). Also these results are in agreement with previous reports (Minervini *et al.*, 2001; Welthagen and Viljoen, 1998; Viljoen and Greyling, 1995). Moreover it can be said that, the presence some of large number of yeasts during the late stages of maturation suggests that some yeast species may play an important role in the ripening of cheese species especially tulum and Kashkaval cheese as observed in this study.

Yoghurt and Strained Yoghurt

Yeasts apparently are indigenous to yoghurt and strained yoghurt and their microbial flora form an interesting microbial ecosystem. The high acid content of yoghurts, which is enough to inhibit bacterial growth and the limited access of air to the yoghurt in the containers during cold storage make it a special microbial habitat that is most suitable for the growth of yeasts. Especially the high yeast counts in strained yoghurt produced by the traditional method directly after packaging suggest a high degree of contamination and mishandling of the product, as previously mentioned by Yamani and Abu-Jaber (1994). This problem occurred in yoghurt and skinned yoghurt samples could be associated to many factor such as the use of yoghurt starter cultures, yoghurt or both that were contaminated by yeasts or to the undesirable hygienic conditions prevailing during straining of the whey, especially resourced by the cleanliness of the bags and apparatus used. Lack of desirable hygienic practices during producing of fermented dairy products may also contribute to the contamination risk by yeasts.

Table 2 shows the yeasts found in yoghurt and strained yoghurt samples, the seven most commonly isolated species and the counts at which these species occurred. The isolates were distributed in seven genera as follows: *Saccharomyces* (23 isolates), *Trichosporon* (13 isolates),

Kluyveromyces (8 isolates), *Candida* (7 isolates), *Debaryomyces* (7 isolates), *Geotricum* (7 isolates), *Pichia* (3 isolates). According to data obtained all yeasts species, from yoghurt and strained yoghurt samples could be usually considered to be psychrophilic because of their ability to grow at refrigeration conditions. Especially the predominance of *S. cerevisiae* in strained yoghurt can be explained by its ability to ferment glucose and galactose and to assimilate glucose, which occurs in appreciable amounts in yoghurt whey as mentioned by Yamani and Abu-Jaber (1994). Also *S. cerevisiae* exhibited from other related dairy products, yoghurt and yoghurt starter cultures. The frequent isolations *Kluyveromyces marxianus* important criteria governing the growth of yeasts in yoghurts is the ability to grow well in cold storage and to ferment glucose, galactose and lactose (Table 2). These criteria are consistent with the storage of yoghurt types under refrigerated conditions and the presence of lactose and sucrose as the major carbohydrates. These findings about the isolations of the lactose fermenting yeast, *Kluyveromyces* spp, is in agreement with previous studies on yeasts in yoghurt starter culture (Suriyarachi and Fleet, 1981), yoghurt (Spillmann and Geiges, 1983), strained yoghurt (Yamani and Abu-Jaber, 1994), kefir grains (Engel *et al.*, 1986).

The remaining yeasts species occurred in this study were previously isolated and identified from different dairy products; *G. candidum*, *T. cutaneum* and *T. brassicae* were isolated from raw milk, quarg and cheese species (Fleet and Mian, 1987; Roostita and Fleet, 1996; Engel *et al.*, 1980; Engel, 1986). *C. blankii*, *C. lipolytica*, *C. tropicalis*, *C. kefir* were isolated from pasteurized milk, yoghurt, cheese (Fleet and Mian, 1987; Hatting and Viljoen, 2002). *Debaryomyces hansenii*, *Pichia farinosa*, *P. anomala* were isolated from yoghurt, labaneh, butter milk and cheese (Suriyarachi and Fleet, 1981; Yamani, and Abu-Jaber, 1994; Hatting and Viljoen, 2002; Mihyar *et al.*, 1997).

Conclusions

The occurrence of yeasts in dairy products is not usually uncommon. Mainly psychrotrophic yeasts are the main cause of spoilage of traditionally produced dairy products. Use of alternative producing techniques, instead of the traditional methods, may help to avoid the adverse effects of yeasts growth. The use of yeasts-free starter cultures, application of accurate hygienic measures during processing and packing and maintenance of an effective cold chain from production until consumption could help control spoilage yeasts. On the other hand, the presence of a large number of yeasts during the late stages of ripening suggests that some yeasts species may play an important role in the ripening of medium and long term ripened cheese varieties. Also the occurrence of toxigenic fungi (*Penicillium expansum*, *P. roquefortii* and *Aspergillus versicolor* etc.) from medium and long term ripened Turkish cheese varieties may be a reason for concern linked to the risk of occurring mycotoxin residues in the end product (Lopez-Diaz *et al.*, 1995).

References

- Brocklehurst, F. and B.M. Lund, 1985. Microbiological changes in cottage cheese varieties during storage at 7°C. *Food Microbiol.*, 2: 207-233.
- Corbo, M.R., R. Lanciotti, M. Albenzio and M. Sinigaglia, 2001. Occurrence and characterization of yeasts isolated from milks and dairy products of Apulia region. *Intl. J. Food Microbiol.*, 69: 147-152.
- Deak, T., 1986. A Simplified Schema for the Identification of Yeasts. *Methods for the Mycological Examination of Foods* King, A.D., J.I. Pitt, L.R. Beuchat and J.E.L. Coeey, (Eds.), Plenum. New York.

- Deak, T. and L.R. Beuchat, 1987. Identification of foodborne yeasts. *J. Food Prot.*, 50: 243.
- Engel, G., U. Krusch and M. Teuber, 1986. Microbiologische Zusammensetzung Von Kefir. I. Hefen. *Milchwissenschaft*, 41: 418.
- Engel, G., M. Teuber, G. Hoffmeisten and K.I. Neumann, 1980. Entwicklung von hefen und schimmelpilzen in speisequark. *Milchwissenschaft*, 35: 13.
- Engel, G., 1986. Hefen in silagen und rochmilch. *Milchwissenschaft*, 41: 633.
- Hatting, A.L. and B.C. Viljoen, 2002. Survival of dairy associated yeasts in yoghurt and yoghurt related products. *Food Microbiol.*, 19: 597-604.
- Fadda, M.E., V. Mossa, M.B. Pisano, M. Deplano and S. Cosentino, 2004. Occurrence and characterization of yeasts isolated from artisanal *Fiore sardo* cheese. *Intl. J. Food Microbiol.*, 95: 51-55.
- Fadda, M.E., S. Cosentino, M. Deplano and F. Palmas, 2001. Yeasts populations in *Sardinian Feta* cheese. *Intl. J. Food Microbiol.*, 69: 153-156.
- Faticenti, F., J.L. Bergere, P. Deiana and G.A. Farris, 1983. Antagonistic activity of *Debaryomyces hansenii* towards *Clostridium tyrobutyricum* and *C. butyricum*. *J. Dairy Res.*, 50: 449-457.
- Fleet, G. and M.A. Mian, 1987. The occurrence and growth of yeasts in dairy products. *Intl. J. Food Microbiol.*, 4: 145-155.
- Fleet, G.H., 1990. Yeasts in dairy products-a review. *J. Applied Bacteriol.*, 68: 199.
- Gadaga, T.H., A.N. Mutukumira and J.A. Narvhus, 2000. Enumeration and identification of yeasts isolated from Zimbabwean traditional fermented milk. *Intl. Dairy J.*, 10: 459-466.
- Hatting, A.L., and B.C. Viljoen, 2002. Survival of dairy associated yeasts in yoghurt and yoghurt related products. *Food Microbiol.*, 19: 597-604.
- Jacobsen, M. and J. Narvhus, 1996. Yeasts and their possible beneficial and negative effects on the quality of dairy products. *Intl. Dairy J.*, 6: 755-768.
- Lopez-Diaz, T.M., J.A. Santas, M. Prieto, M.L.G. Lopez and A. Otero, 1995. Mycoflora of a traditional Spanish blue cheese. *Netherlands. Milk Dairy J.*, 49: 191-199.
- Loureiro, V. and A. Querol, 1999. The prevalence and control of spoilage yeasts in foods and beverage. *Trends Food Sci. Technol.*, 10: 356-365.
- Lodder, J., 1970. *The Yeasts. A Taxonomic Study*. North Holland Publishing Co. Amsterdam.
- Deak, T., 1986. A Simplified Scheme for the Identification of Yeasts. *Methods for the Mycological Examination of Foods*. King, A.D., J.I. Pitt, L.R. Beuchat and J.E.L. Corry, (Eds.), Plenum. New York.
- Mihyar, G., M.I. Yamani and A. Al-sa'Ed, 1997. Resistance of yeast flora of labaneh to potassium sorbate and sodium benzoate. *J. Dairy Sci.*, 80: 2304-2309.
- Minervini, F., M.T. Montagna, G. Spilotros, L. Monaci, M.P. Santacroce and A. Viskonti, 2001. Survey on mycoflora of cow and buffalo dairy products from southern Italy. *Intl. J. Food Microbiol.*, 69: 141-146.
- Pereira-Dias, S., M.E. Potes, A. Marinho, M. Malfeito-Ferreira and V. Loureino, 2000. Characterization of yeasts flora isolated from an artisanal Portuguese ewes cheese. *Intl. J. Food Microbiol.*, 60: 55-63.
- Psomas, E., C. Andrighetto, E. Litropoulou-Tzanetaki, A. Lambard and N. Tzanetakis, 2001. Some probiotic properties of yeasts isolates from infant faeces and Feta cheese. *Intl. J. Food Microbiol.*, 69: 125-131.
- Roostita, R. and G.H. Fleet, 1996. Growth of yeasts in milk and associated changes to milk composition. *Intl. J. Food Microbiol.*, 31: 205-219.

- Suriyarachi, V.R. and G.H. Fleet, 1981. Occurrence and growth of yeasts in yoghurt. *Applied Environ. Microbiol.*, 42: 574-579.
- Spillmann, V. and O. Geiges, 1983. Identifikation von Hefen und schimmelpilzen aus bombierten joghurt-Packungen. *Milchwissenschaft*, 38: 129-133.
- Sarais, I., D. Piussi, V. Aquili and M.L. Stechini, 1996. The behavior of yeasts populations in stracchino cheese packaged under various conditions. *J. Food Prot.*, 59: 541-544.
- Vander Walt, J.P. and D. Yarrow, 1984. *Methods for the Isolation Maintenance, Classification and Identification of Yeasts*. Yeast Technol., Academic Press, pp: 45-104.
- Vanden Tempel, T. and M. Jakobsen, 1998. Yeasts associated with Danablu. *Intl. Dairy J.*, 8: 25-31.
- Viljoen, B.C. and T. Greyling, 1995. Yeasts associated with Cheddar and Gouda making. *Intl. J. Food Microbiol.*, 28: 79-88.
- Welthagen, J.J. and B.C. Viljoen, 1998. Yeasts profile in Gouda cheese during processing and ripening. *Intl. J. Food Microbiol.*, 41: 185-194.
- Wojtatowicz, M., J. Chrzanowska, P. Juszzyk, A. Skiba and A. Gdula, 2001. Identification and biochemical characteristic of yeasts microflora of Rokpol cheese. *Intl. J. Food Microbiol.*, 69: 135-140.
- Yamani, M.I. and M. Abu-Jaber, 1994. Yeast flora of Labaneh produced by in-bag straining of cow milk set yoghurt. *J. Dairy Sci.*, 77: 3558-3564.