



International Journal of Meat Science

ISSN 2071-7113

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Changes of Fatty Acid Profile During Gamma Irradiation of Rainbow Trout (*Oncorhynchus mykiss*) Fillets

S. Javan and A.A. Motallebi

Department of Veterinary, Science and Research Branch, Islamic Azad University, Tehran, Iran

Corresponding Author: S. Javan, Department of Veterinary, Science and Research Branch, Islamic Azad University, Tehran, Iran Tel: +98(13)44506781

ABSTRACT

The present study was conducted to evaluate the effect of different doses of gamma rays (0, 0.75, 1.5, 2.25, 3, 3.75 and 4.5 kGy) on fatty acids composition of Rainbow trout fillet. Irradiation of fillets was performed by gamma rays from a Co_{60} source. The results showed that total saturated fatty acid concentrations increased significantly ($p < 0.001$) with increasing irradiation dose, so that the control sample and fish muscles that irradiated with 4.5 kGy had the lowest and highest amounts of total saturated fatty acids (20.454 ± 0.011 and $19.228 \pm 0.040\%$, respectively). The amount of total Polyunsaturated Fatty Acids (PUFAs) in irradiated samples were significantly lower than control sample ($p < 0.001$) and amounts of total monounsaturated (MUFAs) were significantly higher than control samples ($p < 0.001$). The results indicated that the highest content of MUFAs and PUFAs were in samples irradiated with 3.75 kGy ($37.783 \pm 0.092\%$) and control samples ($37.677 \pm 0.104\%$), respectively. The lowest level of MUFAs and PUFAs were in control samples ($36.596 \pm 0.024\%$) and 4.5 kGy ($36.459 \pm 0.047\%$), respectively. The results of this study showed that Gamma irradiation influenced the fatty acid composition especially polyunsaturated fatty acids in rainbow trout fillet significantly ($p < 0.001$).

Key words: Effect, irradiation, Rainbow trout, fillet, fatty acid, profile

INTRODUCTION

Food irradiation is a means of food preservation that has been in development since the early part of the 20th century. Irradiation as a method of meat, poultry and fish preservation has excellent potential to improve meat safety and extend the shelf life (Fu *et al.*, 2000; Chawla *et al.*, 2003; Jeevanandam *et al.*, 2001; Mahrour *et al.*, 2003; Chouliara *et al.*, 2004, 2005; Morehouse, 2002). It is capable of improving the safety of many foods and extending their shelf life. There has been worldwide interest in using irradiation for preservation of various foods, including fishery products (Kreuzer, 1969; Kilgen, 2001). Irradiation of food up to an overall dose of 10 kGy is accepted in several countries for commercial food processing (Lacroix and Quattara, 2000).

The quality of meat is influenced by its lipid content and fatty acid composition. Fish meat is more or less susceptible to oxidative deterioration, depending on the degree of lipid saturation. Polyunsaturated fatty acids (PUFAs) benefit human health but also increase susceptibility to lipid oxidation. Oxidation leads to negative effects on quality parameters, such as flavour, colour, texture and nutritive value (Buckley *et al.*, 1995). During recent decades, research on n-3 highly unsaturated fatty acid (n-3 HUFA) in marine foods has intensified due to their beneficial effects

on human health. They have been shown to have curative and preventive effects on cardiovascular diseases, mortality and neurodevelopment in infants (Kinsella, 1986; Kinsella *et al.*, 1990; Hu *et al.*, 2001; Okada and Morrissey, 2007).

Gamma rays can convert water molecules to ions and free radicals (OH, H, H₃O, e⁻). Double bonds of carbon-carbon on unsaturated fatty acids and carbonyl groups of fatty acid and amino acids, are especially, prone to attack of free radicals. These unsaturated fatty acids are the primary source for the oxidation of lipids. With increasing irradiation dose, the amount of unsaturated fatty acids decreased at the end time of exposure (Brewer, 2009).

A review of the scientific and technical literature revealed some information about the effect of irradiation and other processes on fatty acid composition in fish. Ozden (2005) investigated about changes in fatty acid composition of marinated fish during its shelf-life. He found that total saturated fatty acid concentrations increased significantly in marinated fish during chilled storage. Haliloglu *et al.* (2002) compare fatty acids profiles of muscle lipids of three trout species (*Salvelinus alpinus*, *Salmo trutta fario* and *Oncorhynchus mykiss*) fed the same commercial diet and reared under the same conditions. They found that Palmitic acid (16:0) in total Saturated Fatty Acids (SFAs) and oleic acids (18:1 n-9) in monounsaturated fatty acids (MUFAs) were the most abundant FAs and significant differences were observed between fish species. Gamma irradiation at 50 kGy of vacuum-packed herring fillets at 0 °C did not affect the proportion of polyunsaturated fatty acids (Adam *et al.*, 1982). Recently, Kalyoncu *et al.* (2012) were studied on the seasonal fluctuations of fatty acid compositions of rainbow trout and ω3/ω6 fatty acids ratio of this fish species. Yildiz *et al.* (2006) worked on the effects of irradiation (0, 1, 3, 5 and 7 kGy) on trans fatty acids of ground beef and showed the greater doses from 3 enhanced the fatty acid oxidation. Mbarki *et al.* (2009) reported that low-dose irradiation had no adverse effect on the nutritionally important polyunsaturated fatty acids (PUFAs) of Mediterranean horse mackerel.

The objective of this study was evaluation of different doses of gamma rays effects on fatty acid composition of *O. mykiss* fillet samples.

MATERIALS AND METHODS

Fish sample: Cultured fresh Rainbow trout (*Oncorhynchus mykiss*) brought from Ranghinkaman farm in Guilan province in Iran during September, 2010. The average weight and length of the fish were 390±53 g and 33±2.2 cm, respectively. Then, fish were eviscerated, washed and packaged in polyethylene pouches and were packed in polystyrene boxes with ice (ice to fish ratio was 2:1) samples were transported to irradiation process within 18 h.

Irradiation: Samples were irradiated at the Iranian Atomic Energy Organization, Tehran, Iran, using a Gamma cell facility with 60 Cobalt radiation source. The applied doses in this study were 0, 0.75, 1.5, 2.25, 3, 3.75 and 4.5 kGy. Exposure time showed in Table 1 (Dose rate: 4.6 Gy/sec, Transit dose: 20.0 Gy, Activity: 19823.7 Ci). The absorbed dose was monitored by an alanine transfer dosimeter. Dosimetry of irradiation process was showed that samples were irradiated with deviation of ±20%.

Fish samples were maintained at 2±1°C during irradiation by using sealed ice covering the samples. Internal temperature of the facility was 20±1°C. For simulation of condition, non-irradiated fish muscle samples (control) were kept in polystyrene boxes in room temperature (20-22°C) until the end of irradiation process. Irradiation process and all analyses were performed using three samples (pouches) per treatment.

Table 1: Exposure time of irradiation of rainbow trout muscle

Irradiation dose (kGy)	Exposure time	
	Minute	Second
0.75	2	20
1.50	5	41
2.25	8	35
3.00	11	22
3.75	13	42
4.50	17	6

Lipid extraction and fatty acid compositions: Fatty acid composition was determined after methylation, by gas chromatography (Agilent 6890, Italy) using a BPX-Column (70120 m, 250 μm i.d, 0.2 μm film thickness). The chromatographic conditions were, as follows; injection volume: 0.5 mL, injection temperature: 250°C, detector and detector temperature: FID-280°C, column temperature, 120°C for 2 min, programmed at 5°C/min up to 220°C for 8 min. Fatty acids were identified by comparison of their retention time with those of authentic standard (Sigma Cod No. 189-19) and their contents were calculated on a weight percentage basis (ISO 5508, ISIRI 4091) (Christie, 1989). All chemical (Methanol, n-Hexane, Potassium hydroxide pellets) brands were MEREK.

Statistical analysis: The complete experiment was performed in triplicate. Significant differences between the samples were calculated by Spss 13.00 by One way analysis of variance (ANOVA). Comparison of means were based on Post Hoc multiple test (Duncan's test). Level of significance was set at $p < 0.001$.

RESULTS

Fatty acid composition in non-irradiated and irradiated rainbow trout muscle samples with 0.75, 1.5, 2.25, 3, 3.75 and 4.5 kGy are shown in Table 2. Content of total Saturated Fatty Acids (SFAs) in the non irradiated muscle of rainbow trout was 19.228%. The highest and lowest content of SFAs between treatments were in samples irradiated with 4.5 kGy (20.454%) and 0.75 kGy (19.043%), respectively. The predominant saturated fatty acid in all treatments was Palmitic acid ($C_{16:2}$). The content of this fatty acid had a significant differences between treatments ($p < 0.001$) and the lowest and the highest amount of this fatty acid were in control sample ($12.245 \pm 0.013\%$) and in samples irradiated with 4.5 kGy ($13.338 \pm 0.431\%$), respectively.

There were a significant differences ($p < 0.001$) in the content of total monounsaturated fatty acids (MUFAs) between treatments. The lowest and highest content of MUFAs in muscle of rainbow trout were in control samples (36.596%) and those irradiated with 3.75 kGy (37.783%), respectively. The most dominant MUFAs in muscle of rainbow trout were *cis*-oleic acid ($C_{18:1\omega9}$) and

The *cis*-11-Eicosaenoic acid ($C_{20:1\omega9}$). The difference these fatty acids between treatments were statistically significant ($p < 0.001$).

The quantity of polyunsaturated fatty acids (PUFAs) in control samples and samples irradiated with 0.75, 1.5, 2.25, 3, 3.75 and 4.5 kGy were 38.425, 38.468, 37.484, 37.277, 37.152, 37.152, 36.677, 36.459%, respectively.

The dominant polyunsaturated fatty acids in control sample were *cis*-Linoleic acid $C_{18:2\omega6}$ ($25.447 \pm 0.580\%$), α -Linolenic acid $C_{18:3\omega3}$ ($3.352 \pm 0.008\%$) and *cis*-5, 8, 11, 14, 14-Eicosapentaenoic acid $C_{20:5\omega3}$ ($3.507 \pm 0.003\%$).

Table 2: Variations in total fatty acid composition of irradiated filets of *Oncorhynchus mykiss* (% of total FA)

Fatty acid	Dose (kGy)						
	0	0.75	1.5	2.25	3	3.75	4.5
C12:0, lauric acid ***	0.020±0.001 ^b	0.013±0.003 ^{ab}	0.011±0.002 ^a	0.019±0.005 ^b	0.022±0.003 ^{bc}	0.030±0.002 ^c	0.029±0.000 ^c
C14:0, myristic acid	0.972±0.003 ^d	0.860±0.026 ^b	0.878±0.016 ^{bc}	0.872±0.008 ^{bc}	0.918±0.012 ^c	0.870±0.004 ^b	0.707±0.006 ^a
C15:0, pentadecanoic acid	0.181±0.002 ^c	0.176±0.005 ^{bc}	0.167±0.003 ^a	0.170±0.002 ^{abc}	0.169±0.001 ^{ab}	0.172±0.003 ^{bc}	0.181±0.005 ^{bc}
C16:0, palmitic acid	12.245±0.013 ^a	12.303±0.045 ^a	12.836±0.553 ^{ab}	12.270±0.047 ^a	12.839±0.105 ^{ab}	12.931±0.341 ^{ab}	13.388±0.431 ^b
C17:0, heptadecanoic acid	0.830±0.003 ^a	0.083±0.006 ^a	0.071±0.006 ^a	0.082±0.008 ^a	0.078±0.003 ^a	0.310±0.006 ^a	0.080±0.003 ^b
C18:0, stearic acid	4.471±0.014 ^c	4.256±0.040 ^a	4.636±0.121 ^e	4.545±0.035 ^d	4.325±0.022 ^b	4.251±0.039 ^a	4.489±0.014 ^{cd}
C20:0, arachidic acid	0.321±0.002 ^c	0.307±0.006 ^b	0.322±0.003 ^c	0.308±0.006 ^b	0.290±0.004 ^a	0.291±0.003 ^a	0.333±0.003 ^c
C22:0, behenic acid	0.823±0.003 ^b	0.943±0.003 ^c	0.863±0.036 ^d	1.155±0.115 ^f	0.780±0.023 ^a	0.840±0.025 ^c	1.158±0.094 ^f
C24:0, lignoceric acid	0.111±0.002 ^{ab}	0.102±0.002 ^{ab}	0.112±0.003 ^c	0.103±0.006 ^{ab}	0.098±0.004 ^b	0.080±0.004 ^a	0.108±0.036 ^{ab}
Total saturated	19.228	19.043	19.897	19.525	19.520	19.777	20.454
C14:1, myristoleic acid	0.059±0.002 ^{ab}	0.040±0.005 ^c	0.060±0.000 ^{ab}	0.069±0.004 ^a	0.059±0.004 ^{ab}	0.060±0.003 ^{ab}	0.058±0.003 ^b
C15:1,	0.039±0.007 ^a	0.041±0.003 ^a	0.042±0.003 ^a	0.039±0.001 ^a	0.040±0.001 ^a	0.041±0.002 ^a	0.039±0.002 ^a
C16:1, palmitoleic acid	2.093±0.005 ^c	2.030±0.036 ^c	2.408±0.011 ^{ab}	2.525±0.005 ^a	2.498±0.012 ^a	2.320±0.073 ^b	2.343±0.061 ^b
C17:1, heptaenoic acid	0.249±0.006 ^{ab}	0.241±0.004 ^b	0.248±0.008 ^{ab}	0.262±0.003 ^a	0.262±0.003 ^a	0.250±0.002 ^{ab}	0.250±0.004 ^{ab}
t.C18:1, <i>trans</i> -oleic acid-ω9	0.143±0.003 ^a	0.171±0.007 ^b	0.137±0.006 ^a	0.13±0.001 ^a	0.140±0.002 ^a	0.140±0.002 ^a	0.130±0.001 ^a
c.C18:1, <i>cis</i> -oleic acid-ω9	32.121±0.213 ^a	32.148±0.451 ^a	32.340±0.017 ^{ab}	32.177±0.449 ^a	32.189±0.142 ^a	32.813±0.105 ^b	32.150±0.442 ^a
C20:1, <i>cis</i> -11-eicosaenoic acid-ω9	1.381±0.009 ^c	1.470±0.010 ^d	1.576±0.005 ^b	1.625±0.014 ^a	1.529±0.006 ^c	1.560±0.004 ^b	1.615±0.015 ^a
C22:1, erucic acid-ω 9	0.402±0.003 ^d	0.474±0.008 ^c	0.570±0.013 ^a	0.565±0.005 ^a	0.508±0.008 ^b	0.489±0.009 ^b	0.538±0.013 ^a
C24:1, nervonic acid-ω 9	0.110±0.005 ^{ab}	0.158±0.003 ^b	0.109±0.002 ^{ab}	0.133±0.003 ^a	0.101±0.003 ^{ab}	0.110±0.002 ^{ab}	0.139±0.004 ^{ab}
Total monounsaturated	36.596	37.773	37.490	37.526	37.326	37.783	37.262
t.C18:2, <i>trans</i> -linoleic acid-ω6	0.218±0.003 ^c	0.169±0.010 ^a	0.189±0.002 ^b	0.180±0.004 ^{bc}	0.180±0.005 ^{bc}	0.176±0.005 ^{bc}	0.250±0.005 ^d
c.C18:2, <i>cis</i> -linoleic acid-ω6	24.166±0.306 ^{ab}	24.447±0.586 ^a	24.547±0.172 ^a	24.296±0.004 ^{ac}	24.103±0.025 ^{ab}	24.105±0.031 ^{ab}	23.830±0.295 ^b
gamma.C18:3, γ-linolenic acid-ω6	0.573±0.015 ^{ab}	0.564±0.018 ^{ab}	0.595±0.007 ^a	0.579±0.002 ^{ab}	0.605±0.005 ^a	0.547±0.006 ^b	0.535±0.005 ^b
alpha.C18:3, α-linolenic acid ω3	3.033±0.015 ^d	3.352±0.008 ^b	3.334±0.013 ^b	3.380±0.005 ^b	3.507±0.040 ^a	3.226±0.007 ^c	3.318±0.020 ^b
C20:3, <i>cis</i> -8, 11, 14-eicosatrienoic acid ω6	0.010±0.010	0.010±0.010	0.010±0.010	0.020±0.010	0.010±0.010	0.010±0.010	0.010±0.010
C20:3, <i>cis</i> -11, 14, 17-eicosatrienoic acid ω3	1.170±0.050	1.180±0.08	1.160±0.07	1.150±0.010	1.200±0.010	1.200±0.05	1.100±0.010
C20:4, arachidonic acid ω6	1.935±0.015	2.010±0.31	1.970±0.10	2.080±0.010	2.010±0.030	1.890±0.04	1.971±0.021
C20:5 <i>cis</i> -5, 8, 11, 14, 17-eicosapentaenoic ω3 (EPA)	3.392±0.008 ^b	3.507±0.033 ^a	3.435±0.053 ^{ab}	3.440±0.002 ^{ab}	3.427±0.006 ^{ab}	3.450±0.017 ^{ab}	3.373±0.012 ^b
C22:4 <i>cis</i> -7, 10, 13, 16-docosatetraenoic acid ω6	0.395±0.005 ^b	0.453±0.015 ^a	0.448±0.008 ^a	0.450±0.004 ^a	0.388±0.008 ^b	0.470±0.006 ^a	0.338±0.008 ^c
C22:6 <i>cis</i> -4, 7, 10, 13, 16, 19-docosahexaenoic ω3 (DHA)	1.785±0.005	1.733±0.067	1.798±0.053	1.702±0.023	1.722±0.055	1.800±0.050	1.734±0.005
Total polyunsaturated	37.677	37.425	37.468	37.277	37.152	36.874	36.459
Not detected	6.499	5.145	5.336	5.672	6.002	5.566	5.825

*Average of three lots analyzed, **Values reported are Mean±SD, ***Abcd values for each sample with different superscript letters in the same rows are significantly different at p<0.001

At the end of irradiation process, the content of cis-5,8,11,14,14-Eicosapentaenoic acid C_{20:5 ω 3} irradiated samples with 0.75, 1.5, 2.25, 3, 3.75 and 4.5 kGy, were 3.435 \pm 0.053, 3.450 \pm 0.017, 3.440 \pm 0.002, 3.427 \pm 0.006, 3.392 \pm 0.053 and 3.373 \pm 0.012%, respectively. These values for cis-4, 7, 10, 13, 16, 19-Docosahexaenoic acid C_{22:6 ω 3} were 1.733 \pm 0.067, 1.798 \pm 0.053, 1.800 \pm 0.050, 1.702 \pm 0.023, 1.722 \pm 0.055, 1.785 \pm 0.005 and 1.734 \pm 0.005%, respectively.

The differences between the values of the former fatty acid were significant (p<0.001) but this difference for the latter fatty acid was not significant (p<0.001).

DISCUSSION

The effects of irradiation on oxidation of lipids are well known. The results of our study were showed that the different doses of irradiation especially high doses (3-4.5 kGy) affect fatty acid composition in rainbow trout muscle. It is stated that there was relationship between irradiation dose and lipid oxidation whereas with increasing of irradiation dose, lipid oxidation will urge.

In our study, amount of palmitic acid (SFA) significantly increased during increasing irradiation dose and Oleic acid content was not significantly changed. However, Katta *et al.* (1991) found significant decrease in amount of Palmitic acid and increase in Oleic acid, as irradiation dose level increased (0.5-3 kGy) in chicken meat. Similar results for other fish species have also been reported (Rahnan *et al.*, 1995). Rady *et al.* (1988) showed no significant difference in total saturated and unsaturated fatty acids between irradiated (1, 3, 6 kGy) and non-irradiated frozen chicken muscle. Armstrong *et al.* (1994) reported no changes in fatty acid compositions of two species of Australian marine fish irradiated at doses up to 6.0 kGy. This finding was contradicted with our results.

Yilmaz and Gecgel (2007) showed that concentration of total trans fatty acids in irradiated ground beef samples had higher than the control samples and irradiated ground beef samples with 7 kGy had the highest total trans fatty acids. Hau and Liew (1993) reported that irradiation at 10 kGy caused a 16% decrease in the linoleic contents of grass prawns but Linolenic acid content didn't changed. Ozden and Erkan (2010) reported that total saturated fatty acid contents were increased in irradiated fish samples. These results were similar to our finding about SFAs. These study results generally agreed that irradiation of seafoods and meats had marginal effects on the lipids, including essential fatty acids. In this study.

Maxwell and Rady (1989) also reported a steady increase in oleic acid in the polar fractions with increasing doses of gamma irradiation. However, Hafez *et al.* (1985) did not find changes in the fatty acids (C16:0, C18:1 and C18:2) of soybeans at different radiation doses (1, 5, 10, 20, 40, 60, 80 and 100 kGy). Katta *et al.* (1991) found significant decrease in amount of Palmitic acid and increase in oleic acid as irradiation dose level increased (0.5-3 kGy) in chicken meat. These authors determined levels of other fatty acids notably polyunsaturated fatty acid (linoleic and arachidonic acid) did not change.

Al-Kahtani *et al.* (1996) reported Influence of irradiation on chemical components of tilapia and Spanish mackerel whereas Irradiation of tilapia at 1.5-10 kGy caused a decrease in some fatty acids (C14:0, C16:0 and 16:1) and In Spanish mackerel, 16:0 and 16:1 fatty acids decreased, when irradiated at 1.5-10 kGy.

CONCLUSION

In this study, results revealed a raise in fatty acid oxidation linked to raise dose of gamma irradiation.. Also irradiation was changed fatty acids composition especially MUFAs and PUFAs fatty acids in fish fillet. The 3.75 and 4.5 kGy irradiation induced the formation of polysaturated fatty acids.

ACKNOWLEDGMENT

This study was supported by the Iranian Fisheries Research Organization (Project No: 2-12-12-88064). We wish to thank Mrs. Gorji Fard Company Director, for performed the irradiation. Gas chromatographic analyses were performed by the oily seeds development and cultivation company, Tehran, Iran.

REFERENCES

- Adam, S., G. Paul and D. Ehlermann, 1982. Influence of ionizing radiation on the fatty acid composition of herring fillets. *Radiat. Phys. Chem.*, 20: 289-295.
- Al Kahtani, H.A., H.M. Abu Tarboush, A.S. Bajaber, M. Atia, A.A. Abou Arab and M.A. El Mojaddidi, 1996. Chemical changes after irradiation and post-irradiation storage in tilapia and spanish mackerel. *J. Food Sci.*, 61: 729-733.
- Armstrong, S.G., S.G. Wyllie and D.N. Leach, 1994. Effects of preservation by gamma-irradiation on the nutritional quality of Australian fish. *Food Chem.*, 50: 351-357.
- Brewer, M.S., 2009. Irradiation effects on meat flavor: A review. *Meat Sci.*, 81: 1-14.
- Buckley, D.J., P.A. Morrissey and J.I. Gray, 1995. Influence of dietary vitamin E on the oxidative stability and quality of pig meat. *J. Anim. Sci.*, 73: 3122-3130.
- Chawla, S.P., D.H. Kim, C. Jo, J.W. Lee, H.P. Song and M.W. Byun, 2003. Effect of gamma irradiation on the survival of pathogens in Kwamegi, a traditional Korean semidried seafood. *J. Food Protect.*, 66: 2093-2096.
- Chouliara, I., I.N. Sawaidis, N. Panagiotakis and M.G. Kontominas, 2004. Preservation of salted, vacuum-packaged, refrigerated sea bream (*Sparus aurata*) fillets by irradiation: Microbiological, chemical and sensory attributes. *Food Microbiol.*, 21: 351-359.
- Chouliara, I., I.N. Savvaidis, K. Riganakos and M.G. Kontominas, 2005. Shelf life extension of vacuum-packaged sea bream (*Sparus aurata*) fillets by combined γ -irradiation and refrigeration: Microbiological, chemical and sensory changes. *J. Sci. Food Agric.*, 85: 779-784.
- Christie, W.W., 1989. *Gas Chromatography and Lipids: A Practical Guide*. The Oily Press, Ayr, Scotland.
- Fu, J., W. Shen, J. Bao and Q. Chen, 2000. The decontamination effects of gamma irradiation on the edible gelatin. *Radiat. Phys. Chem.*, 57: 345-348.
- Hafez, Y.S., A.I. Mohamed, G. Singh and F.M. Hewedy, 1985. Effects of gamma irradiation on proteins and fatty acids of soybean. *J. Food Sci.*, 50: 1271-1274.
- Haliloglu, H.I., N.M. Aras and H. Yetim, 2002. Comparison of muscle fatty acids of three trout species (*Salvelinus alpinus*, *Salmo trutta fario*, *Oncorhynchus mykiss*) raised under the same conditions. *Turk. J. Vet. Anim. Sci.*, 26: 1097-1102.
- Hau, L.B. and M.S. Liew, 1993. Effects of γ -irradiation and cooking on vitamins B₆ and B₁₂ in grass prawns (*Penaeus monodon*). *Radiat. Phys. Chem.*, 42: 297-300.
- Hu, F.B., J.E. Manson and W.C. Willett, 2001. Types of dietary fat and risk of coronary heart disease: A critical review. *J. Am. Coll. Nutr.*, 20: 5-19.
- Jeevanandam, K., A. Kakatkar, S.N. Doke, D.R. Bongirwar and V. Venugopal, 2001. Influence of salting and gamma irradiation on the shelf-life extension of threadfin bream in ice. *Food Res. Int.*, 34: 739-746.
- Kalyoncu, L., Y. Yaman and A. Aktumsek, 2012. Determination of the seasonal changes on total fatty acid composition of rainbow trout, *Oncorhynchus mykiss* in Ivriz Dam Lake, Turkey. *Afr. J. Biotechnol.*, 9: 4783-4787.

- Katta, S.R., D.R. Rao, G.R. Sunki and C.B. Chawan, 1991. Effect of gamma irradiation of whole chicken carcasses on bacterial loads and fatty acids. *J. Food Sci.*, 56: 371-372.
- Kilgen, M.B., 2001. Irradiation Processing of Fish and Shellfish Products. In: *Food Irradiation: Principles and Applications*, Molins, R.A. (Ed.). Wiley, New York, USA., pp: 193-212.
- Kinsella, J.E., 1986. Food components with potential therapeutic benefits: The n-3 polyunsaturated fatty acids of fish oils. *Food Technol.*, 40: 89-97.
- Kinsella, J.E., B. Lokesh and R.A. Stone, 1990. Dietary n-3 polyunsaturated fatty acids and amelioration of cardiovascular disease: Possible mechanisms. *Am. J. Clin. Nutr.*, 52: 1-28.
- Kreuzer, R., 1969. *Freezing and Irradiation of Fish*. Fishing News (Books) Ltd., London, UK., Pages: 519.
- Lacroix, M. and B. Ouattara, 2000. Combined industrial processes with irradiation to assure innocuity and preservation of food products: A review. *Food Res. Int.*, 33: 719-724.
- Mahrour, A., S. Caillet, J. Nketsa-Tabiri and M. Lacroix, 2003. Microbial and sensory quality of marinated and irradiated chicken. *J. Food Protect.*, 66: 2156-2159.
- Maxwell, R.J. and A.H. Rady, 1989. Effect of gamma irradiation at various temperatures on air and vacuum packed chicken tissues II. Fatty acid profiles of neutral and polar lipids separated from muscle and skin irradiated at 2-5°C. *Int. J. Radiat. Applic. Instrument. Part C. Radiat. Phys. Chem.*, 34: 791-796.
- Mbarki, R., S. Sadok and I. Barkallah, 2009. Quality changes of the Mediterranean horse mackerel (*Trachurus mediterraneus*) during chilled storage: The effect of low-dose gamma irradiation. *Radiat. Phys. Chem.*, 78: 288-292.
- Morehouse, K.M., 2002. Food irradiation-US regulatory considerations. *Radiat. Phys. Chem.*, 63: 281-284.
- Okada, T. and M.T. Morrissey, 2007. Production of n-3 polyunsaturated fatty acid concentrate from sardine oil by lipase-catalyzed hydrolysis. *Food Chem.*, 103: 1411-1419.
- Ozden, O., 2005. Changes in amino acid and fatty acid composition during shelf-life of marinated fish. *J. Sci. Food Agric*, 85: 2015-2020.
- Ozden, O. and N. Erkan, 2010. Impacts of gamma radiation on nutritional components of minimal processed cultured sea bass (*Dicentrarchus labrax*). *Iran. J. Fish. Sci.*, 9: 265-278.
- Rady, A.H., J. Maxwell, E. Wiebicki and J.G. Phillips, 1988. Effect of gamma irradiation at various temperatures and packaging conditions on chicken tissues. I. Fatty acid profiles of neutral and polar lipids separated from muscle irradiated at -20°C. *Radiat. Phys. Chem.*, 31: 195-202.
- Rahnan, S.A., T.S. Huah, O. Nassan and N.M. Daud, 1995. Fatty acid composition of some Malaysian freshwater fish. *Food Chem.*, 54: 45-49.
- Yılmaz, I. and U. Gecgel, 2007. Effects of gamma irradiation on *trans* fatty acid composition in ground beef. *Food Control*, 18: 635-638.
- Yildiz, M., E. Sener and H. Gun, 2006. Effect of refrigerated storage on fillet lipid quality of rainbow trout (*Oncorhynchus mykiss* W.) Fed a diet containing different levels of DL α -tocopherol acetate. *Turk J. Vet. Anim. Sci.*, 30: 143-150.