



# International Journal of Pharmacology

ISSN 1811-7775

**science**  
alert

**ansinet**  
Asian Network for Scientific Information

## **An Update on Pharmacology of *Satureja* Species; From Antioxidant, Antimicrobial, Antidiabetes and Anti-hyperlipidemic to Reproductive Stimulation**

<sup>1,2</sup>Saeideh Momtaz and <sup>2</sup>Mohammad Abdollahi

<sup>1</sup>Department of Plant Science, Faculty of Natural and Agricultural Science,  
University of Pretoria, Pretoria, South Africa

<sup>2</sup>Faculty of Pharmacy and Pharmaceutical Sciences Research Center,  
Tehran University of Medical Sciences, Tehran, Iran

**Abstract:** Since ancient, the genus *Satureja* L. is well recognized for its therapeutic values. Only recently, scientists have been aware of its new medicinal aspects. In our last review, we evaluated existing scientific data about this genus. Regarding ongoing reports and new multi-functional properties of this plant we were interested to update pharmacology of *Satureja*. Data were gathered using of scientific books, journals, articles and websites including Pubmed, Scopus and Google Scholar up to 25th April 2010. Of initial search, 1540 records were reviewed for inclusion or exclusion in study, of which 71 publications were included. Different species of *Satureja* are famous for their analgesic, antiseptic, antimicrobial, antiviral, antioxidant, antiproliferative, antiprotozoal, antifungal, antidiarrheal, anti-inflammatory, anti-nociceptive, anticholinesterase and vasodilatory activities. The valuable therapeutic aspects of this genus are mostly correlated to the existence of essential oils, flavonoids and triterpenoids. The combination of two or more subspecies acting by different mechanisms to produce a synergistic effect should be considered. Based on the conducted assessments so far, antibacterial, antifungal, antiviral, antioxidant, anti-diabetic, vasodilatory and analgesic activities are characterized as the most applicable properties of *Satureja* subspecies. Further studies are needed to confirm novel pharmacological aspects of this genus in both animal and human.

**Key words:** *Satureja*, pharmacology, toxicology, animal, human

### **INTRODUCTION**

Accessibility and affordability of the medicinal herbs have made them as fundamental part of many people's life all over the world. The selection of medicinal plants is a conscious process, which has led to an enormous number of medicinal plants being used by the numerous cultures of the world (Heinrich *et al.*, 2004). Surveys conducted in Australia and US indicates that almost 48.5 and 34% of individuals had used at least one form of unconventional therapy, including medicine, respectively (Kim *et al.*, 2007). According to the World Health Organization (WHO), about 65-80% of the world's population in developing countries, due to poverty and lack of access to modern medicine, depends essentially on plants for their primary health care (Calixto, 2005). Regarding the lack of safe modern drugs, evaluation of effective plants for diseases like diabetes has been recommended by WHO (Kim *et al.*, 2007). In the recent years, efficacy of herbal medicines in diseases like inflammatory bowel disease (Rahimi *et al.*, 2009; Rezaie *et al.*, 2007), obesity

(Hasani-Ranjbar *et al.*, 2009a), diabetes (Rahimi *et al.*, 2005), pancreatitis (Monfared *et al.*, 2009), inflammatory and oxidant-related diseases (Hasani-Ranjbar *et al.*, 2009b; Rahimi *et al.*, 2010) has been systematically reviewed. It is estimated that close to 25% of the active compounds in currently prescribed synthetic drugs were first identified in plant resources (Halerstein, 2005) and 20000 plants have been used for medicinal proposes which, 4000 have been used commonly and 10% of those are commercial.

The genus *Satureja* L. (savory) belonging to the family Lamiaceae (labiatae), subfamily Nepetoideae and tribe Mentheae contains about 200 species of aromatic herbs and shrubs that are grown in Middle East, Mediterranean region to Europe, West Asia, North Africa, the Canary Islands and South America. Over 30 species of this genus are distributed in eastern parts of Mediterranean area (Cantino *et al.*, 1992). The known characteristic of the subfamily Nepetoideae is that its representatives contain more than 0.5% of essential oil. Moreover, remarkable differences between and within the

chemical composition of the essential oils of *Satureja* subspecies has been found (Slavkovska *et al.*, 2001). Phytochemical studies have revealed volatile oils, tannins, phenolic compounds, sterols, acids, gum, mucilage and pyrocatechol as the main components of *Satureja* species. The major constituents of the essential oils from savory species are phenols, carvacrol, thymol and flavonoids (Sefidkon *et al.*, 2004). Regarding existence of high amount of thymol and carvacrol in *Satureja* species, good smelling and simple cultivation, they are used as a flavoring compound in food, pharmaceutical and cosmetic industries. Savory species have been traditionally used as muscle pain reliever, tonic and carminative in treating stomach and intestinal disorders such as cramps, nausea, indigestion and diarrhea (Zargari, 1990). In the last five years, other properties of *Satureja* species such as antibacterial, antifungal, antioxidant, anti-diabetes, anti-HIV, anti-hyperlipidemic, reproduction stimulatory, expectorant and vasodilatory activities have been demonstrated (Abdollahi *et al.*, 2003; Amanlou *et al.*, 2005; Basiri *et al.*, 2007; Deans and Svoboda, 1989; Hajhashemi *et al.*, 2000; Sahin *et al.*, 2003; Yamasaki *et al.*, 1998; Vagionas *et al.*, 2007; Vosough-Ghanbari *et al.*, 2008).

An extensive review of the literature regarding biological and composition of *Satureja* species has been already published. The aim of this review is to complete on previous data specially those provided since the publication of the above-mentioned reviews. The present work is an update on pharmacology of *Satureja* up to May 2010.

## MATERIALS AND METHODS

The data presented in this review were collected using all scientific data come from encyclopedia books, journals, articles and websites Pubmed, Scopus, Scirus and Google Scholar up to 25th April 2010. Of initial search, 1540 records were reviewed for inclusion or exclusion in study, of which 71 publications were included.

## RESULTS AND DISCUSSION

As *Satureja* sp. are consumed commonly, their possible beneficial health effects have faced a great deal of attention. Recently, the consumption of natural antioxidant that occur in all higher plants and in all parts of them, has risen up regarding the side effects of synthetic ones (Bakkali *et al.*, 2008). Antioxidant activity of *Satureja* subspecies have been confirmed in many studies (Exarchou *et al.*, 2002; Madsen *et al.*, 1998; Radonic and Milos, 2003). The oils of genus *Satureja* are being filled with isopropanoids such as carvacrol, thymol,

flavonoids,  $\beta$ -caryophyllene,  $\gamma$ -terpinene, p-cymene and linalool, which are expected to possess strong antioxidant effects (Ruberto and Baratta, 2000). The constituents containing hydroxyl group exhibited relatively strong antioxidant effect in *S. Montana* (Radonic and Milos, 2003). Newly, *S. khuzestanica* essential oils (SKEO) was found to protect rat reproductive system from toxicity of cyclophosphamide that is currently used an anticancer alkylating agent having adverse effect on male reproductive tract (Rezvanfar *et al.*, 2008). In a related study, SKEO protected rats from cyclophosphamide-induced Hemorrhagic Cystitis (HC) by reduction of free radical-induced toxic stress (Rezvanfar *et al.*, 2010). Interestingly, the essential oil of *S. cilicica* has been suggested for usage as a natural antioxidant and aroma agent in butter in a dose dependent manner (Ozkan *et al.*, 2007).

Generally, production of essential oils by plants is a source of self protection against environmental pathogens. Antibacterial activity of essential oils depends on the type, composition and concentration of the oil, the type and concentration of the target microorganism, the composition of the substrate and the processing/storage conditions (Baydar *et al.*, 2004). The concentration of essential oils in *S. thymbra* and *S. parnassica* is known to be fluctuated. Those oils which were obtained during the flowering period were found the most potent in exhibiting the lowest Minimum Inhibitory Concentration (MIC) values and retaining considerable antibactericidal properties (Chorianopoulos *et al.*, 2006).

As yet, carvacrol and thymol play the fundamental roles in antimicroorganism activity of this genus. In an interesting study in Turkey, carvacrol the major constituent of the essential oil of *S. spicigera* was tested against 25 phytopathogenic bacterial strains and exhibited potent antibacterial activity over a broad spectrum of the bacteria. Their results demonstrated that *S. spicigera* oil could be used to control agricultural plant pathogenic bacteria and as a seed disinfectant (Kotan *et al.*, 2010). Of fifty-five compounds isolated from the oil of *S. cuneifolia*, linalool and  $\alpha$ -pinene were found to be the dominant compounds. The oil exhibited a complete inhibition of the growth of mostly all tested microorganisms at the lowest supplied concentration.

Antifungal activity of the different species of savory have been reported frequently (Boyraz and Ozcan, 2006; Sokovic *et al.*, 2002). Potent antiviral activities of *S. boliviana* (against viruses-HSV-1 and VSV) and *S. montana* (against HIV-1) have been proven by Abad *et al.* (1999) and Yamasaki *et al.* (1998), respectively.

Up to date, limited investigations have noticed to the anti-diabetes effects of *Satureja* sp. Anti-diabetic activity of *S. khuzestanica* has been proven in several studies.

It is thought that this activity is mostly mediated through antioxidant potential of this genus (Basiri *et al.*, 2007; Saadat *et al.*, 2004). The antioxidant property of *S. khuzestanica* is also known as a reasonable explanation for its triglyceride-lowering effects (Abdollahi *et al.*, 2003). The investigation showed that carvacrol, thymol and flavonoids (the main constituents of *S. khuzestanica* essential oil) are responsible to marked reduction of serum cholesterol levels in diabetic patients. Confirming those findings, Mchedlishvili *et al.* (2005) explained flavonoids isolated from *S. hortensis* responsible for reduction of cholesterol level in rabbits. Study on the effect of SKEO on male rat fertility has revealed significant improvements in potency, fecundity, fertility index and litter size. Additionally a significant decrease in post implantation loss has been observed (Haeri *et al.*, 2006).

Lamiaceae is well recognized as a family with great antispasmodic and pain reliever properties. Due to the broad distribution of the species of the genus *Satureja*, various compounds are recognized responsible for its analgesic effect. For instance, three known flavonoids (naringenin, eriodictyol and luteolin) isolated from *S. obovata* exhibited significant relaxant effects and (De Rojas *et al.*, 1999) while eriodictyol (5,7,3',4' -tetrahydroxyflavanone) isolated from same species showed slight vasodilatory activity in rats and (De Rojas *et al.*, 1999). Two studies clarified SKEO

has beneficial anti-inflammatory effects comparable to those of prednisolone, indomethacin and morphine (positive controls) (Amanlou *et al.*, 2005; Ghazanfari *et al.*, 2006). In contrast, some species does not show any anti-inflammatory effects. The essential oil of *S. thymbra* L. did not show any anti-inflammatory effect while it caused central analgesic activity in mice and rats (Karabay-Yavasoglu *et al.*, 2006).

Despite the confictions about the safety and toxicity of various species of *Satureja*, many people are consuming the subspecies as herbal tea, spices and additives every day worldwide. As earlier mentioned it is necessary to differ between the cytotoxic effects of the subspecies on microorganisms and its toxic effects in eukaryotic cells. The essential oils of *S. hortensis* and *S. khuzestanica* exhibited protective effects against the toxicity of hydrogen peroxide and malathion in rats, respectively (Behravan *et al.*, 2006; Basiri *et al.*, 2007). Another study revealed that SKEO protects reproductive system from toxicity of cyclophosphamide. Co-administration of SKEO significantly improved cyclophosphamide-induced changes in plasma testosterone, sperm quality, spermatogenesis, fertility, toxic stress and DNA damage (Rezvanfar *et al.*, 2008). The biological activities and the major components isolated from the most common species of *Satureja*. have been depicted in Table 1.

Table 1: *Satureja* L. subspecies with their therapeutic indications

Botanical and species	Local name	Therapeutic indications	Effective component	Reference
<i>S. acinos</i> L.	-	-	Triterpenes, oleanolic, ursolic, crataegolic acid, naringenin, rutinoid and eriodictyol	Escudero <i>et al.</i> (1985)
<i>S. aintabensis</i>	-	Antibacterial Antifungal	p-cymene	Azaz <i>et al.</i> (2005)
<i>S. bachtiarica</i>	Marzeh	Antibacterial	Carvacrol and p-cymene	Sefidkon <i>et al.</i> (2004)
<i>S. boissieri</i> Hausskn. ex Boiss.	-	Flavoring	Carvacrol, $\gamma$ -terpinene and p-cymene	Kurkcuoglu <i>et al.</i> (2001)
<i>S. boliviana</i> Briq.	-	Antibacterial Antiviral	$\Gamma$ -terpinene, $\beta$ -caryophyllene and germacrene D	Laurent <i>et al.</i> (1997), Viturro <i>et al.</i> (2000)
<i>S. brownei</i> (SW.) Briq.	-	Antibacterial	Pulegone and menthone	Caceres <i>et al.</i> (1991), Rojas and Usabillaga (2000)
<i>S. coerulea</i> Janka	-	Antibacterial	Germacrene-D	Turnen <i>et al.</i> (1998)
<i>S. cuneifolia</i> Ten.	Wild savory or cuneate Turkish savory	Antibacterial Antidiabetes Carminative	Carvacrol, $\gamma$ -terpinene, linalool and $\alpha$ -pinene	Aydn <i>et al.</i> (1995), Baydar <i>et al.</i> (2004)
<i>S. douglasii</i> (Benth.) Briq.	Douglas savory	-	Monoterpenes	Duke and Beckstorm-Sternberg (2001)
<i>S. edmondii</i>	Marzeh	Antibacterial	Carvacrol and thymol	Sefidkon <i>et al.</i> (2004)
<i>S. forbesii</i> (Benth.) Briq.	-	Antioxidant	Geranial, neral	Ortet <i>et al.</i> (2009)
<i>S. gilliesii</i>	-	Antiprotozoal	Mono and sesquiterpene peroxides	Labbe <i>et al.</i> (1993)
<i>S. gtabella</i> (Michx.) BRIQ.	Smooth savory	-	Germacrene D, isomethanol, menthone and limonene	Duke and Beckstorm-Sternberg (2001)
<i>S. grandiflora</i> L.	French savory	-	Germacrene D, isomethanone, pulegone and methanol	Duke and Beckstorm-Sternberg (2001)
<i>S. hortensis</i> L.	Summer savory	Analgesic Antibacterial Antifungal Antioxidant Antihyperglycemic Flavoring	Carvacrol, $\gamma$ -terpinene, thymol and p-cymene	Hajhashemi <i>et al.</i> (2002), Zargari (1990) Gulluce <i>et al.</i> (2003), Sahin <i>et al.</i> (2003), Behravan <i>et al.</i> (2006), Mchedlishvili <i>et al.</i> (2005)

Table 1: Continued

Botanical and species	Local name	Therapeutic indications	Effective component	Reference
<i>S. icarica</i> P.H. Davis	-	Antibacterial	Carvacrol	Tumen <i>et al.</i> (2000)
<i>S. intermedia</i> C.A. Mey	-	-	Thymol, $\gamma$ -terpinene and p-cymene	Sefidkon and Jamzad (2004)
<i>S. khuzestanica</i> Jamzad.	-	Antibacterial Antidiabetes antioxidant Antihyperglycemic Antiinflammatory Antinociceptive	P-cymene and carvacrol	Haeri <i>et al.</i> (2006), Saadat <i>et al.</i> (2004)
<i>S. kitabelii</i> Wierzb. ex Heuff.	-	-	p-cymene, geraniol and $\beta$ -elemene	
<i>S. macrantha</i> C.A. Mey.	-	Antibacterial Antifungal	Carvacrol, p-cymene, limonene and thymol	Azaz <i>et al.</i> (2005), Sefidkon and Jamzad (2004)
<i>S. montana</i> L.	Winter savory	Antibacterial Antifungal Carminative Antioxidant Anticholinesterase Flavoring	Thymol, p-cymene, carvacrol, $\gamma$ -terpinene, flavonoids, triterpenes	Oussalah <i>et al.</i> (2007), Mastelic and Jerkovic (2002)
<i>S. mutica</i> Fisch. and C. A. Mey.	-	Antibacterial Antimikotic	Carvacrol, thymol, $\gamma$ -terpinene, p-cymene and methyl thymol	Behravan <i>et al.</i> (2004), Gohari <i>et al.</i> (2005), Sefidkon and Jamzad (2004), Sefidkon <i>et al.</i> (2004)
<i>S. obovata</i> Lag.	Iberian savory	Carminative	Flavonoids	Duke and Beckstorm-Stenberg (2001), Sanchez-de-Rojas <i>et al.</i> (1996)
<i>S. parnassica</i> Heldr. and Sart ex Boiss.	-	Antibacterial	Caryophyllene, carvacrol, caryophyllene oxide, spathulenol, p-cymene and linalool	Chorianopoulos <i>et al.</i> (2006), Tzakou and Skaltsa (2003)
<i>S. parvifolia</i> (Phil.) Epl.	Small-leaf Pampa savory	Analgesic Antibacterial Antifungal	Piperitone oxide	Duke and Beckstorm-Stenberg (2001), Viturro <i>et al.</i> (2000), Zygadlo and Grosso (1995)
<i>S. pilosa</i> Velen	-	Antibacterial	-	Tumen <i>et al.</i> (2000)
<i>S. punctata</i>	-	Antiprotozoal	Geraniol, neral, alpha-bisabolol and (E)-nerolidol	Tariku <i>et al.</i> (2010)
<i>S. sahendica</i> Bomm.	-	Culinary herbs	Thymol, p-cymene and $\gamma$ -terpinene	Sefidkon <i>et al.</i> (2004)
<i>S. spicigera</i> (C. Koch) Boiss.	-	Antibacterial	Thymol, p-cymene, $\gamma$ -terpinene and carvacrol	Sefidkon and Jamzad (2004), Azaz <i>et al.</i> (2005), Eftekhar <i>et al.</i> (2009)
<i>S. spinosa</i> L.	-	Antibacterial	Monoterpene hydrocarbons and phenolic monoterpenes	Chorianopoulos <i>et al.</i> (2004)
<i>S. subspicata</i> Vis.	-	Antibacterial Antifungal	Carvacrol, $\alpha$ -pinene, p-cymene, $\gamma$ -terpinene and thymol methyl ether	Skocibusic <i>et al.</i> (2008)
<i>S. thymbra</i> L.	Goat oregano	Antifungal Antiinflammatory Antinociceptive	Carvacrol and $\gamma$ -terpinene	Glamoclija <i>et al.</i> (2006), Sokovic, <i>et al.</i> (2002)
<i>S. wiedemanniana</i> (Lallen.) Velen	-	Antibacterial	Carvacrol, thymol, p-cymene and $\gamma$ -terpinene	Baser <i>et al.</i> (2001)

## CONCLUSIONS

Clearly, herbal-based pharmaceuticals, cosmetics and flavoring are more welcoming every day. Numerous studies have been conducted to brighten both theoretical and practical aspects of the traditional usage of *Satureja* sp. There are remarkable varieties in chemical composition and therapeutic values of different species of this plant. In a recent investigation in Ethiopia, the oil of *S. punctata* contained 67 compounds of which geraniol and neral were the main components with significant antiprotozoal activities (Tariku *et al.*, 2010). Similarly, these compounds were the major constituents of the essential oil of *S. forbesii* with potent antioxidant activity (Ortet *et al.*, 2009). Sefidkon *et al.* (2009) presented carvacrol and p-cymene as the most frequent component in *S. bachtiarica* oil while carvacrol and thymol were the

most fluctuated elements in the oil of *S. mutica*. Their results showed the high antibacterial effect of these oils. Based on the conducted assessments so far, antibacterial, antifungal, antiviral, antioxidant, anti-diabetic, vasodilatory and analgesic activities are characterized as the most applicable properties of *Satureja* subspecies. In 2009 the inhibitory activity of acetyl- and butyrylcholinesterase by *S. montana* extracts was assessed as a potential indicator for the control of Alzheimer's disease. The result indicated a significant anticholinesterase activity of this species (Silva *et al.*, 2009).

According to the literature reviews, more than 90% of the present investigations only emphasized on the antibacterial activities of savory sp. while the other biological effects also seem promising. Monitoring and standardization is needed to direct studies passing

through clinical trials. Various molecular, cellular and whole animal models should be used to demonstrate more valuable significance of *Satureja* sp. Future exploration are critical to assess the synergistic effects, efficacy and edibility of these essential oils.

#### ACKNOWLEDGMENT

No financial support was obtained for this study.

#### REFERENCES

- Abad, M.J., P. Bermejo, E. Gonzales, I. Iglesias, A. Irurzun and L. Carrasco, 1999. Antiviral activity of Bolivian plant extracts. *Gen. Pharmacol.*, 23: 499-503.
- Abdollahi, M., A. Salehnia, S.H.R. Mortazavi, M. Ebrahimi and A. Shafiee *et al.*, 2003. Antioxidant, antidiabetic, antihyperlipidemic, reproduction stimulatory properties and safety of *Satureja Khuzestanica* essential oil as in rat *in vivo*: A toxicopharmacological study. *Med. Sci. Monitor*, 9: 331-335.
- Amanlou, M., F. Dadkhah, A. Salehnia, H. Farsam and A.R. Dehpour, 2005. An anti-inflammatory and anti-nociceptive effects of hydrochloric extract of *Satureja khuzistanica* Jamzad extract. *J. Pharm. Pharmaceut. Sci.*, 8: 102-106.
- Aydn, S., Y. Oztürk, R. Beis and K.H.C. Baer, 1995. Investigation of *Origanum onites*, *Sideritis congesta* and *Satureja cuneifolia* essential oils for analgesic activity. *Phytother. Res.*, 10: 342-344.
- Azaz, A.D., M. Kurkcuoglu, F. Satil, K. Baser and G. Tumen, 2005. *In vitro* antimicrobial activity and chemical composition of some *Satureja* essential oils. *Flavour Fragr. J.*, 20: 587-591.
- Bakkali, F., S. Averbeck, D. Averbeck and M. Idaomar, 2008. Biological effects of essential oils-a review. *Food Chem. Toxicol.*, 46: 446-475.
- Baser, K.H., G. Tumen, N. Tabanca and F. Demirci, 2001. Composition and antibacterial activity of the essential oils from *Satureja wiedemanniana* (Lallem.) Velen. *Z Naturforsch. C*, 56: 731-738.
- Basiri, S., H. Esmaily, S. Vosough-Ghanbari, A. Mohammadirad, N. Yasa and M. Abdollahi, 2007. Improvement by *Satureja khuzestanica* essential oil of malathion-induced red blood cells Pesticide *Biochem. Physiol.*, 89: 124-129.
- Baydar, H., O. Sagdic, G. Ozkan and T. Karadogan, 2004. Antibacterial activity and composition of essential oils from *Origanum*, *Tymbra* and *Satureja* species with commercial importance in Turkey. *Food Control*, 15: 169-172.
- Behravan, J., M. Ramezani, J. Kasaian and Z. Sabeti, 2004. Antimycotic activity of the essential oil of *Satureja mutica* Fisch and C.A. Mey from Iran. *Flavour Fragr. J.*, 19: 421-423.
- Behravan, J., F. Mosaffaa, G. Karimi and M. Iranshahi, 2006. Antigenotoxic effects of *Satureja hortensis* L. on rat lymphocytes exposed to oxidative stress. *Planta Med.*, 72: 113-125.
- Boyras, N. and M. Ozcan, 2006. Inhibition of phytopathogenic fungi by essential oil, hydrosol, ground material and extract of summer savory (*Satureja hortensis* L.) growing wild in Turkey. *Int. J. Food Microbiol.*, 107: 238-242.
- Caceres, A., A.V. Alvarez, A.E. Ovando and B.E. Samayoa, 1991. Plants used in Guatemala for the treatment of respiratory diseases. 1. Screening of 68 plants against gram-positive bacteria. *J. Ethnopharmacol.*, 31: 193-208.
- Calixto, J.B., 2005. Twenty-five years of research on medicinal plants in Latin America: A personal view. *J. Ethnopharmacol.*, 100: 131-134.
- Cantino, P.D., R.M. Harley and S.J. Wagstaff, 1992. Genera of *Labiatae* Status and Classification. In: *Advances in Labiatae Science*, Harley, R.M. and T. Reynolds (Eds.). Royal Botanic Gardens Press, Kew., UK, pp: 511-522.
- Chorianopoulos, K., E. Kalpoutzakis, N. Aligiannis, S. Mitaku, G. Nychas and S.A. Haroutounian, 2004. Essential oils of *Satureja*, *Origanum* and *Thymus* species: Chemical composition and antibacterial activities against foodborne pathogens. *J. Agric. Food Chem.*, 52: 8261-8267.
- Chorianopoulos, N., E. Evergetis, A. Mallouchos, E. Kalpoutzakis, G.J. Nychas and S.A. Haroutounian, 2006. Characterization of the essential oil volatiles of *Satureja thymbra* and *Satureja parnassica*: Influence of harvesting time and antimicrobial activity. *J. Agric. Food Chem.*, 54: 3139-3145.
- De Rojas, V.R.S., B. Somoza, T. Ortega, A.M. Villar and T. Tejerina, 1999. Vasodilatory effect in rat aorta of eriodictyol obtained from *Satureja obovata*. *Planta Med.*, 65: 234-238.
- Deans, S.G. and K.P. Svoboda, 1989. Antibacterial activity of summer savory (*Satureja hortensis* L.) essential oil and its constituents. *J. Hort. Sci.*, 64: 205-210.
- Duke, J.A. and S.M. Beckstorm-Sternberg, 2001. *Handbook of Medicinal Mints (Aromathematics) Phytochemicals and Biological Activities*. CRC Press, USA., pp: 273-290.
- Eftekhari, F., F. Raei, M. Yousefzadi, S.N. Ebrahimi and J. Hadian, 2009. Antibacterial activity and essential oil composition of *Satureja spicigera* from Iran. *Z Naturforsch. C*, 64: 20-24.

- Escudero, J., C. López, R. M. Rabanal and S. Valverde, 1985. Secondary metabolites from *Satureja* species. new triterpenoid from *Satureja acinos*. J. Nat. Prod., 48: 128-131.
- Exarchou, V., N. Nenadis, M. Tsimidou, I.P. Gerothannassis, A. Troganis and D. Boskou, 2002. Antioxidant activities and phenolic composition of extracts from Greek oregano, Greek sage and summer savory. J. Agric. Food Chem., 50: 5294-5299.
- Ghazanfari, G., B. Minaie, N. Yasa, L. Ashtaral Nakhai and A. Mohammadirad *et al.*, 2006. Biochemical and histopathological evidences for beneficial effects of *Satureja khuzestanica jamzad* essential oil on the mouse model of inflammatory bowel diseases. Toxicol. Mech. Methods, 16: 365-372.
- Glamoclija, J., M. Sokovic, J. Vukojevic, I. Milenkovic and L.J.L.D. Van Griensven, 2006. Chemical composition and antifungal activities of essential oils of *Satureja thymbra* L. and *Salvia pomifera* sp. calycina (Sm.). Hayek J. Essential Oil Res., 18: 115-117.
- Gohari, A., A. Hajakhondi, A. Shafee and E.S. Ebrahimi, 2005. Chemical composition of the essential oils of *Satureja atropatana* and *Satureja mutica* growing wild in Iran. J. Essential Oil Res., 17: 17-18.
- Gulluce, M., M. Sokmen, D. Daferera, G. Agar, H. Ozkan and N. Kartal, 2003. *In vitro* antibacterial, antifungal and antioxidant activities of the essential oil and methanol extracts of herbal parts and callus cultures of *Satureja hortensis* L. J. Agric. Food Chem., 51: 3958-3965.
- Haeri, S., B. Minaie, G. Amin, S. Nikfar, R. Khorasani, H. Esmaily, A. Salehnia and M. Abdollahi, 2006. Effect of *Satureja khuzestanica* essential oil on male rat fertility. Fitoterapia, 77: 495-499.
- Hajhashemi, V., A. Ghannadi and S.K. Pezeshkian, 2002. Antinociceptive and anti-inflammatory effects of *Satureja hortensis* L. extracts and essential oil. J. Ethnopharmacol., 82: 83-87.
- Hajhashemi, V., H. Sadrae, A.R. Ghannadi and M. Mohseni, 2000. Antispasmodic and anti-diarrhoeal effect of *Satureja hortensis* L. essential oil. J. Ethnopharmacol., 71: 187-192.
- Halerstein, R.A., 2005. Medicinal plants: Historical and cross-cultural usage patterns. Ann. Epidemiol., 15: 686-699.
- Hasani-Ranjbar, S., B. Larijani and M. Abdollahi, 2009a. A systematic review of the potential herbal sources of future drugs effective in oxidant-related diseases. Inflamm. Allergy Drug Targets, 8: 2-10.
- Hasani-Ranjbar, S., N. Nayeibi, B. Larijani and M. Abdollahi, 2009b. A systematic review of the efficacy and safety of herbal medicines used in the treatment of obesity. World J. Gastroenterol., 15: 3073-3085.
- Heinrich, M., J. Barnes, S. Gibbons and E.M. Williamson, 2004. Fundamentals of Pharmacognosy and Phytotherapy. Elsevier Science Ltd., Oxford.
- Karabay-Yavasoglu, U.N., S. Baykan, B. Ozturk, S. Apaydin and I. Tuglular, 2006. Evaluation of the antinociceptive and anti-inflammatory activities of *Satureja thymbra* L. essential oil. Pharm. Biol., 44: 585-591.
- Kim, J.D., S.M. Kang, M.Y. Park, T.Y. Jung, H.Y. Choi and S.K. Ku, 2007. Ameliorative anti-diabetic activity of dangnyosoko, a Chinese herbal medicine, in diabetic rats. Biosci. Biotechnol. Biochem., 71: 1527-1534.
- Kotan, R., A. Cakir, F. Dadasoglu, T. Aydin and R. Cakmakci *et al.*, 2010. Antibacterial activities of essential oils and extracts of Turkish Achillea, *Satureja* and Thymus species against plant pathogenic bacteria. J. Sci. Food Agric., 90: 145-160.
- Kurkcuglu, M., G. Tumen and K.H.C. Baser, 2001. Essential oil constituents of *Satureja boissieri* from Turkey. Chem. Nat. Compd., 37: 329-331.
- Labbe, C., M. Castillo and J.D. Connolly, 1993. Mono and sesquiterpenoids from *Satureja gilliesii*. Phytochemistry, 34: 441-444.
- Laurent, D., A. Vilaseca, J. Chantraine, C. Ballivian, G. Saavedra and R. Ibanez, 1997. Insecticidal activity of essential oils on *Triatoma infestans*. Phytother. Res., 11: 285-290.
- Madsen, H.L., B. Sorensen, L.H. Skibsted and G. Bertelsen, 1998. The antioxidative activity of summer savory (*Satureja hortensis* L.) and rosemary (*Rosmarinus officinalis* L.) in dressing stored exposed to light or in darkness. Food Chem., 63: 173-180.
- Mastelic, J. and I. Jerkovic, 2002. Gas chromatography-mass spectrometry analysis of free and glycoconjugated aroma compounds of seasonally collected *Satureja montana* L. Food Chem., 80: 135-140.
- Mchedlishvili, D., Z. Kuchukashvili, T. Tabatadze and G. Davitaia, 2005. Influence of flavonoids isolated from *Satureja hortensis* L. on hypercholesterolemic rabbits. Res. Lett., 37: 259-260.
- Monfared, S.S.M.S., H. Vahidi, A.H. Abdolghaffari, S. Nikfar and M. Abdollahi, 2009. Antioxidant therapy in the management of acute, chronic and post-ERCP pancreatitis: A systematic review. World J. Gastroenterol., 15: 4481-4490.
- Ortet, R., E.L. Regalado, O.P. Thomas, J.A. Pino, M.D. Fernandez and J.J. Filippi, 2009. Composition and antioxidant properties of the essential oil of the endemic cape verdean *Satureja forbesii*. Nat. Prod. Commun., 4: 1277-1280.

- Oussalah, M., S. Caillet, L. Saucier and M. Lacroix, 2007. Inhibitory effects of selected plant essential oils on the growth of four pathogenic bacteria: *E. coli* O157: H7, *Salmonella typhimurium*, *Staphylococcus aureus* and *Listeria monocytogenes*. Food Control, 18: 414-420.
- Ozkan, G., B. Simsek and H. Kuleasan, 2007. Antioxidant activities of *Satureja cilicica* essential oil in butter and *in vitro*. J. Food Eng., 79: 1391-1396.
- Radonic, A. and M. Milos, 2003. Chemical composition and *in vitro* evaluation of antioxidant effect of free volatile compounds from *Satureja montana* L. Free Radical Res., 37: 673-679.
- Rahimi, R., S. Nikfar, B. Larijani and M. Abdollahi, 2005. A review on the role of antioxidants in the management of diabetes and its complications. Biomed. Pharmacother., 59: 365-373.
- Rahimi, R., S. Mozaffari, M. Abdollahi, 2009. On the use of herbal medicines in management of inflammatory bowel diseases: A systematic review of animal and human studies. Dig. Dis. Sci., 54: 471-480.
- Rahimi, R., S. Ghiasi, H. Azimi, S. Fakhari and M. Abdollahi, 2010. A review of the herbal phosphodiesterase inhibitors: future perspective of new drugs. Cytokine, 49: 123-129.
- Rezaie, A., R.D. Parker and M. Abdollahi, 2007. Oxidative stress and pathogenesis of inflammatory bowel disease: An epiphenomenon or the cause? Dig. Dis. Sci., 52: 2015-2021.
- Rezvanfar, M.A., R.A. Sadrkhanlou, A. Ahmadi, H. Shojaei-Sadee and M.A. Rezvanfar *et al.*, 2008. Protection of cyclophosphamide-induced toxicity in reproductive tract histology, sperm characteristics and DNA damage by an herbal source; evidence for role of free-radical toxic stress. Hum. Exp. Toxicol., 27: 901-910.
- Rezvanfar, M.A., A.A. Farshid, R.A. Sadrkhanlou, A. Ahmadi, M.A. Rezvanfar, A. Salehnia and M. Abdollahi, 2010. Benefit of *Satureja khuzestanica* in subchronically rat model of cyclophosphamide-induced hemorrhagic cystitis. Exp. Toxicol. Pathol.,
- Rojas, L.B. and A. Usubillaga, 2000. Composition of the essential oil of *Satureja brownei* (SW.) Briq. from Venezuela. Flavour Fragr. J., 15: 21-22.
- Ruberto, G. and M.T. Baratta, 2000. Antioxidant activity of selected essential oil components in two lipid model systems. Food Chem., 69: 167-174.
- Saadat, M., S. Pournourmohammadi, M. Donyavi, R. Khorasani, G. Amin, A. Salehnia and M. Abdollahi, 2004. Alteration of rat hepatic glycogen phosphorylase and phosphoenolpyruvate carboxykinase activities by *Satureja khuzestanica* Jamzad essential oil. J. Pharm. Pharmaceut. Sci., 7: 310-314.
- Sahin, F., I. Karaman, M. Gulluce, H. Ogutcu and M. Sengul *et al.*, 2003. Evaluation of antimicrobial activities of *Satureja hortensis* L. J. Ethnopharmacol., 87: 61-65.
- Sanchez-de-Rojas, V., B. Somoza, T. Ortega, A.M. Villar and T. Tejerina, 1996. Isolation of vasodilatory active flavonoids from the traditional remedy *Satureja obovata*. Planta Med., 62: 272-274.
- Sefidkon, F. and J. Jamzad, 2004. Essential oil composition of *Satureja spicigera* (koch) Boiss. from Iran. Flavour Fragrance J., 19: 571-573.
- Sefidkon, F. and Z. Jamzad and M. Mirza, 2004. Chemical variation in the essential oil of *Satureja sahendica* from Iran. Food Chem., 88: 325-328.
- Sefidkon, F., F. Askari, L. Sadeghzadeh and P. Oulia, 2009. Antimicrobial effects of the essential oils of *Satureja mutica*, *S. edmondi* and *S. bachtiarica* against *Salmonella paratifi* A and B. Iran. J. Biol., 22: 249-259.
- Silva, F.V., A. Martins, J. Salta. N.R. Neng and J.M. Nogueira *et al.*, 2009. Phytochemical profile and anticholinesterase and antimicrobial activities of supercritical versus conventional extracts of *Satureja montana*. J. Agric. Food Chem., 57: 11557-11563.
- Skocibusic, M., N. Bezic and V. Dunkic, 2008. Phytochemical composition and antimicrobial activities of the essential oils from *Satureja subspicata* Vis. growing in Croatia. Food Chem. Toxicol., 46: 446-475.
- Slavkovska, V., J. Jancic, S. Bojovic, S. Milosavljevic and D. Djokovic, 2001. Variability of essential oils of *Satureja montana* L. and *Satureja kitaibelii* Wierzb. ex Heuff. from the central part of the Balkan peninsula. Phytochemistry, 57: 71-76.
- Sokovic, M., O. Tzakou, D. Pitarokili and M. Couladis, 2002. Antifungal activities of selected aromatic plants growing wild in Greece. Nahrung, 46: 317-320.
- Tariku, Y., A. Hymete, A. Hailu and J. Rohloff, 2010. Essential-oil composition, antileishmanial and toxicity study of artemisia abyssinica and *Satureja punctata* ssp. punctata from Ethiopia. Chem. Biodivers, 7: 1009-1018.
- Tumen, G., K.H.C. Baser, B. Demirci and N. Ermin, 1998. The essential oils of *Satureja coerulea* Janka and *Thymus aznavourii* Velen. Flavour Fragr. J., 13: 65-67.
- Tumen, G., F. Satil, H. Duman and K.H.C. Baser, 2000. Two new records for turkey: *Satureja icarica* P.H. Davis, *Satureja pilosa* Velen. Turk. J. Bot., 24: 211-214.
- Tzakou, O. and H. Skaltsa, 2003. Composition and antibacterial activity of the essential oil of *Satureja parnassica* subsp. Parnassica. Planta Med., 69: 282-284.



- Vagionas, K., K. Graikou, O. Ngassapa, D. Runyoro and I. Chinou, 2007. Composition and antimicrobial activity of the essential oils of three *Satureja* species growing in Tanzania. *Food Chem.*, 103: 319-332.
- Vituro, C.I., A. Molina, I. Guy, B. Charles, H. Guinaudeau and A. Fournet, 2000. Essential oils of *Satureja boliviana* and *S. parvifolia* growing in the region of Jujuy, Argentina. *Flavour Fragrance J.*, 15: 377-382.
- Vosough-Ghanbari, S., R. Rahimi., S. Kharabaf., S. Zeinali and A. Mohammadirad *et al.*, 2008. Effects of *Satureja khuzestanica* on serum glucose, lipids and markers of oxidative stress in patients with type 2 diabetes mellitus: A double-blind randomized controlled trial. *Evidence Based Complimentary Alternative Med.*, 10.1093/ecam/nen018
- Yamasaki, K., M. Nakamo, T. Kawahata, H. Mori and T. Otake *et al.*, 1998. Anti-HIV-1 activity of herbs in Labiatae. *Biol. Pharm. Bull.*, 21: 829-833.
- Zargari, A., 1990. *Medicinal Plants*. 4th Edn., Tehran University Publications, Tehran, pp: 42-45.
- Zygadlo, J.A. and N.R. Grosso, 1995. Comparative study of the antifungal activity of essential oils from aromatic plants growing wild in the central region of Argentina. *Flavour Fragrance J.*, 10: 113-118.