

ISSN 1682-8356
ansinet.org/ijps



INTERNATIONAL JOURNAL OF
POULTRY SCIENCE

ANSI*net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan
Mob: +92 300 3008585, Fax: +92 41 8815544
E-mail: editorijps@gmail.com

Productive Performance and Immune Response of Laying Hens as Affected by Dietary Propolis Supplementation

A. Galal¹, A.M. Abd El - Motaal², A.M.H. Ahmed¹ and T.G. Zaki³

¹Department of Poultry Production, Faculty of Agriculture, Ain Shams University, Cairo, Egypt

²Department of Poultry Nutrition Research, Animal Production Research Institute, Agricultural Research Center, Giza, Egypt

³Department of Poultry Breeding Research, Animal Production Research Institute, Agricultural Research Center, Giza, Egypt

Abstract: An experiment was conducted to evaluate the efficacy of supplemental Propolis on productive performance and immune response of laying hens. One hundred and twenty Hy-Line White strain were divided into four groups of 30 each. They were fed 0, 50, 100 and 150 mg of supplemental Propolis for 8 weeks (46-54wk). Chickens in all groups were reared under the same environmental, managerial and hygienic conditions. Feed and water were supplied *ad libitum*. The average high and low ambient temperatures recorded during the experimental period were 22.6 and 15.8°C, respectively. The performance data revealed that the laying hens fed diet containing 100 and 150 mg Propolis were significantly consumed more feed than control-group. Similar trend, but not statistically significant, was observed for hens fed diet added 50 mg Propolis. With respect to egg mass, it could be noticed that the laying hens fed diet containing 100 and 150 mg Propolis significantly produced heaviest egg mass compared to control-group. The hens fed diet added 50 mg Propolis were intermediate. The increase feed intake and egg mass in propolis groups, resulting in significantly improve feed conversion ratio compared to control-group. Eggshell quality was significantly affected by Propolis supplementation, whereas the percentage and thickness of eggshell were significantly increased in the eggs produced from hens fed diet containing medium or high level of Propolis. In accordance to hematological parameters, the medium or high level of dietary Propolis significantly increased hematocrit level, plasma total protein and globulin. Conversely, the plasma cholesterol and liver enzymes were significantly reduced when laying hens fed diet containing 100 or 150 mg Propolis. *In vivo* cell-mediated immune response as measured by PHA-P stimulation (wattle) revealed that the laying hens fed diet added 50, 100 and 150 mg Propolis/kg had significantly hyper responder compared to control-fed group. Concerning white blood cells differentiation, the present results speculated that the 100 and 150 mg Propolis supplementation significantly decreased heterophils count and increased lymphocytes count when compared with the control-group. In conclusion, it can be concluded from the above study that supplementation of Propolis at 100 or 150 mg is beneficial for improving the performance and immunity and for exploiting the full genetic potential of the commercial laying hens.

Key words: Laying hens, propolis, performance and immunocompetence

Introduction

Commercial laying hens are economic agriculture field production units in which the objective is to maximize field performance. Propolis, or "bee glue," is a well-known substance that beekeepers find in their hives. There are many factors affecting propolis composition such as collecting location, time and plant source (Greenaway *et al.*, 1991; Markham *et al.*, 1996). Propolis according to research has shown to be effective against a variety of bacteria (Velikova *et al.*, 2000), viruses (Amoros *et al.*, 1994), fungi (Murad *et al.*, 2002) and molds (Miyataka *et al.*, 1997). It has been shown to be a non-specific immunostimulant (Dimov *et al.*, 1991). Literature survey revealed that flavonoids, aromatic acids, diterpenic acids and phenolic compounds appear to be the principal components responsible for the

biological activities of Propolis samples. Flavonoids are found in most fruits and vegetables, especially citrus fruits, onions, apples, kale, broccoli, grapes, red wine and tea. They act as antioxidants, prevent blood clotting and protect veins, lowering levels of harmful estrogen (Middleton and Kandaswami, 1993; Hanasaki *et al.*, 1994; Duarte *et al.*, 1993). The effect of Egyptian Propolis on chicken body weight and lymphoid organs, Hegazi *et al.* (1996) revealed an increase in body weight after one week post injection and increase thymus weight after 14 days post injection up to the end of the experiment. While spleen weight slightly affected, but bursa and caecal tonsils weights were increased at the last two weeks of the experiment. The highest phagocytic activity was at 14 days post injection with Propolis (65% vs. 59%). Also, there was increase in the stimulation index

of the peripheral lymphocytes as detected by lymphocyte transformation in case of Propolis group. Hegazi *et al.* (1995) studied the effect of some bee products on immune response of chicken infected with virulent NDV. They found that, the mortality rate was reduced in-groups infected with virulent NDV and subsequently treated either with Propolis or honey if compared with the infected groups only. It was clear that, Propolis acts actively as antiviral agent than honey. The treatment with Propolis and honey of NDV infected chicken groups induced increase in the antibody titres and phagocytic percentage. Glinnik and Gapanovich (1981) showed that, in chickens, Propolis was effective against *S. aureus* and *S. epidermidis* in vitro. When Hubbard Golden comb hens were given a meal mixture containing 10, 20 and 30 mg Propolis/kg diet, Bonomi *et al.* (1976) reported that 30 mg Propolis in diet significantly increased egg production, egg weight, feed utilization and weight gain by 6.07, 1.27, 5.46 and 6% respectively, compared with control-group. In Ross broiler chicks, Shalmany and Shivazad (2006) indicated that average weight gain, feed consumption and feed efficiency were significantly higher for Propolis fed birds and inclusion of propolis also reduced mortality rate in comparison to control diet. Therefore, the present experiment was conducted to evaluate the efficacy of supplemental Propolis on productive performance and immune response of laying hens.

Materials and Methods

This experiment was carried out at poultry breeding farm, Poultry Production Department, Faculty of Agriculture, Ain Shams University. One hundred and twenty 46-wk-old, Hy-line White strain were randomly assigned to four groups of 30 hens each. The hens were individually housed in individually cages, on a 16-hour light schedule. Chickens in all groups were reared under the same environmental, managerial and hygienic conditions. Feed and water were supplied *ad libitum*. The hens received a typical layer diet containing 2800 ME kcal/kg and 18% CP to meet or slightly exceed the nutrient requirement recommended by NRC (1994). The hens were fed basal diet (control) or basal diet containing 50, 100 and 150 mg Propolis/kg. The composition and calculated chemical analysis of the experimental diet are presented in Table 1. The average high and low ambient temperatures recorded during the experimental period were 22.6 and 15.8°C, respectively.

Measurements and observations

Productive parameters: Body weight was individually recorded at 46, 50 and 54 weeks of age. Also, number and weight of eggs were recorded daily from the 46 to 54 weeks of age. Internal and eggshell quality was determined at 54 weeks of age. The egg length (long axis) and width (short axis) were measured with the

Table 1: Composition and calculated chemical analysis of the experimental diet

Ingredients	%
Ground yellow corn	61.80
Soybean meal 44%	19.30
Corn gluten meal	2.90
Decorticated cottonseed meal	2.00
Corn gluten feed	4.00
Bone meal	1.80
Limestone	7.42
Salt	0.32
Vitamin and mineral premix*	0.40
DL-Methionine	0.04
L-Lysine	0.02
Total	100
Calculated chemical analysis	
ME (kcal/kg)	2800
Crude protein (%)	18.00
Crude fat (%)	2.90
Crude fiber (%)	2.80
Calcium (%)	3.75
Available phosphorus (%)	0.40

*Each 2.5 kg of vitamin and minerals premix contain: vit. A, 12 mIU; vit. D₃, 4 mIU; vit E, 15g; vit. K, 2g; vit. B₁, 1g; vit B₂, 8g; vit. B₆, 6g; vit B₁₂, 10mg; niacin, 30g; biotin, 150mg; folic acid, 1g; pantothenic acid, 10g; choline chloride, 40mg; zinc, 60g; manganese, 70g; iron, 15g; copper, 5g; iodine, 1g; selenium, 0.15g.

electronic caliper. The width to length ratio was shown in percentage points and constituted the egg shape index. The height of thick albumen (H) and the egg weight (W) were used to calculate Haugh units from the formula of Williams (1997): $HU = 100 \log (H+7.7-1.7 W^{0.37})$, where H = thick albumen height, W = egg weight. Yolk diameter along the chalazae line was determined with the caliper (mm). The eggshell, after the removal of the egg content, was dried. Subsequently the eggshell was weighed to the nearest 0.01g. Eggshell thickness without inner membranes was measured (mm) with the micrometer. The albumen weight was calculated from the difference between the entire egg weight and the yolk and eggshell weight. The contents of yolk, albumen and the eggshell were expressed as percentages from the weight of a fresh egg. The breaking strength was measured according to Fathi and El-Sahar (1996) which assessed the resistance of the egg to crushing.

Blood constituents: At 54 weeks of age, blood samples were taken from the brachial vein into heparinized tubes for all birds. A portion of the fresh blood was used for hematocrit determination using capillary tubes and a microhematocrit centrifuge. The hematocrit Figures were measured after spinning microhematocrit for 12 min. Plasma was obtained from the blood samples by centrifugation for 10 min. at 4000 rpm and was stored at -20°C until the time of analysis. The frozen plasma was allowed to thaw at room temperature prior to analysis. Plasma calcium, phosphorus, total protein, albumin, cholesterol, GOT and GPT were determined by

Galal *et al.*: Dietary Propolis Supplementation

Table 2: Productive parameters of laying hens as affected by Propolis supplementation

	Age (wk)	Propolis (mg/kg)				Pooled SEM	Prob.
		0	50	100	150		
Body weight, g	46	1453.6	1450.2	1418.7	1453.9	51.45	NS
	50	1469.7	1480.9	1440.8	1484.5	67.18	NS
	54	1510.0	1500.3	1498.7	1512.2	57.83	NS
Egg number, no.	46-50	22.54 ^b	22.84 ^b	25.19 ^a	25.67 ^a	0.31	0.01
	50-54	21.43 ^c	22.32 ^b	25.12 ^a	25.81 ^a	0.25	0.01
	46-54	43.97 ^c	45.16 ^b	50.31 ^a	51.48 ^a	0.30	0.01
Egg weight, g	46-50	61.53 ^b	61.69 ^b	62.19 ^a	62.60 ^a	0.22	0.01
	50-54	61.72 ^b	61.82 ^b	62.93 ^a	63.10 ^a	0.23	0.01
	46-54	61.63 ^b	61.76 ^b	62.56 ^a	62.85 ^a	0.20	0.01
Egg mass, g	46-50	1386.9 ^a	1409.0 ^c	1566.6 ^b	1606.9 ^a	15.22	0.01
	50-54	1322.7 ^c	1379.8 ^c	1580.8 ^b	1628.6 ^a	21.14	0.01
	46-54	2709.9 ^c	2789.1 ^c	3147.4 ^b	3235.5 ^a	27.16	0.01
Feed consumption, g	46-50	3298.7 ^b	3268.9 ^b	3321.3 ^a	3310.3 ^a	62.14	0.01
	50-54	3227.3 ^b	3215.0 ^b	3335.6 ^a	3338.7 ^a	87.12	0.01
	46-54	6526.0 ^b	6483.9 ^b	6656.6 ^a	6649.0 ^a	75.86	0.01
Feed conversion ratio	46-50	2.38 ^a	2.32 ^a	2.12 ^b	2.06 ^b	0.02	0.01
	50-54	2.44 ^a	2.33 ^a	2.11 ^b	2.05 ^b	0.01	0.01
	46-54	2.41 ^a	2.32 ^a	2.11 ^b	2.05 ^b	0.02	0.01
Egg production rate	46-50	80.50 ^b	81.57 ^b	89.96 ^a	91.68 ^a	0.20	0.01
	50-54	76.54 ^c	79.71 ^b	89.71 ^a	92.18 ^a	0.32	0.01
	46-54	78.52 ^c	80.64 ^b	89.84 ^a	92.57 ^a	0.28	0.01

^{a,b,c} Means within the same row with different letters are significantly differ.

Table 3: Interior and eggshell quality parameters as affected by Propolis supplementation

Item	Propolis (g/kg diet)				Pooled SEM	Prob.
	0	50	100	150		
Egg weight, g	61.83 ^b	62.01 ^b	63.14 ^a	63.45 ^a	0.25	0.01
Albumen, %	60.61	60.22	60.11	59.43	0.87	NS
Yolk, %	29.52	29.86	29.77	30.15	0.23	NS
Haugh unit	81.56 ^b	84.12 ^a	84.17	85.63 ^a	1.14	0.01
Shape index	76.30	76.45	76.54	77.01	2.12	NS
Shell, %	9.87 ^b	9.92 ^b	10.12	10.42	0.36	NS
Shell thickness, mm	0.320 ^b	0.323 ^b	0.357 ^a	0.365 ^a	0.04	0.01
Breaking strength, kg/cm ²	3.17 ^b	3.26 ^b	3.65 ^a	3.71 ^a	0.08	0.01

^{a,b} Means within the same row with different letters are significantly differ.

enzymatic colorimetric methods using available commercial kits. The plasma globulin was calculated as the difference between plasma total protein and albumin.

In vivo cell - mediated immunity: A phytohemagglutinin-P (PHA-P) injection assay (Cheng and Lamont, 1988) was used to evaluate *in vivo* T-cell-mediated immune response of Hy-Line laying hens. Birds were injected intradermally in the wattle with 0.5 mg of PHA-P (Sigma Chemical Co., St. Louis, Missouri) in 0.1 ml of phosphate buffered saline (PBS) after marking the injection site. The thickness of wattle was measured (to nearest 0.01mm) at 0, 24, 48 and 72hrs after PHA-P injection. Wattle swelling was calculated as the difference between the thickness of the wattle prior to and after injection of PHA-P.

Heterophils / lymphocytes ratio: At 54 week of age, blood samples were obtained from each treatment for heterophil (H) and lymphocyte (L) enumeration based on the procedures of Gross and Siegel (1983). Briefly, one drop of blood being smeared on each of glass slides. The smears were stained using Wright's stain. Two hundred leukocytes, including granular (heterophils) and nongranular (lymphocytes) ones, were counted on different microscopic fields representing 200 cells and the heterophil to lymphocyte ratio was calculated.

Statistical analysis: Data were subjected to a one-way analysis of variance with treatment group effect using the General Linear Model (GLM) procedure of SAS User's Guide, 2001. When significant differences among means were found, means were separated using Duncan's multiple range tests.

Results and Discussion

Productive parameters: Body weight, egg production parameters, feed consumption and feed conversion ratio of laying hens fed different levels of Propolis are summarized in Table 2. The present results revealed that there was no significant difference among treated groups for body weight. However, supplemental Propolis at 100 and 150 mg significantly increased number and weight of eggs compared to control-group. These reflected on egg mass, whereas the hens fed diet containing 100 or 150 mg Propolis produced significantly heaviest egg mass by 437.5 and 525.7 g, respectively compared to control-group. Hubbard Golden

Galal *et al.*: Dietary Propolis Supplementation

Table 4: Hematological parameters of laying hens fed different levels of propolis.

Item	Propolis (mg/kg diet)				Pooled SEM	Prob.
	0	50	100	150		
Hematocrit level, %	32.15 ^b	32.56 ^b	33.12 ^a	33.42 ^a	1.12	0.01
Total protein, g/dl	5.67 ^b	5.68 ^b	6.18 ^a	6.25 ^a	0.23	0.01
Albumin, g/dl	3.57	3.44	3.51	3.23	0.10	NS
Globulin, g/dl	2.10 ^b	2.24 ^b	2.67 ^a	3.02 ^a	0.06	0.01
Calcium, mg/dl	21.45	21.46	22.10	22.32	0.98	NS
Phosphorus, mg/dl	10.12	10.43	10.51	10.17	0.45	NS
Cholesterol, mg/dl	132.14 ^a	131.12 ^a	125.46 ^b	119.18 ^c	3.14	0.01
GOT, U/L	42.16 ^a	41.35 ^a	39.13 ^b	38.75 ^b	1.10	0.01
GPT, U/L	12.14 ^a	12.10 ^a	10.48 ^b	10.52 ^b	0.65	0.01

^{a,b,c}Means within the same row with different letters are significantly differ

comb hens aged 5.5 months were given a meal mixture without and with Propolis (10, 20 and 30 mg/kg diet), Bonomi *et al.* (1976) reported that supplemental 30 mg Propolis in diet significantly increased egg production, egg weight, feed utilization and weight gain by 6.07, 1.27, 5.46 and 6.0% respectively, compared with control group. With respect to feed consumption and feed conversion ration, the present results indicated that the laying hens fed diet containing 100 and 150 mg Propolis/kg diet were significantly consumed more feed than control-group. Similar trend, but not statistically significant, was observed for hens fed diet added 50 mg Propolis. Moreover, the increase feed intake and egg mass in Propolis groups, resulting in significantly improve feed conversion ratio compared to control-group. Shalmany and Shivazad (2006) reported that chicks fed Propolis containing diets consumed significantly higher feed. Bonomi *et al.* (1976) found an increase in feed intake when laying hens were fed Propolis versus control groups. Also, they concluded that increase in feed intake in the Propolis groups may be due to improved birds health and higher palatability of Propolis diets due to mixture of resin, wax, honey and vanillin content of Propolis. Ghisalberti (1979) report additional weight gains for broiler chickens of up to 20% when 500 ppm of propolis was added to their diets. They said that this improved effect is partially due to its high content of flavonoids and increase feed intake of Propolis diets than the control. Experimental work of Buhatel *et al.* (1983) showed that Propolis supplementation to the ration of pullets improved feed conversion. This effect is due to high content of flavonoids and healthy conditions of birds fed Propolis.

Internal and eggshell quality: Internal and eggshell quality as affected by Propolis supplementation is presented in Table 3. The egg weight was significantly affected by supplemental Propolis at 100 and 150 mg compared to control-group. Similar trend, but not statistically significant, was observed for 50 mg Propolis. There was no significant difference among treated groups for both albumen and yolk percentages. The Haugh units were significantly affected by Propolis

supplementation, whereas the eggs produced from laying hens fed diet containing Propolis at all levels recorded the highest Haugh units compared to control-group. With respect to eggshell quality, it could be observed that both 100 and 150 mg supplemental Propolis significantly increased the eggshell percentage compared to control-group. Similar trend was noticed for eggshell thickness, whereas the eggs produced from hens fed diet added 100 or 150 mg Propolis recorded highest shell thickness compared to other group. In accordance to breaking strength, the present results showed that the breaking strength of eggs produced from hens fed diet containing 100 or 150 mg Propolis significantly highest than those of other groups. It could be concluded that the supplemental Propolis at 100 or 150 mg improved the eggshell quality.

Hematological parameters: Effect of supplemental Propolis on some blood parameters of laying hens are summarized in Table 4. The higher hematocrit level may have enhanced oxygen delivery to the tissue (Zongo and Petitjean, 1990). Also, this increase is supposed to be caused by increased blood volume as a reaction to increasing body oxygen requirement. The present result revealed that there was no significant difference between control and 50 mg propolis diet for hematocrit level. Inversely, the hematocrit level of hens fed diet containing 100 or 150 mg propolis was significantly higher than that of control-group. Also, it could be noticed that supplemental Propolis increased plasma calcium and phosphorus compared to control-group, but the difference did not statistically significant. With respect to plasma total protein, it could be speculated that the supplemental propolis at 100 or 150 mg significantly increased plasma total protein compared to control group. Similar reports were drawn by Giurgea *et al.* (1981). They indicated that daily administration of propolis extract to chickens changed the blood concentration of cholesterol, total proteins and amino acid. Also, Propolis stimulated mammalian tissue regeneration, as it caused strong activation of mitosis of cells cultured in vitro and it enhanced protein biosynthesis (Scheller *et al.*, 1977 and Gabrys *et al.*,

Table 5: Wattle swelling (difference) of laying hens fed different levels of propolis

Time	Propolis (mg/kg diet)				Pooled SEM	Prob.
	0	50	100	150		
24 hr post PHA-P injection	0.31 ^d	0.42 ^c	0.54 ^b	0.68 ^a	0.02	0.001
48 hr post PHA-P injection	0.22 ^d	0.36 ^c	0.42 ^b	0.52 ^a	0.04	0.001
72 hr post PHA-P injection	0.08 ^d	0.19 ^c	0.30 ^b	0.36 ^a	0.01	0.001

^{a-d} Means within the same row with different letters are significantly differ.




Fig. 1: Wattle swelling of laying hens as affected by propolis supplementation.




Fig. 2: Heterophils (H), lymphocytes (L) and H/L ratio of laying hens fed different level of Propolis.

1986). Similar trend was not observed for albumin, whereas there was no significant difference among treated groups for plasma albumen. Concerning plasma globulin, our result revealed that the plasma globulin was significantly increased when the propolis added to the diet at 100 or 150 mg. The globulins are composed of three fractions, designated alpha, beta and gamma. Alpha-globulins are a group of proteins manufactured almost entirely by the liver. Normally, these proteins increase with acute nephritis, severe active hepatitis, active, usually systemic inflammation, malnutrition and in nephritic syndromes (Margaret, 2001). The gamma-globulin fraction contains most of the immuno-proteins, including IgM, IgA, IgE and IgG. These usually elevate with ongoing antigenic stimulation, usually from infectious agents (Margaret, 2001).

There was no significant difference between 0 and 50 mg Propolis for plasma cholesterol. Inversely, the supplemental propolis at 100 and 150 mg significantly reduced plasma cholesterol compared to control-group. With respect to liver function, it could be noticed that supplemental Propolis at 100 and 150 mg significantly reduced GOT and GPT concentration compared to control-group. Hegazi *et al.* (1997) showed that, administration of Egyptian and Bulgarian propolis induces an antibacterial activity *in vivo* as well as *in vitro*. The ethanolic extract of propolis has a weak general effect on estimated parameters in normal rats and it is not a toxic substance. Both types of propolis exerted an anabolic effect for protein synthesis by liver cells. Both

types of infections with *S. aureus* and *E. coli* caused an increase in the activity in serum AST and ALT and consequently decrease their activity in the liver. On the other hand, the activity of ALT and AST returned to the control level after administration of propolis in rats infected with *S. aureus* and *E. coli*.

Immunocompetence parameters

***In vivo* cell - mediated immunity:** The PHA intradermally reaction, a T-lymphocyte-dependent response, has been well researched and has been shown to be a reliable indicator of *in vivo* cellular immunity in poultry (Goto *et al.*, 1978; McCorkle *et al.*, 1980). The skin response reflects a complex series of physiological events such as mitogen-receptor and lymphocyte-macrophage interactions, release of chemical mediators, cellular proliferation and changes in vascularity (Chandra and Newberne, 1977). Histologically, PHA is strongly mitogenic to T-lymphocytes and intradermal injections elicit macrophage infiltration and dense perivascular accumulations of lymphocytes 24h post-injection in chickens (Goto *et al.*, 1978; McCorkle *et al.*, 1980). The increased infiltration by basophils and eosinophils 24h post-injection has been described as a cutaneous basophil hypersensitivity response (Stadeckerm *et al.*, 1977). *In vivo* cell-mediated immune response as measured by PHA stimulation (wattle) is presented in Fig. 1 and Table 5. It could be noticed that the hens fed diet added 50, 100 and 150 mg Propolis/kg diet had significantly hyper responder to PHA-P injection

compared to control-fed group. Propolis according to research has shown to be effective against a variety of bacteria, viruses, fungi and molds. It has been shown to be a non-specific immunostimulant. The delayed hypersensitivity skin test using propolis as sensitizing antigen showed specific stimulation to propolis after 72 hours after inoculation with specific antigen. Egyptian propolis gave the typical delayed hypersensitivity when inoculated to the sensitized chickens. The thickness index was 0.90 mm thickness if compared with non-sensitized control group 0.12 mm thickness (Hegazi *et al.*, 1996). Hegazi *et al.* (1995) studied the effect of some bee products on immune response of chicken infected with virulent NDV. They found that, the mortality rate was reduced in-groups infected with virulent NDV and subsequently treated either with propolis or honey if compared with the infected groups only. It was clear that, propolis acts actively as antiviral agent than honey. The treatment with propolis and honey of NDV infected chicken groups induced increase in the antibody titres and phagocytic percentage. The inoculation of different antigens in the footpad of sensitized and non-sensitized chickens induced different degrees of footpad thickness as well as cellular and vascular reaction depending on the type of inoculation with NDV antigen.

White blood cells differentiation: White blood cells differential count for laying hens fed different levels of propolis are presented in Fig. 2. It could be noticed that there was no significant difference between control and low level of propolis supplementation for both heterophils and lymphocytes count. Conversely, the medium and high level of propolis supplementation significantly decreased the heterophils count and increased the lymphocytes count when compared with the control-group. In birds, the heterophil are phagocytic cells whose main is protection against invading microorganisms, whereas primary functions of lympho-involve cell-mediated and humoral immunity. Heterophils increase and lymphocytes decrease when are stressed, so that the ratio between them is a index of response to a stressor (Gross and Siegel, 1985). In accordance to H/L ratio, our results showed that the propolis supplementation at 100 and 150 g/kg diet significantly decreased the H/L ratio of laying hens. The H/L ratio is a recognized measure of stress in birds (Davison *et al.*, 1983; Gross and Siegel, 1983; Maxwell, 1993) that has become a valuable tool in stress research especially when combined with the convenience and repeatability of automated blood cell counts.

In conclusion, it can be concluded from the above study that supplementation of Propolis at 100 or 150 mg is beneficial for improving the performance and immunity and for exploiting the full genetic potential of the commercial laying hens.

References

- Amoros, M., E. Lurton, J. Boustie, L. Girre, F. Sauvager, and M. Cormier, 1994. Comparison of the anti-Herpes Simplex Virus activities of propolis and 3-methyl-but-2-enyl caffeate. *J. Nat. Prod.*, 57: 644-647.
- Bonomi, A., F. Morletto and M. Binachi, 1976. Propolis in feeds for laying hens. *Avicoltura*, 54: 43-54.
- Buhatel, T., S. Vesa, A. Dimitrin and I. Moldovan, 1983. Contribution to knowledge of the stimulative effect of propolis on piglet and pullets. *Buletinul Institutului Agronomic Cluj Napoca Zootehnie Medicina Veterinara*, 37: 45-48.
- Chandra, R.K. and P.M. Newberne, 1977. Nutrition, immunity and infections. Plenum Press, New York.
- Cheng, S. and S.J. Lamont, 1988. Genetic analysis of immunocompetence measures in a white Leghorn chicken line. *Poult. Sci.*, 67: 789-995.
- Davison, T.F., L.G. Rowell and J. Rea, 1983. Effects of dietary corticosterone on peripheral blood lymphocyte and granulocytes populations in immature domestic fowl. *Res. Vet. Sci.*, 34: 236-239.
- Dimov, V., N. Ivanovska, N. Manolova, V. Bankova, N. Nikolov and S. Popov, 1991. Immunomodulatory action of propolis. Influence on anti-infectious protection and macrophage function. *Apidologie*, 22: 155-162.
- Duarte, J., E.P. Vizcaino, P. Utrilla, J. Jimenez, J. Tamargo and A. Zarzuelo, 1993. Vasodilatory effect of flavonoids in rat aortic smooth muscles. Structure activity relationships. *Biochem. Pharmacol.*, 24: 857-862.
- Fathi, M.M. and E.A. EL - Sahar, 1996. Determining the strength of eggshell by using an appropriate apparatus and an equation to calculate egg surface depending on its dimensions. *Egypt. Poult. Sci.*, 16: 285-303.
- Gabrys, J., Z. Konecki, W. Krol, S. Scheller and J. Shani, 1986. Free amino acids in bee live product (propolis) as identified and quantified by gas - Liquid chromatography. *Pharmacol. Res. Commun.*, 18: 513-518.
- Ghisalberti, E.L., 1979. Propolis: A review. *Bee World*, 60: 59-84.
- Giurgea, R., V. Toma, H. Popescu and C. Linicencu, 1981. Effects of standardized propolis extracts on certain blood constituents in chickens. *Clujul - Med.*, 54: 151-154.
- Glinnik, A.V. and V.Y.A. Gapanovich, 1981. Antibacterial properties of propolis. *Zhurnal Ushnykh Nosovykh Gorlovykh Boleznj No.*, 4: 75-76.
- Goto, N., H. Kodama, K. Okada and Y. Fujimoto, 1978. Suppression of phytohemagglutinin skin response in thymectomized chicken. *Poult. Sci.*, 57: 246-250.

- Greenaway, W., J. May, T. Scaysbrook and F.R. Whatley, 1991. Identification by gas chromatography-mass spectrometry of 150 compounds in propolis. *Zeitschrift fur Naturforschung*, 42c: 111-121.
- Gross, W.B. and P.B. Siegel, 1983. Evaluation of the heterophil/lymphocyte ratio as a measure of stress in chickens. *Avian Dis.*, 27: 972-979.
- Gross, W.B. and P.B. Siegel, 1985. Selective breeding of chickens for corticosterone response to social stress. *Poult. Sci.*, 64: 2230-2233.
- Hanasaki, Y., S. Ogawa and S. Fukui, 1994. The correlation between active oxygen scavenging and antioxidative effects of flavonoids. *Free Radical Biol. Med.*, 16: 845-850.
- Hegazi, A.G., H.F. El Miniawy and F.A. El Miniawy, 1995. Effect of some honeybee products on immune response of chicken infected with Virulent NDV. *Egyptian J. Immunol.*, 2: 79-86.
- Hegazi, A.G., M. Khalefa and E. Toussun, 1997. Influence of various types of propolis on some biochemical changes of normal and bacterial infected rats. *International Symposium On Apitherapy*, Cairo 8-9th, March, 1997.
- Hegazi, A.E., F.A. El Miniawy and F.K. Abd El Hady, 1996. Influence of administration of propolis on chicken immune status. *The Egypt. J. Immunol.*, 3: 111-116.
- Margaret, A.W., 2001. *Avian Plasma Proteins*. <http://www.exoticpetvet.net>.
- Markham, K.E., K.A. Mitchel, A.L. Wilkins, J.A. Daldy and Y. Lu, 1996. HPLC and GC-MS identification of the major organic constituents in New Zealand propolis. *Phytochemistry*, 42: 205-211.
- Maxwell, M.H., 1993. Avian blood leukocyte responses to stress. *World's Poult. Sci. J.*, 49: 34-43.
- McCorkle, F., I. Olah and B. Glick, 1980. The morphology of the phytohemagglutinin-induced cell response in the chicken's wattle. *Poult. Sci.*, 59: 616-623.
- Middleton, E. and C. Kandaswami, 1993. The impact of plant flavonoids on mammalian biology: Implications for immunity, inflammation and cancer. In *the flavonoids: Advances in Research Science 1986*. J.B. Harborne, (Ed.). Chapman and Hall, London, UK., pp: 619-652.
- Miyataka, H., M. Nishiki, H. Matsumoto, T. Fujimoto, M. Matsuka and T. Satoh, 1997. Evaluation of propolis. 1. Evaluation of Brazilian and Chinese propolis by enzymatic and physico-chemical methods. *Biol. Pharm. Bull.*, 20: 496-501.
- Murad, J.M., S.A. Calvi, A.M.V.C. Soares, V. Bankova and J.M. Sforcin, 2002. Effect of propolis from Brazil and Bulgaria on fungicidal activity of macrophage against *paracoccidioides brasiliensis*. *J. Ethnopharm.*, 79: 331-334.
- National Research Council, 1994. *Nutrient Requirements of Poultry*. 9th Revised Edn. National Academy Press, Washington, DC.
- SAS Institute, 2001. *SAS/STAT User's Guide Version 8.2 Edn. Statistics*. SAS Institute Inc., Cary, NC.
- Scheller, S., E. No Lewajka, M. Panasiewicz, D. Dzieka, J. Tustanowski and A. Stojko, 1977. Biological properties and clinical application of propolis. IV. The action of ethanol extract of propolis on cells cultured *in vitro*. *Arzneim Forsch.*, 27: 1547-1549.
- Shalmany, S.K. and M. Shivazad, 2006. The effect of diet Propolis supplementation on Ross broiler chicks performance. *Int. J. Poult. Sci.*, 5: 84-88.
- Stadeckerm, J., M. Luk, C.A. Dvorak and S. Leskowitz, 1977. The cutaneous basophil response to phytohemagglutinin in chickens. *J. Immunol.*, 118: 1564-1568.
- Velikova, M., V. Bankova, I. Tsvetkova, A. Kujumgiev and M.C. Marcucci, 2000. Antibacterial ent-kaurene from Brazilian propolis of native stingless bees. *Fitoterapia*, 71: 693-690.
- Williams, K.C., 1997. Some factors affecting albumen quality with particular reference to Haugh unit score. *World's Poult. Sci. J.*, 48, 1: 5-16.
- Zongo, D. and M. Petitjean, 1990. Effects associated with the Na (naked neck) gene in the domestic cock. I. live weight, comb, haematocrit, sexual maturity and semen characters in temperature environment. *Bulletin of Anim. Health Prod. Afr.*, 38: 3, 259-263.