



Journal of Applied Sciences

ISSN 1812-5654

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Effect of Human Activities and Oil Pollution on the Microbiological and Physicochemical Quality of Udu River, Warri, Nigeria

H.S.A. Aluyi, F.O. Ekhaize and D.M. Adelusi

Department of Microbiology, Faculty of Science, University of Benin, Benin City, Nigeria

Abstract: The effect of human activities and oil pollution on the microbiological quality of Udu River was investigated from September 2002 to January 2003. The microbiological parameters were carried out using the standard microbiological techniques. The bacterial and fungal loads were shown to be highest in the inhabited stations (2 and 3) with downstream (station 4) suffering from dilution effects of human activities and upstream (station 1) showing lower counts. Stations 2 and 3 were shown to record the highest heterotrophic microbial load 10.7×10^8 cfu mL⁻¹ and 10.4×10^8 cfu mL⁻¹ and 8.0×10^8 cfu mL⁻¹ and 9.0×10^8 cfu mL⁻¹ for bacterial and fungal isolates respectively. Total coliform counts ranges from $1.5 - 4.3 \times 10^6$ and $1.4 - 2.8 \times 10^6$ MPN100 mL⁻¹ for stations 2 and 3. Microorganism isolated and characterized includes nine bacterial and four fungal genera. The bacterial isolates were *Staphylococcus*, *Streptococcus*, *Proteus*, *Escherichia*, *Clostridium*, *Acetobacter*, *Salmonella*, *Pseudomonas*, *Micrococcus* and the predominant fungal isolates includes, *Aspergillus*, *Penicillium*, *Geotrichum* and *Helminthosporium*. The concentrations of total suspended solids, Biochemical Oxygen Demand (BOD₅), conductivity, total iron, total solids, total copper and total dissolved solids were shown to be higher in stations 2 and 3 than station 1 which have no direct human activities. Similarly, the values obtained for salinity, total hardness and Dissolved Oxygen (DO) of the inhabited stations were shown to be lower than station 1. Correlation coefficient shows positive relationship between the different microbial populations and some physicochemical parameters studied.

Key words: Human activities, oil pollution, heavy metals

INTRODUCTION

Water constitutes an integral part of man's life as well as the earth's surface, hence the need for continuous research about impact on its quality due to one form of human activity or the other. Of all the water found on earth, marine constitutes the largest with about 97%. Fresh waters, although a small part of the waters on earth, are extremely important as source of drinking water. In many locations the contamination of surface and subsurface waters by domestic and industrial wastes causes environmental problems (Prescott *et al.*, 2000).

The Udu River which is a tributary of the Warri River is located in Warri South Local Government Area of Delta State and spans over 48 km². The river has its source from Aboh in Delta north, lying between longitude 6.20°N and latitude 5.45°E. The river flows southwest to Warri where it is joined by a large number of creeks from where it empties into the very brackish Forcados River and finally into the Atlantic Ocean (Udo, 1975).

The river flows across a large uninhabited stretch of land. Companies involved in one form of marine activity

or the other basically occupy the inhabited expanse of land. This rapid industrialization has led to alterations in the water quality of the river thus causing great concern for the immediate users of the river water and the larger society. The location of a jetty where people take boat straight from Owina to Igbudu (a central position of Warri main town) thereby avoiding the numerous traffic in the town has resulted in increased human activities along the course of the river. This is evident in higher faecal contamination of the river, urban run offs, abattoir effluent discharges and organic and inorganic dumps around or directly into the river. The dumping of chemicals by companies like Nissco and Globester involved in the fabrication of oilrig platforms, has further led to great alterations of the bacteriological and physicochemical quality of the river.

The importance of water to humans has resulted in the sitting of most rural communities and industries as well as individual homes along the river course (Egborge, 1982). Large amounts of water consumed by humans are from surface water, which includes rivers, streams, lakes, wetlands, groundwater

(Thurman *et al.*, 1998). Water pollution is basically rendering the water unfit for human consumption and recreational purpose. The Udu River serves communities like Ujevuwu, Egin, Aladja and Ovwian where fishing serves as a major source of occupation for most inhabitants of these communities. Other activities like swimming, bathing, dumping of sawdust and washing of clothes by the river are also a common feature of water pollution observed here. Thurman *et al.* (1998) reported similar results where inhabitants in the rural communities rely on untreated natural water for use, especially in the developing countries of the world.

Increased human activities, the accidental and intentional dumping of pollutants, effluent discharges into this river has created growing environmental concerns, especially loss of commercial fisheries, contaminated seafood and species extinctions. This alteration in the water quality has increased the anxiety of the rural inhabitants, who have to travel great distance to other creeks to fish as well as shortage of good quality drinking water. Types and concentrations of contaminants in drinking water differ with location. In surface water the difference can also be attributed to nature of contaminants around the water body (Miller, 2001). Toxic chemicals and microbial pollutants are widespread. Lead, commonly used in plumbing conduit and fixtures until recently, was a serious contaminant especially to foetus and children. Anson and Ware, (1974) stated that the bacterial population in water is greatly influenced by oxygen, pH and temperature of a water body being one of the most important characteristics which determines to a considerable extent, the trends and tendencies of changes in its quality (Anonymous, 1971). This paper reports the effect of human activities and oil pollution on the microbiological and physicochemical qualities of Udu River, Warri, Nigeria.

MATERIALS AND METHODS

Source of samples: Water samples were collected between September, 2002 and January, 2003 from four sampling stations designated 1 to 4 along the course of the river. Station 1 is an upstream and uninhabited with less human activities. The vegetation around this station is composed mainly of trees with branches extending and creating shade around the immediate bank of the river. Station 2 is 1 km downstream of station 1 and is located just by the Udu Bridge, the water here is polluted with high human activities like swimming, washing of clothes and logs, as well as discharge from boats buildings. Station 3 is about 1.2 km downstream of Station 2, with a moderate flow rate and less human activities. The

vegetation around this station is composed mainly of thick grasses, trees and others. The water is turbid, probably due to the effect of human activities such as disposal of sawdust from upstream. Station 4 is about 500 m downstream of station 3. It has a consistent flow rate with no direct human activity. The water is less turbid, clearer than upstream. The vegetation is composed mainly of thick palm trees and grasses.

Sample collection: Samples for bacteriological analysis were collected directly from the river in sterilized plastic containers. The cover of the bottle was removed and held in one hand while the bottle was held at the base with the other hand. It was then plunged neck downwards below the surface of the water. It was tilted until the neck pointed slightly upwards, with the mouth being directed towards the current. Samples for Dissolved Oxygen (DO) and Biochemical Oxygen Demand (BOD) were collected in 250 mL bottles with stoppers, one millimeter each of Winkler's solutions A and B were added to samples on site to fix the oxygen (Anonymous, 1989). The samples were collected between 7.00 and 9.00 am and were transported to the laboratory for both bacteriological and physicochemical analyses and were analyzed with 6 h of collection.

The bacteriological parameters monitored include total viable aerobic count to isolate heterotrophic bacterial and fungal, total coliform counts, *Escherichia coli* counts, faecal *Streptococci* counts and *Clostridium perfringens* counts according to the methods of Gerhardt *et al.* (1994). The isolation and identification of bacterial and fungal isolates were carried out in accordance with Bergey's Manual of Determinative Bacteriology (Buchanan and Gibbons, 1974; Barnett and Hunter, 1972; Gerhardt *et al.*, 1994). The physicochemical parameters were studied in accordance with the standard methods of APHA (Anonymous, 1989).

RESULTS

The highest total heterotrophic bacteria population of 10.7×10^8 cfu mL⁻¹ was recorded in late January at downstream station 2 and the lowest count of 5.3×10^8 cfu mL⁻¹ was recorded in the upstream station 1 in mid September. The highest aerobic pathogenic bacterial densities 9.1×10^8 cfu mL⁻¹ was recorded in January at downstream station 2 and the lowest 3.2×10^8 cfu mL⁻¹ in September in station 1. It was observed that both total heterotrophic and aerobic pathogenic bacterial population increased along the stations, from station 1 to station 3 with a short drop in station 4 (Table 1).

Table 1: Total bacterial and fungal population of Udu River

Station	Total heterotrophic bacteria x 10 ⁸ cfu mL ⁻¹	Total aerobic pathogenic bacteria x 10 ⁵ cfu mL ⁻¹	Total hydrocarbon degraders x 10 ⁸ cfu mL ⁻¹	Total coliform x 10 ⁶ MPN100 mL ⁻¹	Total <i>E. coli</i> x 10 ⁶ MPN100 mL ⁻¹	Total <i>Strept. faecalis</i> x 10 ⁴ MPN100 mL ⁻¹
1	5.3-6.7	3.2-4.5	3.1-4.5	1.1-2.2	0.9-1.4	0.7-2.1
2	8.6-10.7	6.1-9.1	3.6-5.4	1.5-4.3	1.1-3.9	1.5-3.9
3	8.7-10.4	6.8-8.6	3.6-5.3	1.4-2.8	0.9-3.9	0.9-3.9
4	7.4-9.5	6.1-7.6	2.6-4.4	0.7-1.4	0.7-2.1	0.7-2.8

Continued

Station	Total <i>C. perfringens</i> x 10 ⁴ MPN100 mL ⁻¹	<i>A. niger</i> x 10 ⁸ cfu mL ⁻¹	<i>A. flavus</i> x 10 ⁸ cfu mL ⁻¹	<i>Penicillium</i> sp. x 10 ⁸ cfu mL ⁻¹	<i>G. candidum</i> x 10 ⁸ cfu mL ⁻¹	<i>Helminthosporium</i> sp. x 10 ⁸ cfu mL ⁻¹
1	1.1-2.1	2.0-7.5	1.0-6.0	3.0-7.0	1.0-6.0	0-3.0
2	1.5-2.8	2.0-6.0	2.0-4.0	3.0-8.0	0-4.0	0-3.0
3	1.4-2.8	3.0-7.0	1.0-6.0	4.0-9.0	1.0-5.0	1.0-4.0
4	0.9-2.8	0-3.0	1.0-3	1.0-5.0	0-4.0	0-3.0

Table 2: Occurrence and spatial distribution of microbial isolates in Udu River

Isolate	Station	Sept.	Sept.	Oct.	Oct.	Nov.	Nov.	Dec.	Dec.	Jan.	Jan.
<i>S. aureus</i>	1	-	-	+	-	-	+	+	-	+	-
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	-	+	-	+	+	+	+	-
<i>Proteus</i> sp.	1	+	+	-	-	-	-	-	+	-	-
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	-	-	+	+	+	+	-	-
<i>S. typhi</i>	1	-	-	-	-	+	-	-	-	-	+
	2	+	+	-	+	+	+	+	-	+	+
	3	+	-	+	+	+	-	-	-	+	+
	4	+	-	+	+	-	-	+	-	-	+
<i>C. perfringens</i>	1	+	+	+	+	+	+	+	+	+	+
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	+	+	+	+	+	+	+	+
<i>S. faecalis</i>	1	+	+	+	+	+	+	+	+	+	+
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	+	+	+	+	+	+	+	+
<i>Pseudomonas</i> sp.	1	+	-	+	+	+	-	+	+	+	+
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	+	+	+	+	+	+	+	+
<i>Microoccus</i> sp.	1	-	-	+	-	+	+	-	-	-	-
	2	+	+	+	+	-	+	+	+	-	+
	3	+	+	-	-	+	+	+	+	+	-
	4	+	+	-	+	+	+	-	-	+	+
<i>Acetobacter</i> sp.	1	-	-	-	+	-	-	+	-	-	-
	2	+	-	+	+	+	+	-	+	+	-
	3	+	+	-	+	-	-	+	+	-	+
	4	-	-	+	+	-	+	-	-	+	+
<i>A. niger</i>	1	+	+	+	+	+	+	+	+	+	+
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	-	+	+	+	-	+	+	+	+
<i>A. flavus</i>	1	+	+	+	+	+	+	+	+	+	+
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	+	+	+	+	+	+	+	+
<i>Penicillium</i> sp.	1	+	+	+	+	+	+	+	+	+	+
	2	+	+	+	+	+	+	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	+	+	+	+	+	+	+	+
<i>G. candidum</i>	1	+	-	-	+	+	+	+	+	+	+
	2	-	+	+	+	+	-	+	+	+	+
	3	+	+	+	+	+	+	+	+	+	+
	4	+	+	-	+	+	+	-	-	+	+

+ = Present, - = Not detected

Table 3: Mean±SE of the physicochemical properties of sample stations, Sept. 2002-Jan., 2003

Properties physical	Station 1 n (10)	Station 2	Station 3	Station 4
Atmos. temp.	27.7±0.76	28.5±0.84	27.9±0.83	27.95±0.684
Water temp. (°C)	26.5±0.69	27±0.75	26.45±0.807	26.75±0.573
Water velocity (°C)	0.721±0.04	0.59±0.023	0.57±0.025	0.498±0.020
Total suspended solids (mg L ⁻¹)	24.03±0.84	67.35±77.93	59.75±7.3	52.22±7.312
Total dissolved solids (mg L ⁻¹)	42.98±3.45	105.06±6.51	97.31±6.22	88.37±6.213
Total solids	67.01±4.24	172.35±14.31	157.01±13.27	140.35±13.31
pH	6.73±0.029	6.15±0.049	6.14±0.048	6.52±0.066
Conductivity	51±2.67	92.8±6.17	84.7±4.876	70.8±5.280
Hardness	12.08±0.12	10.81±0.029	10.52±0.105	10.24±0.096
Dissolved oxygen	9.1±0.35	7.68±0.0228	7.08±0.267	8.1±0.301
Biological oxygen demand	5.07±0.19	6.83±0.122	6.28±0.154	6.37±0.109
Nitrate	0.156±0.019	0.157±0.011	0.132±0.008	0.085±0.005
Phosphate	0.028±0.023	0.218±0.016	0.18±0.015	0.138±0.012
Sulphate	0.39±0.032	0.38±0.010	0.36±0.019	0.326±0.016
Iron	0.25±0.013	0.545±0.016	0.537±0.018	0.352±0.010
Lead	0.435±0.028	0.29±0.007	0.28±0.013	0.226±0.01
Chromium	0.029±0.004	0.03±0.05	0.03±0.005	0.015±0.002
Copper	0.43±0.013	0.49±0.017	0.50±0.019	0.42±0.009
Manganese	0.39±0.003	0.049±0.0043	0.05±0.004	0.023±0.003
Zinc	0.465±0.021	0.577±0.018	0.56±0.016	0.495±0.016
Cadmium	0.071±0.005	0.087±0.008	0.09±0.007	0.063±0.008

The total hydrocarbon degrading microbial isolates showed a similar pattern as total heterotrophic and aerobic pathogenic bacteria, the microbial counts ranges from 3.1×10^8 cfu mL⁻¹ – 5.4×10^8 cfu mL⁻¹. The highest count 5.4×10^8 cfu mL⁻¹ was recorded in downstream station 2 in late January and the lowest value was recorded in early September at station 4.

The total coliform counts were observed to fluctuate throughout the sampling period. The results showed that the total coliform population was lowest (0.7×10^6 MPN100 mL⁻¹) in mid September at station 4 while the highest density (4.3×10^6 MPN100 mL⁻¹) was recorded in January at station 2. Similar trend was observed in the population densities of faecal streptococci *Clostridium perfringens* in all the water-sampling stations. The population of *E. coli* in all the water sampling stations was moderate but significantly lower than that of coliforms. There was variation in the fungal population, which ranges from species to species, as some species of fungi were found more in uninhabited station than inhabited stations. *Aspergillus niger* and *Geotrichum candidum* were seen as the most predominant fungal species. Eleven microbial isolates were identified and characterized and these include nine bacterial and four fungal genera, *Staphylococcus*, *Streptococcus*, *Proteus*, *Escherichia*, *Clostridium*, *Acetobacter*, *Salmonella*, *Pseudomonas*, *Micrococcus*, *Aspergillus*, *Penicillium*, *Geotrichum* and *Helminthosporium*. The eleven microbial isolates were frequently isolated from the upstream-downstream station. The results of the occurrence and distribution of the microbial isolates is showed in Table 2. Station 1 showed a scanty distribution of the bacterial isolates, with the exception of *C. perfringens*. Stations 2 and 3 were seen to be heavily

loaded with the bacterial isolates, which is an evidence of the high human activities and the oil pollution effects of the different oil companies operating in the area. Among the fungal isolates, *Aspergillus niger*, *A. flavus* and *Penicillium* sp. were predominantly present in the first three stations. The physicochemical qualities of Udu River studied are showed in (Table 3). During the periods of sampling, the flow rate ranged from 0.498 to 0.721 ms⁻¹. The flow rate was highest at station 1 (0.721 ms⁻¹) and lowest at station 4 (0.498 ms⁻¹). Air and water temperature varied from 27.7-28.5°C and from 26.45 to 27.0°C. The pH value of water varied from 6.14 to 6.73. The Total Dissolved Solids (TDS), Total Suspended Solids (TSS) and Total Solids (TS) values varied from 24.03 to 67.35; from 42.98 to 105.06 and from 67.01 to 172.35, respectively with stations 2 and 3 recording the highest values (Table 3). In contrast, conductivity, hardness, Dissolved Oxygen (DO), nitrate, sulphate, lead, chromium, cadmium decreased from September to January along the sampling stations. The BOD values of the water samples varied from 5.07-6.83, the highest value was recorded in station 2. The correlation coefficient tests conducted for all the sampled stations between the microbiological and physicochemical parameters at 95% probability level showed mostly relationships in all the parameters.

DISCUSSION

This research has showed the various effects of human activities and oil pollution on the microbiological and physicochemical qualities of Udu River. There was increase in the most of the parameters studied with little fluctuations. In the study, human activities and oil

pollution effect were observed to play significant role in the influence on the qualities of the parameters investigated. Benka-Coker and Ohimian (1995) noted that increased human activities could bring about increase in organic matter thereby leading to high microbial load. The introduction of waste water, increased amount of organic matter and essential nutrients brings about changes in the microflora (Rheinheimer, 1991). The heterotrophic bacteria and aerobic pathogenic bacteria make up the total bacterial load of river systems. Sayler *et al.* (1975) reported that high microbial population in an aquatic system is a reflection of the input of microorganisms from extraneous sources and availability of growth supporting organic matter. Anson and Ware (1974) noted that high counts of bacterial load reflected the level of water pollution as it gave indication of the amount of organic matter present. These findings correlate with the results of the effect of human activities and oil pollution on the microbiological and physicochemical qualities of Udu River, Warri, Nigeria.

The presence of hydrocarbon utilizing microorganisms in Udu River could be attributed to the regular transportation of oil and its products across the river as well as the presence of hydrocarbons of biogenic origin. The counts for this group of microorganisms in stations 2 and 3 were again higher than those of stations 1 and 4, which may be due to the increased human activities around these stations. Atlas and Bartha (1993) noted that outside the natural input of carbon into aquatic habitats, organic compounds enter the ecosystem through human activities Benka-Coker and Ojior (1995).

The presence of total coliforms, *E. coli*, faecal *Streptococci* and *Clostridium perfringens* in the water samples during sampling period indicated faecal pollution of the water due to human activities. The high densities of faecal indicator organisms observed at all the station in station 2 is indicative of faecal pollution. Akpata and Ekundayo (1978) reported high faecal load with concentration of *E. coli* in the Lagos Lagoon.

The increased microbial load during the dry season may be attributed to surface water runoff, urban runoff, reduced water volume and increased activities.

The physicochemical parameters investigated showed some variation along the sampling stations (Table 3). The temperature range obtained throughout the sampling period was within the WHO range of 10°C to 50°C for surface water (Anonymous, 1971). The conductivity range of the various stations varied considerably between inhabited and uninhabited stations. Station 2 showed the highest value and therefore decreased along the sampling stations, probably due to effect of dilution and removal of soluble salts by

biological utilization. Similar results have been reported by other workers Egborge and Fagade (1999) and Nkwodimah (1985).

The concentrations of nitrate, sulphate and phosphate in all the sampling stations were quite low, except in station 2, which was reported to be due to increased human activities and oil pollution effect. Concentration of dissolved oxygen was highest in station 1 with stations 2 and 3 having lower counts than station 4, this could be due to increased waste disposal at stations 2 and 3 and other form of human activities that may enhance their growth and proliferation, leading to consumption of available oxygen.

Conclusively, it was observed that human activities has impacted negative influence on the quality of water around stations 2 and 3 with station 4 suffering from impact of the upstream stations. The high microbial load as well as high densities of hydrocarbon-utilizing microbes suggested oil pollution of the water and high BOD₅ of the inhabited stations as well as high anions concentrations along the river course and other observations can be used to classify the Udu river as polluted, unfit for direct human consumable needs in accordance with the international standards for classifying surface waters.

REFERENCES

- Anonymous, 1971. International Standard for Drinking Water. 3rd Edn., WHO. Geneva, Switzerland, pp: 15-62.
- Anson, A.E. and G.L. Ware, 1974. Survey of distribution of bacterial population in bristol Channel. J. Applied Bacteriol., 37: 657-661.
- Akpata, V.I. and J.A. Ekundayo, 1978. Faecal pollution of Lagos Lagoon. Nig. J. Sci., 12: 39-53.
- Anonymous, 1989. Standard methods for examination of water and waste. 17th Edn., APHA. Washington, DC. pp: 1325.
- Atlas, R.M. and R. Bartha, 1993. Microbial Ecology: Fundamentals and Applications. 3rd Edn., Benjamin/Cummings Publishing Company. California. pp: 563.
- Barnett, H.L. and B.B. Hunter, 1972. Illustrated genera imperfect fungi. 3rd Edn. Burgess Publishing Company. New York. pp: 241.
- Buchanan, R.E. and N.E. Gibbons, 1974. Bergey's Manual of Determinative Bacteriology. 8th Edn., Williams and Wilken Co., Baltimore., pp: 3100.
- Benka-Coker, M.O. and O.O. Ojior, 1995. Effect of slaughterhouse wastes on the quality of Ikpoba River, Benin City. Biores. Technol., 52: 5-12.

- Benka-Coker, M.O. and E.I. Ohimian, 1995. Investigation of spatial variation in bacterial distribution in surface waters of Ikpoba River. *Nig. J. Microbiol.*, 10: 27-32.
- Egborge, A.B.M., 1982. Hydrobiology of Warri River. Unpublished, pp: 120.
- Egborge, A.B.M. and S.O. Fagade, 1999. Notes on the hydrobiology of the Wikki Warm Spring, Yankari Game Reserve, Nigeria. *Pollut. Arch. Hydrobiol.*, 26: 313-322.
- Gerhardt, P., E.G.R. Murray, A.W. Wood and R.N. Krieg, 1994. *Methods for General and Molecular Bacteriology*. ASM. Press. Washington, DC. pp: 791.
- Miller, J.E., 2001. Water pollutants and public health. *Microbiology*. 3rd Edn., John Wiley and Sons. New York, pp: 411.
- Nkwodimah, C.N., 1985. Bacteriological and physicochemical quality of Ojirami Dam. M.Sc. Thesis. University of Benin, Benin City, Nigeria, pp: 32.
- Prescott, L.M., J.P. Harley and D.A. Klein, 2000. *Microbiology*. 4th Edn., McGraw Hill Companies Inc. USA, pp: 963pp
- Rheinheimer, G., 1991. *Aquatic Microbiology*. 4th Edn. John Wiley and Sons. New York, pp: 363.
- Saylor, G.S., J.O. Nelson, A. Justice and R.R. Colwell, 1975. Distribution and significance of faecal indicator organisms in the upper Chasepark Bay. *Applied Indust. Microbiol.*, 30: 625-638.
- Thurman, R., B. Foulkner, D. Veal, G. Gramer and M. Meiklejohn, 1998. Water quality in rural Australia. *J. Applied Microbiol.*, 84: 627-632.
- Udo, K.R., 1975. *Geographical Region of Nigeria*. Heinemann Educational Books, London, pp: 212.