



Journal of Biological Sciences

ISSN 1727-3048

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Effect of Salt, Nickel Chloride Supplementation on the Growth of Silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae)

Md. Rezuanul Islam, ¹Md. Abdul Ohayed Ali, ²Dipak Kumar Paul, ¹Shaheen Sultana,
Nilufa Akhter Banu and Md. Rafiqul Islam

Department of Biotechnology, Islamic University, Kushtia, Bangladesh

¹Department of Zoology, Rajshahi University, Rajshahi, Bangladesh

²Department of Applied Nutrition and Food Science, Islamic University, Kushtia, Bangladesh

Abstract: The present investigation deals with the effect of nickel chloride supplementation on the growth of *Bombyx mori* L. (Lepidoptera: Bombycidae). The weights of mature *Bombyx mori* L. larvae were 1522.10±78.97, 1579.67±61.29, 1685.13±61.77, 1781.60±66.62, 1801.60±60.69 and 1426.07±89.48 mg for nickel chloride at 0 (Control), 100, 200, 400, 800 and 1600 ppm, respectively. The weights of male and female pupae were 648.03±37.97 and 763.57±55.44, 667.50±51.10 and 774.23±54.33, 682.60±61.78 and 811.47±61.13, 712.23±42.46 and 841.33±63.04, 746.77±45.73 and 871.30±50.47, 618.40±30.61 and 713.70±63.16 mg for nickel chloride for above applied doses. The weights of male and female adults were 300.67±70.07 and 546.27±61.67, 317.63±42.98 and 559.50±76.87, 331.90±48.93 and 571.50±80.46, 352.00±55.14 and 597.67±56.53, 373.17±46.53 and 645.23±55.34, 283.40±43.89 and 514.27±75.83 mgs for nickel chloride for above applied doses. The weights of male and female cocoons were 752.07±31.70 and 87.17±55.50, 769.23±49.84 and 889.10±50.40, 795.53±61.69 and 937.10±59.70, 841.33±42.25 and 970.53±62.33, 856.93±48.50 and 979.63±44.09, 717.27±32.13 and 822.13±64.96 mg for nickel chloride at 0 (Control), 100, 200, 400, 800 and 1600 ppm, respectively. The results were statistically significant ($P < 0.001$) among the concentrations. From the above results it may be concluded that nickel chloride can be used at low concentrations for enhancing the economic character of silkworm, *Bombyx mori* L.

Key words: Nickel chloride, *Bombyx mori* L., silkworm

INTRODUCTION

The mulberry silkworm *Bombyx mori* L. is a beneficial insect producing the finest natural silk. Silk is called the “Queen of textiles”. Sericulture is essentially a village based and welfare oriented industry that plays an important role in our national economy.

There are about 1000 varieties of silkworm^[1], among them, bivoltine and multivoltine races are used for rearing in our country. *Bombyx mori* L. is a domesticated variety, which is believed to have been derived from the original mandarina silkworm *Bombyx Mandarina* Moore^[2].

Nutrition is an important growth regulatory factor in silkworms like in any other organisms, the silk worm and the mulberry tree have a great partnership because the silkworm can not thrive without the leaves of mulberry plants and the quality of leaves of mulberry plants and the quality of leaves influence greatly the biology of these worms^[3,4]

Bombyx mori L. is an oligophagous insect that feeds mainly on mulberry leaves. Significant advances in the

research of silkworm nutrition began with development of artificial diets^[5]. Advances in silkworm nutrition and in the development and use of artificial diets will be of great benefit to the silk industry. The effect of artificial diets with different nutrition on better production of cocoon crops and silk were investigated by many workers like Fukuda and Higuchi^[6], Yokoyama^[7] and Hamamura^[8], additions of extranutrients to the mulberry leaf in the silk worm enhance their growth and cocoon yield^[9].

Salts may constitute a limiting factor in the growth of insects upon some diets^[10]. The salt significantly enhanced the growth of the developmental stages, Decreased the developmental period, increased the cocoon characters and increased the reproductive potential of the worms at 0.25 and 0.50%.

Dietary supplementation of copper sulphate, nickel chloride and potassium iodide increased the economic parameters of the silkworm^[11] The present investigation reports the effects of mulberry leaves enriched with nickel chloride, potassium iodide and their combinations on the growth and development of *Bombyx mori* L.

MATERIALS AND METHODS

The insect used for the present study was the mulberry silkworm, *Bombyx mori* L. is collected from the germ plasm bank of the Bangladesh Sericulture research and training Institute (BSRTI), Rajshahi, Bangladesh. Standard rearing techniques were followed.

Effect of salt supplementation on the growth of *Bombyx mori* L.: The growth of various stages of the silkworm, *Bombyx mori* L. is of paramount importance because a successful cocoon crop in sericulture depends mostly on a healthy larval growth. It has been well established that *Bombyx mori* L. requires certain essential sugars, protein, salts, amino acids and vitamins for its normal growth, survival and also for the growth of its silk glands and higher production of good quality silk^[12].

Effect of salt on the growth of *Bombyx mori* L. larvae: To study the effect of salt on the growth of *Bombyx mori* L. larvae the concentration of the salt were prepared by mixing appropriate quantities of nickel chloride [NiCl₂.6H₂O] in distilled water. The concentrations were 0 (Control), 100, 200, 400, 800 and 1600 ppm. Control batches were raised on leaves treated with distilled water only. These were three replication per concentration each with 100 larvae. Treated leaves were chopped and supplied to the worms after hatching up to the third moult four times a day. For the fourth and fifth instar larvae whole leaves were supplied. The growth of larvae was determined after attaining maturity, i.e., one day before spinning ten larvae were taken randomly from each rearing tray and then weighted individually on an electronic balance on milligrams and calculated through statistical analysis.

Effect of salt on the growth of *Bombyx mori* L. pupae: After attaining maturity silk worm larvae formed cocoon within which they were transformed into pupae. To determine the weight of pupae from various concentration of nickel chloride, mature larvae from different trays were transferred to bamboo made mountages (1.8x1.2 m). After the formation of cocoons they were harvested and cut very carefully at one end obliquely with a sharp blade to observe pupal characters. In all the cases 10 males and 10 females were considered and the experiment was repeated thrice. The coefficient of variation (CV) and t-values were calculated by using standard statistical formula.

Effect of salt on the growth of *Bombyx mori* L. adults: After nine to twelve days of spinning the adults emerged from cocoons by rupturing the shells. Freshly emerged

adults were individually weighted on an electronic balance in milligrams and calculated the values.

Effect of salt on the *bombyx mori* L. Cocoon characters: Cocoons were harvested after four to five days of spinning and weighted individually on electronic balance on milligrams and calculated the data.

RESULTS AND DISCUSSION

The mature larval weights are shown in Table 1 and it was observed that the supplementation of mulberry leaves with nickel chloride produced the highest larval weight at 800 ppm and lowest at 1600 ppm.

The weight of male and female pupae (Table 2) *Bombyx mori* L. resulting from various concentration was observed of nickel chloride, the highest and lowest pupal weights were observed at 800 and 1600 ppm, respectively.

The data obtained in the present experiment, for adult, weights of *Bombyx mori* L. resulting from various treatments showed that heaviest and lightest moths were obtained at 800 and 1600 ppm nickel chloride respectively. It was reported that Cobalt exerts favorable effect on the growth of silkworm (Table 3)^[13].

Table 1: Indicating the effect of nickel chloride on the mature larval weight (mg) of *Bombyx mori* L

Concentration (ppm.)	Mean ± SD	CV (%)	t-value	Level of significance
0 (Control)	1522.10±78.92	5.18	—	—
100	1579.67±61.29	3.88	-3.16	P<0.01
200	1685.13±61.77	3.66	-8.91	P<0.001
400	1781.60±66.62	3.74	-13.76	P<0.001
800	1801.60±60.69	3.37	-15.38	P<0.001
1600	1426.07±89.48	6.27	4.41	P<0.001
F-Ratio	145.15			
CD at 0.05	108.52			

Table 2: Indicating the effect of nickel chloride on the mature pupal weight (mg) of *Bombyx mori* L

Concentration (ppm.)	Mean±SD (Male and female respectively)	CV (%)	t-value	Level of significance
0 (Control)	648.03±37.97 (763.57±55.44)	5.85 7.26	—	—
100	667.50±51.10 (774.23±54.33)	7.65 7.01	-1.68 (-0.77)	NS NS
200	682.60±61.78 811.47±61.13	9.05 7.53	-2.61 -3.18	P<0.05 P<0.01
400	712.23±42.46 841.33±63.04	5.96 7.49	-6.17 -5.07	P<0.001 P<0.001
800	746.77±45.73 871.30±50.47	6.12 5.79	-9.19 -7.87	P<0.001 P<0.001
1600	618.40±30.16 713.70±63.16	4.94 8.84	3.38 3.27	P<0.01 P<0.01
F-Ratio	20.33			
	12.78			
CD at 0.05	73.64			
	89.83			

Note: NS= Non significant

Table 3: Indicating the effect of nickel chloride on the mature adult weight (mg) of *Bombyx mori* L.

Concentration (ppm.)	Mean±SD (Male and female respectively)	CV (%)	t-value	Level of significance
0 (Control)	300.67±70.07	23.30	—	—
	546.27±61.67	11.29		
100	317.63±42.98	13.53	-1.13	NS
	559.50±76.87	13.74	-0.74	NS
200	331.90±48.93	14.74	-2.00	NS
	571.50±80.46	9.45	1.36	NS
400	352.00±55.14	15.66	-3.15	P<0.01
	597.67±56.53	9.45	-3.37	P<0.01
800	373.17±46.53	12.47	-4.72	P<0.001
	645.23±55.34	8.57	-6.54	P<0.001
1600	283.40±43.89	15.48	1.14	NS
	514.27±75.83	14.74	1.79	NS
F-Ratio	10.19			
	6.28			
CD at 0.05	83.37			
	109.03			

Note: NS= Non significant

Table 4: Indicating the effect of nickel chloride on the cocoon weight (mg) of *Bombyx mori* L.

Concentration (ppm.)	Mean±SD (Male and female respectively)	CV (%)	t-value	Level of significance
0 (Control)	752.07±31.70	4.21	—	—
	878.17±55.50	6.32		
100	769.23±49.84	6.48	-1.59	NS
	889.10±50.40	5.67	-0.80	NS
200	795.53±61.69	7.75	-3.43	P<0.01
	937.10±59.70	6.37	-3.96	P<0.001
400	841.33±42.25	5.02	-9.26	P<0.001
	970.53±62.33	6.36	-6.66	P<0.001
800	856.93±48.50	5.66	-9.91	P<0.001
	979.69±44.09	4.54	-7.14	P<0.001
1600	717.27±32.17	4.45	4.22	P<0.001
	822.13±64.96	7.90	3.59	P<0.001
F-Ratio	42.53			
	29.01			
CD at 0.05	71.87			
	98.65			

Note: NS= Non significant

The weights of male and female cocoons showed that nickel chloride at 100, 200, 400 and 800 ppm gradually increased the cocoon weight but at the highest concentration, i.e, 1600 ppm weight was significantly reduced (Table 4).

Nutrition is an important growth regulating factor in silkworm, like in any other organisms. Various minerals take part in regulation of osmotic pressure of intra and extra cellular fluids in the maintenance of an ionic balance suitable for the activity of living cells and as co-factors in some enzyme systems and as integral part of other systems. The known essential trace elements for insects growth include iron, nickel, copper, manganese, zinc and iodine^[14].

Our present investigation, it was found that nickel chloride significantly increased the growth of larvae, pupae and adults but higher salt concentrations produced deleterious effects on these parameters. The present

findings, the larvae pupal and adult growth in *Bombyx mori* L. were enhanced at the concentration of 800 ppm and the economic characters (cocoon characters) of the mulberry were enhanced at 800 ppm of nickel chloride.

REFERENCES

1. Barnet, L., 1963. The wonders of life and earth. Parentic and Hall. Landon, pp: 48.
2. Krishnaswami, S., M.N. Narasimahanna, S.K. Suryanarayn and S. Kumararaj, 1973. Sericulture Manual J. Silkworm Rearing, FAO, Rome., pp: 131.
3. Frassie, R. and Arnoux, 1954. Biometrical characters of the cocoon in *Bombyx mori* L. and their variation under the influence of food. J. Silkworm, 6: 41-62.
4. Arai, N. and T. Ito, 1963. Food value of mulberry leaves for the silkworm, *Bombyx mori* L. determined by means of artificial diets. II. Comparison between soft and hard leaves. Bull. Seric. Expt. Sta., Tokyo, 18: 1-16.
5. Ito, T., 1978. Silkworm nutrition. In: The silkworm: An Important Laboratory Tool, (Ed. Y. Tazima), pp: 121-57. Tokyo: Kodansha Ltd., pp: 307.
6. Fukuda, T. and Higuchi, 1963. Artificial food for oak Silkworm rearing. Agric. Biochem., 27: 99-102.
7. Yokoyama, Y., 1964. Alimentation Artificielle pour to ver. A soie. J. Silkworm, 15/16: 1-13.
8. Hamamura, Y., 1964. On the feeding mechanism and prepared food of silkworm, *Bombyx mori* L. J. Silkworm, 14: 49-56.
9. Sharda, K., T. Leena, M.B. Shroof, K. Shtamala, S. Ivenore and J.V. Bhat, 1956. Large scale experiments on the effect of chloromycitine and glycine in the nutrition of silkworm, *Bombyx mori* L. J. Indian Inst. Seri., 38.
10. Wiggleswarth, V.B., 1972. The principles of insect physiology. Chapman and Hall. London, pp: 827.
11. Magadam, S.B., 1987. Studies on the effect of some chemicals and hormones on the gonads and the life cycle of the Silkworm, *Bombyx mori* L. Ph. D. Thesis, Karnataka University, Dharwad. India, pp: 1-30.
12. Sengupta, K., B.D. Singh and C. Mustafij, 1972. Nutrition of Silkworm. *Bombyx mori* L. I. Studies on the enrichment of mulberry leaf with various sugars, proteins, aminoacids and vitamin for vigorous growth of the worm and increased cocoon crop production. Indian J. Sci., 11: 11-27.
13. Takahashi, J., 1956. Vitamin B-12 in Silkworm II. Nippon. Nogei. Kagaku Kaishi, 29: 711-715.
14. Trager, W., 1953. Nutrition, In: Insect physiology (Ed. K.D. Roeder), John wiley and sons. New York, pp: 250-256.