



Journal of Biological Sciences

ISSN 1727-3048

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Promotive Effects of 5-Aminolevulinic Acid (5-ALA) on Fruit Yield and Quality of Date Palm cv., Khalas

¹A.A. Al-Khateeb, ¹S.A. Al-Khateeb, ²R. Okawara and ³I.A. Al-Abdoulhady
¹Date Palm Research Center, King Faisal University, P.O. Box 420, Al-Hassa, 31982
²King AbdulAziz for Science and Technology (KACST)
³Date Palm Research Center, Ministry of Agriculture, Al-Hufuf, Saudi Arabia

Abstract: The effects of spraying 5-ALA on date palm in relation to chlorophyll biosynthesis and fruits yield and quality during 1998/1999 seasons had been investigated. Fruits of Khalas cv. were sprayed with 5-ALA aqueous solutions of 0, 50 and 100 ppm two weeks after fruit setting. Spraying was applied biweekly for six weeks duration. Fruits chlorophyll content in Khalal stage was significantly increased with 5-ALA treatments. Fruit weight, fruit volume and fruit flesh percentage of rutab stage were significantly increased with 100 ppm 5-ALA. On tamer stage, fruit volume was significantly increased with 5-ALA treatment, while fruit weight was not. Total and reducing sugars were significantly increased with 5-ALA treatment in rutab stage, but not in tamer stage.

Key words: 5-Aminolevulinic acid, date palm, CV Khalas, chlorophyll, fruit quality, yield

INTRODUCTION

Date Palm (*Phoenix dactylifera*, L.) is a major fruit tree of Saudi Arabia with a total production of about 830000 tons date fruits FAO (2004). Cultivation of cv. Khalas date palm has tremendously increased in Saudi Arabia in recent years. This cultivar is well known for its high fruit quality. However, with the recent increase in its cultivation, several fruit quality problems have been surfaced. Fruit size is considered one of the major factors that determine the income of the producers. Chemical spraying (Ethrel, GA₃, Ethepon, Naphthalene Acetic Acid) improved date palm fruit size and quality (Rouhani and Bassiri, 1977; El-Hamady *et al.*, 1983, 1992; Al-Khateeb *et al.*, 1993; Hussein *et al.*, 1996; Moustafa and Seif, 1996; Moustafa *et al.*, 1996). However, chemical treatment, particularly GA₃ delayed fruit ripening and decreased total soluble solid and total sugar (Hussein *et al.*, 1996; Moustafa and Seif, 1996).

5-Amino levulinic acid (5-ALA) is a precursor of tetrapyrrole compounds such as chlorophyll, phycobilin, heme and vitamin B12 which are exist in plants. Foliar spraying of 5-ALA at low concentrations improved the growth and yield by 10-60% of crops such as radish, kidney beans, barley, potatoes, garlic, rice and corn (Hotta *et al.*, 1997a, b; Bingshan *et al.*, 1998; Yongin *et al.*, 2003; Al-Thabet, 2006; Al-Khateeb, 2006). However, in plants treated with 5-ALA at relatively high concentration, more than 10 mM, herbicidal properties of

5-ALA were well documented (Hotta *et al.*, 1997a). Effects of 5-ALA in plants have been reported in relation to chlorophyll biosynthesis, photosynthesis activity and suppression of respiration (Hotta *et al.*, 1997a).

Treating fruit of date at khalal stage (green stage) is expected to show some of the previously mentioned effects. If this assumption is true, the contribution of the treated fruits to photosynthetic activity could be higher than untreated ones. Therefore, accumulation of assimilates due to the increased activity of photosynthesis of treated fruits might positively affect the fruit size and chemical properties. Therefore the present study was planned to study the effects of spraying 5-ALA on the fruits of date palm in relation to chlorophyll biosynthesis and fruits yield and quality.

MATERIALS AND METHODS

The study was carried out in 1998/1999 in Al-Hassa. Twelve uniform 25 years old vigorous palm trees cv. Khalas were selected. They were subjected to the normal agricultural practices and thinned to eight bunches/tree. Pollination was conducted using the same male parent for all experimental palm trees. Fruits were sprayed with 5-ALA aqueous solutions of 0, 50 and 100 ppm containing Tween 20 (0.1%). The spraying treatments were started two weeks after fruit setting with approximately 240 mL/tree of aqueous solution (approximately, 30 mL/bunch). Spraying was applied

biweekly in the early morning for a duration of six weeks. Samples from each tree (replicate) were randomly collected at Khalal, rutab and tamer stages. A completely randomized block design with four replicates per treatment was used.

Chlorophyll was determined in khalal stage after six weeks of spraying application. Fruits skin (the grin coat only) was taken by a cork borer and then homogenized in cold (4°C) 80% v/v acetone in water. The homogenate was kept in the dark and centrifuged for 3.0 min to remove the fruit debris. The absorbance of the extract at 647 and 664 nm was taken using a spectrophotometer. For the accurate determination of chlorophyll a, b and the total, the extinction coefficients of Graan and Ort (1984) were used. Physical properties of rutab (two weeks after fruits become yellow in color) and tamer (four weeks after fruits become brown in color) were obtained, i.e., fruit length and diameter (cm), fruit weight (g), fruit size (cm³), fruit flesh (%) and fruit seed (%). Chemical properties including moisture (%), ash content, total sugars and reducing sugars were measured according to AOAC (1984). Collected data were subjected to analysis of variance (ANOVA). SAS computer package (SAS Institute, 2001) has been used. Throughout, p = 0.05 was used to define statistical significance.

RESULTS AND DISCUSSION

Total chlorophyll and chlorophyll a and b contents at khalal stage were significantly increased with ALA treatments (Fig. 1). There was no marked increase in chlorophyll a with increasing 5-ALA concentration from 50-100 ppm. However, 50 ppm 5-ALA treatment showed significantly the highest total chlorophyll and chlorophyll b concentrations. Since a slight drop in all chlorophyll types was noticed with the 100 ppm 5-ALA application, the 50 ppm 5-ALA concentration might have represented a physiological threshold level beyond which chlorophyll deteriorates. The increase in chlorophyll content has been reported in horse radish treated with low concentration of 5-ALA less than 100 ppm, while respiration was suppressed in certain crops with higher levels of 5-ALA, more than 100 ppm (Hotta *et al.*, 1997a).

Treatment of 5-ALA had no significant effects on rutab fruit dimensions. Fruit weight, fruit volume and fruit flesh% were significantly increased with increasing concentrations of 5-ALA, while the seed fruit% was significantly reduced. The significant increase in rutab fruit weight with 5-ALA treatments may be attributed to the increase in flesh fraction under the same treatments. The fruit volume showed slight positive increasing with 5-ALA spraying, while both fruit diameter and length were not. Under this situation, it is not possible to establish a




Fig. 1: Chlorophyll content in Khalal stage of cv. Khalas as affected by ALA treatments. Bars = LSD (5%)

clear positive relation between fruit volume and the two dimensions. However, we can continuously assume that other unobserved factors may have affected the volume parameter in addition to the slight non significant increases in fruit diameter and length. The increase in fruit weight and volume was approximately similar, i.e., 30% more than control. Fruit dimensions of tamer were not significantly affected by 5-ALA treatments. Fruit volume was significantly increased with 5-ALA treatment, while fruit weight was not. Seed and fruit flesh% were not significantly affected by 5-ALA treatment (Table 1). It is quite possible that fruits at this stage may have reached a stable physiological maturation that can not be changed by 5-ALA or any other hormonal treatments. Moreover, the 5-ALA effects are much possibly pronounced at the rutab stage of the fruit. Hussein *et al.* (1996) reported that GA3 treatment significantly increased fruit weight, volume, length and thickness, but there was no definite trend for seed weight in Zaghlolo date cv. Similar effects had been reported by Moustafa and Seif (1996). The proper time of chemical application during fruit or flower development to obtain certain desirable characters of fruits have been well investigated (Rouhani and Bassiri, 1977; El-Hammady *et al.*, 1992; Al-Khateeb *et al.*, 1993). Total and reducing sugars were significantly increased in rutab stage with the application of 5-ALA, but there was no significant difference in non reducing sugar content (Table 2). As was the case in chlorophyll, 50 ppm 5-ALA yielded significantly the highest total and reducing sugars. Liangju *et al.* (2004) found that spraying 5-ALA increased the total soluble solid content of apple fruits. This possibly indicates an active photosynthesis rate and/or reducing respiration rate with this concentration. Ash content was significantly reduced with increasing 5-ALA concentration. Moisture content was significantly

Table 1: Fruit and seed physical properties of Khalas cv. Rutab and Tamer stages as affected by ALA treatments

ALA concentration (ppm)	Weight (g)	Volume (cm ³)	Flesh fruit (%)	Seed fruit (%)	Diameter (cm)	Length (cm)
Rutab						
0	8.81	8.5	88.2	11.8	2.13	3.18
50	10.10	9.9	89.8	10.2	2.19	3.33
100	11.49	11.1	90.1	9.9	2.24	3.42
F-test	*	*	*	*	NS	NS
LSD (5%)	2.27	2.58	1.7	1.7	-	-
Tamer						
0	5.88	5.9	87.7	12.3	1.91	3.1
50	6.33	6.4	88.7	11.1	1.91	3.1
100	6.75	7.2	89.2	10.8	1.93	3.2
F-test	NS	*	NS	NS	NS	NS
LSD (5%)	-	0.9	-	-	-	-

* = Significant at 0.05 level of probability, NS = Not Significant

Table 2: Chemical properties of Khalas Rutab and Tamer stages as affected by ALA treatments

ALA concentration (ppm)	Moisture (%)	Ash (%)	Reducing sugars (%)	Non red. sugars (%)	Total sugars (%)
Rutab					
0	64.0	0.98	24.3	1.6	25.9
50	58.3	0.93	30.6	2.0	32.6
100	62.8	0.82	26.9	1.5	28.4
F-test	*	*	*	NS	*
LSD (5%)	5.2	0.09	4.1	-	4.2
Tamer					
0	6.9	1.78	70.1	4.3	74.4
50	7.7	1.57	71.6	4.2	75.8
100	8.3	1.58	70.5	4.2	74.7
F-test	NS	*	*	NS	NS
LSD (5%)	-	0.17	1.3	-	-

* = Significant at 0.05 level of probability, NS = Not Significant

lower with 50 ppm 5-ALA, but there was no significant difference between control and 100 ppm 5-ALA. The reduction on moisture content with 5-ALA treatment indicates an increase in dry matter of fruits which raise the possibility of higher photosynthetic activity of fruits at khalal stage. The improved photosynthesis efficiency of khalal stage might possibly led to the accumulation of assimilates which might explain the increase in dry matter content of fruits. Total and non-reducing sugars in tamer stage were not much altered with 5-ALA, but there was an increase in reducing sugars with 5-ALA treatment and this increase was significant and best with 50 ppm 5-ALA. Similar to rutab stage, ash content significantly decreased with 5-ALA treatment. Moisture content was not significantly changed with 5-ALA treatment. However, there was a clear trend of an increase in moisture content. This increase in moisture content may explain the significant increase in fruit volume of tamer. Although, there was no significant effect of 5-ALA on fruit weight, a 20% increase in fruit weight was obtained compared to 22% in fruit volume. It is worth mentioning that an increase in fruit volume and weight of tamer might be




Fig. 2: Tamer yield (kg/tree) in cv. Khalas as affected by ALA treatments. Bars = LSD (5%)

obtained due to the flesh fraction. Mustafa and Seif (1996) and Hussein *et al.* (1996) reported that total sugars% has been reduced with GA3 treatment. However, Hussein *et al.* (1996) reported an increase in total and reducing sugars in Zaghlol date treated with cycocel.

Despite the consistent increasing trend, fruit yield (kg/tree) was not significantly affected (Fig. 2). Yield is a combined factor of fruit number, size, weight and other related variables (Soliman and El-Kosary, 2002; Mushtaq *et al.*, 2004). In this study, 5-ALA has improved fruit weight and volume at rutab stage. It is quite probable that increasing of fruit size or weight reduces fruit number as has often been shown in certain fruit crops (El-Hamady *et al.*, 1992; Soliman and ElKosary, 2002; Mushtaq *et al.*, 2004). Improvement of fruit size is becoming detrimental in marketing of date fruits and highly preferred by consumers.

REFERENCES

- Al-Khateeb, A.A., A.S. Elgamdi and O. Eltahir, 1993. Effect of thinning stage on fruit quantity and quality of date palm cv. Khalas Proc. 3rd Symp. On date palm, King Faisal University, Kingdom of Saudi Arabia.
- Al-Khateeb, S.A., 2006. Promotive effect of 5-AmioLevulinic Acid on growth, yield and gas exchange capacity of barley (*Horidium vulgare* L.) grown under drought regimes. J. King Saud Uni. (Agriculture Sciences) (In Press).
- Al-Thabet, S.S., 2006. Promotive effect of 5-AmioLevulinic Acid on growth and yield of wheat grown under drought condition. J. Agron., 5: 45-49.
- AOAC., 1984. Official methods of analysis of the Association Official Analysis Chemists, 13th Edn., Washington, DC., USA.

- Bingshan, L., Y. Hotta, Z. Qu Yinglan, T. Jinsong, Y. Tanaka, Takeuchi and M. Konnai, 1998. Effects of 5-Aminolevulinic acid on the growth and ripening of wheat. *J. Pesticide Sci.*, 23: 300-303.
- El-Hamady, M.M., A.S. Khalifa and A.M. Al-Hamady, 1983. Fruit thinning in date palm with ethephon. Proc. 1st Symp. On date palm. Saudi Arabia, pp: 245-295.
- El-Hamady, M.M., M.A. Al-Maghrabi and M.A. Basha, 1992. Effect of ethephon treatment on fruit thinning and quality of seleg and meneify date palm cultivars. *Annal of Agric. Sci., Cairo*, 37: 531-538.
- FAO, 2004. FAOSTAT Database 2004, (<http://apps.fao.org>).
- Graan, T. and D.R. Ort, 1984. Quantitation of the rapid electron donors to P 700, the functional plastoquinone pool and the ratio of the photosystems in spinach chloroplast. *J. Biol. Chem.*, 259: 14003-14010.
- Hotta, Y., T. Tanaka, H. Takaoka, Y. Takeuchi and M. Konnai, 1997a. Promotive effects of 5-aminolevulinic acid on the yield of several crops. *Plant Growth Regul.*, 22: 109-114.
- Hotta, Y., T. Tanaka, H. Takaoka, Y. Takeuchi and M. Konnai, 1997b. New physiological effects of 5-aminolevulinic acid in plants: The increase of photosynthesis, chlorophyll content and plant growth. *Biosci. Biotech. Biochem.*, 61: 2025-2028.
- Hussein, M.A., H.M. Mahmoud, K.I.A. Amen and M. Mustafa, 1996. Changes in the physical and chemical characteristics of Zaghloul dates during development and maturity as affected by GA3 and CCC under Assuit Governorate conditions. Proceedings of the 3rd Symposium on the Date Palm in Saudi Arabia. January, 17-20: 389- 404.
- Liangju, W., W. Zhonghua, L. Zhiqiang., L. Hui, L. Weiqin., C. Zongyuan, Y. Pu. and S. Dunqian, 2004. Effects of 5-Amiolevulinic Acid on enhancing apple fruit coloration. *J. Fruit Sci.*, 21: 512-515.
- Moustafa, A.A., S.A. Seif and A.I. Abou-El-Azayem, 1996. Date fruit response to naphthalene acetic acid. Proceedings of the 3rd Symposium on the Date Palm in Saudi Arabia. January, 17-20 : 369-378.
- Moustafa, A.A. and S.A. Seif, 1996. Effect of ethrel and GA treatments on yield and fruit quality of seewy date palms, grown in El-Fayoum Governorate. Proceedings of the 3rd Symposium on the Date Palm in Saudi Arabia. January, 17-20: 379- 388.
- Mushtaq, A., M.A. Bashir, N.A. Chaudhry, M.A. Shakir and M. Khan, 2004. Performance of date varieties under Bahawalpur conditions. *Sarhad J. Agric.*, 20: 39-42.
- Rouhani, I. and A. Bassiri, 1977. Effect of ethephon on ripening and physiology of date fruits at different stage of maturity. *J. Hortic. Sci.*, 52: 289-297.
- SAS Institue, 2001. SAS for Windows, SAS user's guide: Statistics. Version 8.0 e. SAS Inst., Inc., Cary, North Carolina.
- Soliman, S.S. and S. El-Kosary, 2002. Effect of different hand pollination methods on fruit retained, yield and fruit quality of some Egyptian dry date palm cultivars. *Egyp. J. Hprtic.*, 29: 281-300.
- Yongin, K., L.G. Seob, C. Sang, H. Tayeak and G Jaock, 2003. Effects of 5-Amiolevulinic Acid on growth and inhibition of various plant species. *Kor. J. Crop. Sci.*, 48: 127-133.