



# Journal of Biological Sciences

ISSN 1727-3048

**science**  
alert

**ANSI***net*  
an open access publisher  
<http://ansinet.com>

## Cold Pretreatment-Induced Changes in Antioxidant Enzyme Activities and Relative Water Content and Soluble Sugars in Shoots and Roots of Soybean Seedlings

Leila Zeinali Yadegari, Reza Heidari and Jirair Carapetian  
Department of Biology, Faculty of Science, Urmia University, Iran

**Abstract:** Physiological responses to chilling, including antioxidative enzyme activity, Relative Water Content (RWC) and soluble sugar contents were investigated in soybean to identify mechanisms of chilling tolerance. Plants were exposed to 15°C (cold-acclimated) or 25°C (non-acclimated) for 24 h, under 250  $\mu\text{mol m}^{-2} \text{sec}^{-1}$  Photosynthetically Active Radiations (PAR). Then all plants were exposed to 4°C (chilling temperature) for 24 h and allowed to recover at 25°C for 24 h. We analyzed the activity of ascorbate peroxidase (APX), catalase (CAT) and guaiacol peroxidase (GPX) and soluble sugar content and RWC in both shoots and roots of soybean seedlings. It revealed that the activity of APX and CAT and GPX induced in leaves and roots. Increased activity in roots is important for cold tolerance as compared to shoots. The amount of RWC decreased in both roots and shoots, but soluble sugar content increased, especially in shoots as compared to control plants. Chilling sensitive soybean plants can be made tolerant to cold by cold-acclimation.

**Key words:** Acclimation, antioxidant, ascorbate peroxidase, catalase, chilling, guaiacol peroxidase, RWC, soluble sugar

### INTRODUCTION

Food crops of tropical and subtropical origins such as soybean (*Glycine max*), corn (*Zea mays* L.) and tomato (*Lycopersicon esculentum* Mill) are cultivated in areas where temperatures fall below the optimum required for their normal growth and development. It is now known that exposure of chilling-sensitive plants, such as maize and tomato, to temperatures that are slightly above chilling, reduces chilling injury (Scebba *et al.*, 1999; Venema *et al.*, 2000). Various mechanisms have been suggested to account for chilling injury or tolerance in plants (Barsa, 2001). There is increasing evidence that chilling causes elevated levels of Active Oxygen Species (AOS), which contribute significantly to chilling damage (Wise and Naylor, 1987a). AOS such as superoxide ( $\text{O}_2^-$ ), hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), hydroxyl radicals ( $\text{OH}^\bullet$ ) and singlet oxygen ( $^1\text{O}_2$ ), are present in plants at various levels at 25°C as a result of normal aerobic metabolism. Plants have evolved antioxidant systems to protect cellular membranes and organelles from damaging effects of AOS (Foyer *et al.*, 1991). Antioxidant enzymes, such as catalase (CAT) and various peroxidases such as guaiacol peroxidase (POX) and ascorbate peroxidase (APX) can react with and neutralize, the activity of AOS (Foyer *et al.*, 1991; Lee and Lee, 2000; Oidaira *et al.*, 2000; Scandalios, 1993). Beside these enzymes, antioxidant compounds such as ascorbate, glutation,  $\beta$ -carotene and  $\alpha$ -tocopherol

also play important roles in the removal of toxic oxygen compounds (Hodges *et al.*, 1996; Wise and Naylor, 1987b). Cold-acclimation increases tolerance to AOS in plants with an increase in antioxidant enzymes (Scebba *et al.*, 1999). In chilling sensitive plants, the ability to defend against oxidative damage is inhibited by the reduction expression of antioxidants such as ascorbate, glutation and  $\alpha$ -tocopherol (Wise and Naylor, 1987b), CAT (Fadzillah *et al.*, 1996) and SOD (Michalski and Kanjuga, 1982). Chilling tolerance improved when GSH peroxidase and CAT levels were enhanced (Upadhyaya *et al.*, 1989). Thus, it is important to determine the activity of various antioxidants during acclimation and chilling to assess their contribution to chilling tolerance. Sugars appear essential in plant cold-acclimation, as shown for example by the inability of an *Arabidopsis* sucrose synthase mutant to cold-acclimation (Welling and Palva, 2006) or the requirement for light in low-nonfreezing temperature-induced cold-acclimation connected to sugar accumulation (Wanner and Junttila, 1999). Precise function of sugars is not known, but their high abundance in cold acclimated plants suggests a role in osmoregulation and less abundant sugars might also have a role in cryoprotection or as signaling molecules (Welling and Palva, 2006). We aimed to determine whether AOS-scavenging enzymes and soluble sugar and relative water contents play a role in soybean tolerance to chilling stress.

## MATERIALS AND METHODS

Seeds of soybean (*Glycine max*) were purchased from oilseeds center, Ardabil and were soaked in water for 6 h at 25°C, then were germinated in Petri dishes on two layers of filter paper for 48 h at 25°C in an incubator. Subsequently the soybean seedlings were transferred to pots containing washed sand (4 seedlings per pot) and were watered with half-strength Hoagland nutrient solution. The plants were grown at 27/25°C (day/night) temperature, 70% relative humidity, with a 16/8 h day/night photoperiod under 250  $\mu\text{mol m}^{-2} \text{sec}^{-1}$  photosynthetic photon flux density. Seedlings at the three-leaf stage, were placed at 15°C (cold-acclimated) or 25°C (non-acclimated) for 24 h. The acclimated and non-acclimated seedlings were then exposed to chilling at 4°C for 24 h and allowed to recover for 24 h at 25°C. Harvesting was done at the same time each day to avoid complications from diurnal fluctuations in biochemical processes. Experiments were conducted from May to July in 2007 at Biochemistry Lab, Department of Biology, Faculty of Science, Urmia University, Iran. Statistical analyses were performed using SAS software. Measurements were mean of 3 replicates  $\pm$  SE.

**Enzyme extraction and assay:** Samples were prepared for CAT, APX and GPX analyses by the method described by Chang and Kao (1998). The amount of 0.5 g of frozen leaf and root materials from all treatments were harvested and homogenized with a mortar and pestle in 6 mL of extraction solution [0.05 M tris-HCl buffer (pH = 7.2), 3 mM  $\text{MgCl}_2$ , 1 mM EDTA]. Then the solution was centrifuged for 20 min at 5000 rpm. The supernatant was used for monitoring antioxidative enzymes mentioned above. Catalase activity was determined by monitoring the disappearance of  $\text{H}_2\text{O}_2$  at 240 nm using the method of Chance and Maehly (1959). The reaction mixture contained 50 mM potassium phosphate buffer (pH = 7.4), 0.1 mL  $\text{H}_2\text{O}_2$  (1%) and 2 mL enzyme extract. APX activity was assayed by monitoring the ascorbic acid dependent reduction of  $\text{H}_2\text{O}_2$  at 240 nm, as described by Asada (1992). The reaction mixture contained 50 mM potassium phosphate buffer (pH = 7.4), 0.5 mM ascorbate, 0.2 mL  $\text{H}_2\text{O}_2$  (1%), 0.1 mL enzyme extraction. GPX activity was determined at 420 nm by the method of Chang and Kao (1998). The reaction mixture contained 50 mM potassium phosphate buffer (pH = 7.4), 1 mL guaiacol (1%), 1 mL  $\text{H}_2\text{O}_2$  (1%) and 2.5 mL enzyme extract.

**Relative water content:** Chilling injury on leaves and roots was evaluated by changes in Relative Water Content (RWC). Relative water content was calculated using the formula has become below (Kuk *et al.*, 2003).

$$\text{RWC} = \left( \frac{1 - \text{Dry weight of tissue}}{\text{Fresh weight of tissue}} \right) \times 100$$

**Soluble sugar content:** Soluble sugars content was measured using phenol sulphuric method (Fales, 1979). The amount of 0.5 g of plant tissues (leaf or root) were sampled and homogenized with a mortar and pestle in 5 mL distilled water and then was filtered through Whatman No. 1 filter paper. Two milliliter of filtered extract transferred into tube and 1 mL of phenol 5% (w/v), 3 mL of sulphuric acid 98% were added. After 1 h the absorbance was recorded at 485 nm using UV-visible spectrophotometer (WPA model S2100).

## RESULTS AND DISCUSSION

It has been reported that some chilling-sensitive plants acclimate if they are exposed to a low temperature slightly above the threshold of chilling temperature, in a process analogous in some respects to the acclimation that occurs in perennial plants in the autumn. Therefore chilling injury can be alleviated if plants are exposed first to a low non-chilling temperature (Rikin *et al.*, 1979). It has been shown that acclimation of maize seedlings to otherwise lethal chilling temperatures by a milder cold pretreatment was accompanied by catalase and peroxidase transcript accumulation (Prasad *et al.*, 1994). Other correlations between cold-acclimation and antioxidant defense have also been reported (Tao *et al.*, 1998). To study the mechanisms of chilling injury or tolerance, most researchers utilize metabolic differences between chilling-sensitive and tolerant varieties as model systems (Pinhero *et al.*, 1997; Saruyama and Tamida, 1995). However, this system is confounded by genetic differences between sensitive and tolerant varieties. Therefore, it is difficult to interpret the observed metabolic differences in relation to mechanisms of chilling tolerance when using different varieties of plants. Using a chilling-sensitive variety that can be cold-acclimated is advantageous for studying the mechanisms involved in chilling tolerance because it eliminates the complexity of genetic differences. In this study, only one variety was used to demonstrate whether chilling tolerance can be induced in soybean plants by cold-acclimation and to examine whether an AOS-scavenging system is involved in tolerance to chilling stress. It has been reported that cold-acclimated rice plants shows higher tolerance to chilling stress than non-acclimated plants (Kuk *et al.*, 2003). Also, a similar phenomenon was demonstrated in other cold-sensitive species, such as maize, tomato and *Arabidopsis thaliana* (L.) Heynh (Venema *et al.*, 2000). Free radicals and other active derivatives of oxygen are

inevitable by-products of biological redox reactions. Active oxygen species inactivate enzymes and damage important cellular component. The increased production of toxic oxygen derivatives is considered to be a universal or common feature of stress conditions. Plants have evolved a wide range of mechanisms to contend with this problem. The antioxidant defence system of the plant comprises a variety of antioxidant molecules and enzymes (Arora *et al.*, 2002). We have investigated the activity of APX, CAT and GPX in both leaves and roots of cold-acclimated and non-acclimated soybean plants. Ascorbate peroxidase activity has mainly been reported from chloroplast and cytosol (Chen and Asada, 1989). However some recent studies have also reported its occurrence in mitochondria as well (Gomez *et al.*, 1999). This enzyme involved in Asada-Halliwell pathway of hydrogen peroxide scavenging that also involves various antioxidant enzymes (Arora *et al.*, 2002). Guaiacol peroxidase removes H<sub>2</sub>O<sub>2</sub> from apoplast and vacuole and utilizes L-ascorbic acid as electron donor. In an abiotic environment it uses phenolic compounds such as guaiacol as redox intermediate (Chanda and Singh, 2000). Catalase is another defense enzyme that is present in peroxisomes (Scandalios, 1993). In this study APX activity in leaves was found to be higher than roots. But significant changes was observed in roots. In recovery phase, cold-acclimated plants returned to normal condition more easily and rapidly than non-acclimated plants. Soybean plants exposed to chilling temperature exhibited all the aforementioned symptoms; however, cold-acclimated plants showed higher tolerance to chilling stress than non-acclimated plants. Cold-acclimated plants generally still showed similar level of injury as non-acclimated ones, but cold-acclimated plants showed the capability to recover from chilling injury. This was indicated by faster recovery of fresh weight of cold-acclimated plants compared with non-acclimated ones. The baseline levels of antioxidative enzyme activities were generally the same between cold-acclimated and non-acclimated leaves. No differences were found in APX, GPX and CAT activities in cold-acclimated and non-acclimated leaves during chilling and recovery period. CAT, APX and GPX activities in cold-acclimated and non-acclimated leaves were similarly affected by chilling temperature (Fig. 1-3). However, recovery in cold-acclimated leaves for APX, GPX and CAT activities were better than in non-acclimated leaves. Significant changes in GPX activity was observed between cold-acclimated and non-acclimated leaves 24 h after acclimation (Fig. 2, 3). GPX and APX activities in cold-acclimated leaves were higher than non-acclimated leaves.




Fig. 1: Changes in leaf CAT activity (U min<sup>-1</sup>) in cold-acclimated and non-acclimated soybean leaves during acclimation, chilling and recovery. Mean of 3 measurements±SE p = 0.05




Fig. 2: Changes in leaf APX activity (U min<sup>-1</sup>) in cold-acclimated and non-acclimated soybean leaves during acclimation, chilling and recovery. Mean of 3 measurements±SE p = 0.05




Fig. 3: Changes in leaf GPX activity (U min<sup>-1</sup>) in cold-acclimated and non-acclimated soybean leaves during acclimation, chilling and recovery. Mean of 3 measurements±SE p = 0.05

Activities of CAT, APX and GPX in cold-acclimated and non-acclimated roots during acclimation and chilling were generally the same (Fig. 4-6). Roots of cold-

acclimated plants showed higher activity of antioxidative enzymes during the phase of the recovery period compared to roots of non-acclimated plants. APX activity




Fig. 4: Changes in root CAT activity ( $\text{U min}^{-1}$ ) in cold-acclimated and non-acclimated soybean roots during acclimation, chilling and recovery. Mean of 3 measurements $\pm$ SE  $p = 0.05$




Fig. 5: Changes in root APX activity ( $\text{U min}^{-1}$ ) in cold-acclimated and non-acclimated soybean roots during acclimation, chilling and recovery. Mean of 3 measurements $\pm$ SE  $p = 0.05$




Fig. 6: Changes in root GPX activity ( $\text{U min}^{-1}$ ) in cold-acclimated and non-acclimated soybean roots during acclimation, chilling and recovery. Mean of 3 measurements $\pm$ SE

in cold-acclimated roots was higher than in non-acclimated roots in recovery phase. GPX activity in cold-acclimated roots was significantly higher than non-acclimated roots (Fig. 6). In all plants APX, CAT and GPX activities in cold-acclimated roots was higher than non-acclimated and cold-acclimated roots had higher potential to recover than non-acclimated plants. However, CAT and APX activity were similar between cold-acclimated and non-acclimated roots during recovery. No significant differences were found in CAT activity between cold-acclimated and non-acclimated roots during acclimation (Fig. 4).

The minimal impact of chilling on Relative Water Content (RWC), high photosynthetic efficiency of cold-acclimated plants and the full recovery of carbon fixation capability of cold-acclimated plants after chilling. Typical symptoms of chilling injury are wilting, yellowing of leaves and inhibition of growth. Leaf fresh weights of cold-acclimated and non-acclimated soybean seedlings were reduced in chilling period (Fig. 7, 8). Leaf and




Fig. 7: Changes in root relative water content (%) in cold-acclimated and non-acclimated soybean roots during acclimation, chilling and recovery. Mean of 3 measurements $\pm$ SE  $p = 0.05$




Fig. 8: Changes in leaf relative water content (%) in cold-acclimated and non-acclimated soybean leaves during acclimation, chilling and recovery. Mean of 3 measurements $\pm$ SE  $p = 0.05$




Fig. 9: Changes in root soluble sugars ( $\text{mg g}^{-1}$  FW) in cold-acclimated and non-acclimated soybean roots during acclimation, chilling and recovery. Mean of 3 measurements  $\pm$  SE  $p = 0.05$




Fig. 10: Changes in leaf soluble sugars ( $\text{mg g}^{-1}$  FW) in cold-acclimated and non-acclimated soybean leaves during acclimation, chilling and recovery. Mean of 3 measurements  $\pm$  SE  $p = 0.05$

root fresh weight of non-acclimated plants rapidly declined when exposed to chilling temperature and did not recover from the chilling treatment properly. Leaf and root fresh weight of cold-acclimated plants also declined significantly 24 h after exposure to chilling temperature. Although the cold-acclimated plants did not completely recover from chilling injury compared with untreated control plants, the cold-acclimated plants showed a general increase in leaf and root weight compared with non-acclimated plants during the recovery period.

Soluble sugars function as cryoprotectants and osmolites that protect cells from freezing damage (Xin and Browe, 2000). Shou *et al.* (2004) showed that 24 and 48 h cold-acclimations significantly increased the soluble sugar levels in plants. Accumulation of sucrose and other simple sugars that occurs with cold-acclimation also contributes to the stabilization of membrane as these

molecules can protect membranes against freeze-damage because increase in sugar content lowers the freezing point of cell solution (Mahajan, 2005). Especially sucrose can increase the effectiveness of the heat stable proteins in preventing protein denaturation (Robertson *et al.*, 1994). Soluble sugar levels were increased in roots. In cold-acclimated roots, soluble sugars content were started to increase in acclimation phase and downed in recovery phase. But in non-acclimated roots they increased in chilling phase suddenly (Fig. 9). In leaves changes were significant (Fig. 10). Soluble sugars content in non-acclimated leaves, in acclimation phase, decreased, then increased suddenly in chilling phase and subsequently decreased in recovery phase.

## REFERENCES

- Arora, A., R.K. Sairam and G.C. Srivastava, 2002. Oxidative stress and antioxidative system in plants. *Curr. Sci.*, 82: 10-25.
- Asada, K., 1992. Ascorbate peroxidase-A hydrogen peroxide scavenging enzyme in plants. *Physiol. Plant.*, 85: 235-241.
- Brasa, A.S., 2001. Crop Responses and Adaptations to Temperature Stress. In: *Mechanisms of chilling Injury and Tolerance*, Prasad, T.K. (Ed.). Food Products Press. New York, pp: 1-34.
- Chance, B. and A.C. Maehly, 1959. The Assay of Catalase and Peroxidase. In: *Methods of Biochemical Analysis*, Click, D. (Ed.). Vol. 1. Interscience Publishers. New York, pp: 357-425.
- Chanda, S.V. and Y. Singh, 2000. Changes in peroxidase and IAAoxidase activities during zea mays development. *Plant Physiol. Biochem.*, 35: 245-250.
- Chang, C.J. and C.H. Kao, 1998.  $\text{H}_2\text{O}_2$  metabolism during senescence of rice leaves: Changes in enzyme activities in light and darkness. *Plant Growth Regul.*, 25: 11-15.
- Chen, G.X. and K. Asada, 1989. Ascorbate peroxidase in tea leaves: Occurrence of two isozymes and differences in their enzymatic and molecular properties. *Plant Cell Physiol.*, 30: 987-998.
- Fadzillah, N.M., V. Gill, R.P. Pinch and R.H. Burdon, 1996. Chilling, oxidative stress and antioxidant responses in shoot cultures of rice. *Planta*, 199: 552-556.
- Fales, F., 1979. The assimilation and degradation of carbohydrates by yeast cells. *J. Biol. Chem.*, 193: 113-124.
- Foyer, C.H., M. Lelandais, C. Galap and K.J. Kunert, 1991. Effect of elevated cytosolic glutathione reductase activity on the cellular glutathione pool and photosynthesis in leaves under normal and stress conditions. *Plant Physiol.*, 97: 863-872.

- Gomez, J.M., J.A. Hernandez, A. Jimenez, L.A. del Rio and F. Sevilla, 1999. Differential responses of antioxidative enzymes of chloroplasts and mitochondria to long-term NaCl stress of pea plants. *Free Radic. Res.*, 31: 11-18.
- Hodges, D.M., C.J. Andrews, D.A. Johnson and R.I. Hamilton, 1996. Antioxidant compound responses to chilling stress in differentially sensitive inbred maize lines. *Physiol. Plant*, 98: 685-692.
- Kuk, Y.I., J.S. Shin, N.R. Burgos, T.E. Hwang, Oksoo, B.H. Cho, S. Jung and J.O. Guh, 2003. Antioxidative enzymes offer protection from chilling damage in rice plants. *Crop Sci.*, 43: 2109-2117
- Lee, D.H. and C.B. Lee, 2000. Chilling stress-induced changes of antioxidant enzymes in the leaves of cucumber: In gel enzyme activity assay. *Plant Sci.*, 159: 75-85.
- Mahajan, S. and N. Tuteja, 2005. Cold, salinity and drought stresses: An overview, 10.018
- Michalski, W.P. and Z. Kanjuga, 1982. Photosynthetic apparatus of chilling sensitive plants: Reversibility by light of cold and dark-induced inactivation of cyanide-sensitive superoxide dismutase activity in tomato leaf chloroplasts. *Biochem. Biophys. Acta*, 680: 250-257.
- Oidaira, H., S. Satoshi, K. Tomokazu and U. Takashi, 2000. Enhancement of antioxidant enzyme activities in chilled rice seedlings. *Plant Physiol.*, 156: 811-813.
- Pinhero, R.G., M.V. Rao, G. Paliyath, D.P. Murr and R.A. Fletcher, 1997. Changes in activities of antioxidant enzymes and their relationship to genetic and paclobutrazol induced chilling tolerance of maize seedlings. *Plant Physiol.*, 114: 685-704.
- Prasad, T.K., M.D. Anderson B.A. Martin and C.R. Stewart, 1994. Evidence for chilling induced oxidative stress in maize seedlings and a regulating role for hydrogen peroxidase. *Plant Cell*, 6: 65-74.
- Rikin, A., D. Atsmon and C. Gitler, 1979. Chilling injury in cotton (*Gossypium hirsutum* L.): Prevention by abscisic acid. *Plant Cell Physiol.*, 20: 1537-1546.
- Robertson, A.J., M. Ishikawa, L.V. Gusta and S.L. Mackenzie, 1994. Abscisic acid-induced heat tolerance in *Bromus inermis* Leyss cell-suspension cultures. *Plant Physiol.*, 105: 181-190.
- Saruyama, H. and M. Tamida, 1995. Effect of chilling on activated oxygen-scavenging enzymes in low temperature-sensitive and-tolerant cultivars of rice (*Oryza sativa* L.). *Plant Sci.*, 109: 105-113.
- Scandalios, G.J., 1993. Oxygen stress and superoxide dismutase. *Plant Physiol.*, 101: 7-12.
- Scebba, F., L. Sebastiani and C. Vitagliano, 1999. Protective enzymes against activated oxygen species in wheat (*Triticum aestivum* L.) seedlings: Responses to cold acclimation. *J. Plant Physiol*, 155: 762-768.
- Shou, H., P. Bordallo, J. Fan, J.M. Yeakley, M. Bibikova, J. Sheen and K. Wang, 2004. Expression of an active tobacco mitogen-activated protein kinase enhances freezing tolerance in transgenic maize. *PNAS*, 101: 3298-3303.
- Tao, D.L., G. Oquist and G. Wingsle, 1998. Active oxygen scavengers during cold-acclimation of Scots pine seedlings in relation to freezing tolerance. *Cryobiology*, 37: 38-45.
- Upadhyaya, A., T.D. Davis, R.H. Walser, A.B. Galbraith and N. Sankhla, 1989. Uniconazole-induced alleviation of low-temperature damage in relation to antioxidant activity. *Hortscience*, 24: 955-957.
- Venema, J.H., L. Villerius and P.R. Van Hasselt, 2000. Effect of acclimation to suboptimal temperature on chilling-induced photo-damage: Comparison between a domestic and a high-altitude wild Lycopersicon species. *Plant Sci.*, 152: 153-163.
- Wanner, L.A. and O. Junttila, 1999. Cold-induced freezing tolerance in *Arabidopsis*. *Plant Physiol.*, 120: 391-399.
- Welling, A. and E.T. Palva, 2006. Molecular control of cold-acclimation in trees. *Physiol. Plant.*, 127: 167-181.
- Wise, R.R. and A.W. Naylor, 1987a. The peroxidative destruction of lipids during chilling injury to photosynthesis and ultrastructure. *Plant Physiol.*, 83: 272-277.
- Wise, R.R. and A.W. Naylor, 1987b. Chilling-enhanced photooxidation. Evidence for the role of singlet oxygen and superoxide in the breakdown of pigments and endogenous antioxidants. *Plant Physiol.*, 83: 278-282.
- Xin, Z. and J. Browse, 2000. *Plant Cell Environ*, 23: 893-902.