

Journal of Biological Sciences

ISSN 1727-3048

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Prevalence of Epstein-Barr Virus Latent Membrane Protein-1 in Jordanian Patients with Hodgkin's Lymphoma and Non-Hodgkin's Lymphoma

¹F. Irshaid, ¹A. Jaran, ¹F. Dilmi, ²K. Tarawneh, ³R. Hadeth and ³Ahad Al-Khatib

¹Department of Biological Sciences, Al Al-Bayt University, P.O. Box 130040, Al-Mafraq 25113, Jordan

²Department of Biological Sciences, Faculty of Science, Mutah University, Mutah, Jordan

³Department of Pathology, Faculty of Medicine, Jordan University of Sciences and Technology, Irbid, Jordan

Abstract: The aim of this study was to estimate the frequency of EBV infection in Hodgkin's Lymphoma (HL) and Non-Hodgkin's Lymphoma (NHL) occurring in Jordanian patients. A total of 55 patients with lymphoma were examined. The diagnosis was confirmed by staining the slides of these cases with hematoxylin and eosin. The presence of EBV was detected by immunostating for expression of latent membrane protein-1 (LMP-1). Of 55 patients, 31 and 25 were diagnosed as HL and NHL, respectively. The majority of the HL cases exhibited Mixed Cellularity (MC) subtype. The high grade was found to be the commonest subtype of NHL. The frequency of LMP-1 expression occurred more frequent in patients with HL (60.0%) than in patients with NHL (32.0%). The frequency of LMP-1 expression was higher in patients with MC subtype (61.11%) than in patients with nodular sclerosis (28.57%). No age or gender difference in occurrence of EBV infection was observed among HL patients. By contrast, the frequency of LMP-1 expression in NHL patients aged below 50 was lower (16.66%) than in NHL patients aged 50 or above (46.15%). In addition, EBV infection was more frequent in females with NHL (38.46%) than in male with NHL (25%). In NHL cases, the frequency of LMP-1 expression in intermediate grade (60.0%) was high compared to low (37.5%) or high grades (16.7%). Therefore, analysis of LMP-1 expression suggests an important role for this viral oncogene in the pathogenesis of EBV-associated malignant lymphomas. These data support the previous findings that people with EBV may develop lymphoma and that efforts to maintain low lymphoma should be considered for people with EBV infection.

Key words: Expression, hematoxylin, infection, malignant, oncogene

INTRODUCTION

Lymphoma is broadly divided into Hodgkin's Lymphoma (HL) and Non-Hodgkin's Lymphoma (NHL) based on classification of the World Health Organization (Harris *et al.*, 1999). The HL is also subclassified into two four categories termed Nodular Sclerosis (NS), Mixed Cellularity (MC), Lymphocyte Depletion (LD) and Lymphocyte-Rich (LR). In addition, the working formulation classification has been adopted in United States. According to this system NHL has been divided into low, intermediate and high grades (Harris *et al.*, 1999; Lu, 2005). In Europe and elsewhere, the Kiel classification of NHL has been accepted widely. Based on the Kiel classification, NHL is divided into B-cell and T-cell lineages based on available immunophenotypic data and into low or high grades based on cell morphology (Lu, 2005).

Data from The International Agency for Research on cancer indicated that at least 287,000 of NHL and 62,000 of HL new cases were annually diagnosed world wide. In USA, NHL is already ranked fifth among white male cancers for both incidence and mortality and sixth for female cancers (Jemal *et al.*, 2002). In addition, the prevalence of NHL has increased over the past two decades, with median age of 55-60 years at presentation and the incidence increases with advancing age. Moreover, according to data from the Jordanian National Cancer Registry in 2004, the estimated number of newly diagnosed cancer cases was approximately 5000 cases per year. This estimate may be less far from the accurate number (Stjernsward *et al.*, 2007). This is because cancer database is generally incomplete or inaccurate and several numbers of cases were officially not recognized or reported. Importantly, because cancer among these patients is commonly discovered in advance stage, about

70% of these patients will die of cancer within the first year of their diagnosis. In addition, approximately 5.7% of these cancer cases were diagnosed with lymphoma.

Recently, there was some evidence of complete cure of lymphoma by irradiation and chemotherapy. The best results were achieved for those patients when intensifying radiotherapy and chemotherapy were administered earlier rather than later in the course of the disease (Gribben, 2007; Travis, 2007). However, it has been become clear that death due to second cancers are now the most common cause of mortality among long-term survivors of HL, followed by cardiac disease (Hemminki *et al.*, 2008; Knight *et al.*, 2009).

The search for a cause of cancers has revealed that the Epstein Barr Virus (EBV) appears to be associated with some cancers including lymphoma (Pattle and Farrell, 2006; Kapatai and Murray, 2007). The EBV is a very common virus that is already infecting high percent of population worldwide and persisting for the lifetime in the host (Cohen, 2000). Importantly, despite the fact that EBV is a common infection, several studies postulated that EBV is linked to the development of multiple cancers, including Burkitt's lymphoma, HL, NHL and some diffuse large B-cell lymphomas (Serraino *et al.*, 2005; Pattle and Farrell, 2006; Kapatai and Murray, 2007; Asano *et al.*, 2009). According to these studies, the association between lymphoma and EBV came from the observation that EBV is the causative agent of infectious mononucleosis. So that patients with a previous history of infectious mononucleosis have an elevated risk of developing lymphoma. Furthermore, a reasonable interpretation for the association of EBV with Hodgkin's disease came from a number of studies which reported that EBV can transform resting human B-cells into permanently proliferating lymphoblastoid cell lines that constitutively expresses LMP-1 (Staudt, 2000; Cohen, 2000; Thorley-Lawson and Gross, 2004; Kim *et al.*, 2006; Kapatai and Murray, 2007; Zhao *et al.*, 2008). These studies also reported that the transcription of this viral oncogene was demonstrated in various malignant lymphoma cells. They also proposed that the expression of LMP-1 might provide growth and survival signals that enhance tumor growth and hence contribute to the malignant phenotype of B-cells. They also postulated that a germinal-center B cell might acquire a mutation during infectious mononucleosis that blocks its differentiation. Thus, we anticipate that an earlier detection and treatment of EBV infection most likely would contribute to decrease the rate of lymphoma incidence among our population. Moreover, little is known about the risk of EBV infection for the substantial majority of lymphoma patients in Jordan. In this study our goal, therefore, was to estimate

the prevalence of EBV infection in Jordanian patients diagnosed with HL and NHL using immunohistochemistry technique with primary monoclonal antibody against EBV-LMP-1.

MATERIALS AND METHODS

Samples collection: A total of 55 samples diagnosed with malignant lymphoma were randomly selected from the archive of pathology laboratory of King Hussein Hospital, Amman, Jordan, between January, 2006 and December, 2008. The samples were available as paraffin embedded tissues. All lymphoma tissues were examined microscopically and then divided into HL and NHL. These HL and NHL samples were further subdivided into different grades or subtypes by Dr. Raji Hadeth, a pathologist from faculty of Medicine, Jordan University of Sciences and Technology, Irbid, Jordan.

Hematoxylin and eosin (H and E) staining: Formalin-fixed paraffin tissues were sectioned at 4 µm thick. These sections were deparaffinized by heating and then immediately cleared in xylene. This was followed by rehydration through serial dilutions of alcohol. These sections were counterstained with hematoxylin and stained with eosin before dehydration. Finally, these sections were mounted with distrenedibutylphthalate-xylene (DPX) and examined under light microscope (Nikon, Japan).

Immunohistochemistry study: Immunohistochemical staining was done on 4 µm thick consecutive sections of formalin fixed, paraffin-embedded tissue specimens. Tissue section slides were deparaffinized and rehydrated as described above. Then, slides were immersed in 3% hydrogen peroxidase in methanol for 10 min to block endogenous peroxidase activity and rinsed in running water. Subsequently slides were treated under pressure in the commercial solution (Reveal 10X, Biocare Medical, CA, USA) in the Decloaking chamber (Biocare Medical) to retrieve the antigens and block endogenous biotin. Then, slides were allowed to cool at room temperature for 20 min. This was followed by washing and incubation with Phosphate Buffer Saline (PBS) (diluted at 1:10) for 15 min. Slides were then incubated with primary mouse monoclonal antibody against EBV-LMP-1 for 30 min (diluted at 1:50, from Biocare medical). One drop of biotinylated goat anti-mouse (diluted at 1:10; from BioGenex, San Ramon, CA, USA) was added to each slide and washed well with PBS. After that, slides were incubated with streptavidin peroxidase (diluted at 1:10, from BioGenex) for 20 min at room temperature and

washed well with PBS. Slides were then incubated with 3,3-Diaminobenzidine chromogen (from BioGenex). This was followed by counterstained with hematoxylin and then stained with eosin before dehydration and mounted with DPX. Finally, all slides were examined under light microscope (Nikon, Japan). A previously well-known negative case for EBV was used as negative control and treated as described above.

Immunostaining assessment and evaluation: In NHL specimens with less than 5% EBV-positive tumor cells were judged as negative and specimens with more than 5% EBV-positive tumor cells were interpreted as positive. It is worth to mention that the percentage of EBV-positive staining whether membrane or paranuclear were taken into consideration. Whereas, in HL, the specimen is considered EBV-positive if any Reed-Sternberg (RS) cell was membrane or paranuclear staining positive. This is because the HL is usually characterized by the scarcity of the neoplastic RS cell (1-10%).

RESULTS AND DISCUSSION

In the present study, we examined 55 samples which fulfilled the lymphoma criteria. There were 30 patients HL and 25 patients were NHL. Male patients with HL were 20 (66.7%) while 10 (33.3%) were female patients with HL. The male versus female ration was 2:1. Of 30 HL cases, 24 (80%) were below the age of 50 years while six out of the 30 cases (20%) occurred in age 50 years or above (Table 1). As can be seen in Fig. 1, the age group distribution of our HL cases was as follows: 0-9 years 1 (3.3%), 10-19 years 10 (33.3%), where the highest incidence occurred, 20-29 years 5 (16.7%), 30-39 years 4 (13.3%), 40-49 years 4 (13.3%), 50-59 years 6 (19.9), 60-69 years 4 (13.3%) and 70 years and above 1 (3.3%). The median age was 28 years and ranged from 4-70 years. On other hand, there were 12 males (48%) and 13 female (52%) with NHL. The male versus female ration was 1:0.92. Of 25 NHL cases, 12 (48%) were below the age of 50 years and 13 (52%) were at the age of 50 years and above. The age group distribution of our NHL cases was as follows (Fig. 1): 0-9 years 4 (16%), 10-19 years 3 (12%), 20-29 years 1 (4%), 30-39 years 1 (4%), 40-49 years 3 (12%),

50-59 years 3 (12%), 60-69 years 4 (16%) and 70 years and above 6 (24%). Unlike HL, no peak of incidences in NHL was noted among the different age groups. The median age was 50 years and ranged from 3-78 years.

The HL samples were further divided into subgroups according to the WHO classification criteria. The HL cases were divided into 5 subtypes. Based on this classification, there were 60% (18/30) mixed cellularity (MC), 23.33% (7/30) Nodular Sclerosis (NS), 6.66% (2/30) Lymphocyte Depletion (LD), 4.33% (1/30) Lymphocyte Rich (LR) and 6.66% (2/30) unclassified (UC), as shown in Table 2. On other hand, NHL was graded into high, intermediate and low grade according to the International Working Formulation system. There were 48% (12/25) of high grade, 20% (5/25) of intermediate and 32% (8/25) of low grade (Table 2).

The presence of EBV was confirmed by the expression of LMP-1 using specific monoclonal antibodies. LMP-1 usually displays a typical membrane staining in tumor cells. In this work, immunohistochemical staining results of the investigated LMP-1 marker were used to classify the HL or NHL samples into positive only when tumor cells stained with the corresponding marker and thus, the absence of such staining pattern was considered negative. The results of expression of LMP-1 of EBV in HL patients are shown in Fig. 2a-c. Based on these staining results, high incidence of EBV-LMP-1




Fig. 1: Distribution of Hodgkin's lymphoma and non-Hodgkin lymphoma patients according to age

Table 2: Distribution of Hodgkin's lymphoma (HL) and non-Hodgkin's lymphoma (NHL) patients according to subtype and grade

Type of lymphoma	Subtype	Results of diagnosis	
		N	%
HL	Mixed cellularity	18	60.00
	Nodular sclerosis	7	23.33
	Lymphocyte depletion	2	6.66
	Lymphocyte rich	1	4.33
	Unclassified	2	6.66
NHL	High	12	48.00
	Intermediate	5	20.00
	Low	8	32.00

Table 1: Distribution of Hodgkin's lymphoma (HL) and non-Hodgkin's lymphoma (NHL) patients according to gender and age at diagnosis

Type of lymphoma	Age groups		Gender	
	<50 years	≥50 years	Male	Female
	N (%)			
HL	24 (80.0)	6 (20.0)	20 (66.6)	10 (33.3)
NHL	12 (48.0)	13 (52.0)	12 (48.0)	13 (52.0)

protein was observed in HL cases 60% (18/30), in contrast to the relatively low incidence was also seen in NHL cases 32% (8/25). Present results also indicated that the EBV-LMP-1 protein expression was observed in 15 of 24 (62.50%) HL cases with below 50 years of age and a closely similar frequency was detected in 3 of 6 (50.00%) HL cases with 50 years of age and above (Table 3). In addition, a similar frequency of EBV-LMP-1 expression was seen in both male 60.00% (12/20) and female 60.00% (6/10) with HL, as shown in Table 3.

With regards to subtype, the presence of EBV-LMP-1 was detected in 11 of 18 (61.11%) MC cases and 2 of 7 (28.57%) NS cases as shown in Table 4. On other hand, expression of LMP-1 of EBV was observed in all cases of

other subtypes, including LD, LR and UC. Along the same line, Fig. 3a-c show the results of expression of LMP-1 of EBV in NHL patients. With respect to gender patients with NHL, the presence of EBV was detected in 3 of 12 (25%) male cases and 5 of 13 (38.46%) female cases (Table 3). Present data also revealed that EBV-LMP-1 was found in 2 of 12 (16.66%) NHL cases with below 50 years of age, in contrast to the relatively high incidence was observed in 6 of 13 (46.15%) NHL cases with 50 years of age and above. With regard to grade, the presence of EBV was detected in 2 of 8 (25.00%) low grade cases, 3 of 5 (60.0%) intermediate grade cases and 3 of 12 (25.00%) high grade cases (Table 4).

Our current study indicated that the HL accounted 55 and 45% for NHL. Previously, it was reported that NHL accounted for as high as 60.7 and 78.4% of all lymphoma cases among the Jordanian population (Haddadin, 2005). Additionally, different studies from the Middle East have also reported higher prevalence of NHL compared to HL, namely Oman with 65% NHL and 35% HL (Bamanikar *et al.*, 1995), Bahrain with 66.7% NHL and 33.3% HL (Shome *et al.*, 2004) and United Emirates with 59% NHL and 41% HL (Castella *et al.*, 2001). This finding might be explained by the nature of our samples. In fact, our samples were randomly selected from 300 available samples, of which 190 samples were found in well preserved and adequate amount that might explain the high incidence of HL. It is also important to note that some of lymphomas previously diagnosed as HL are now classified as NHL and this shift is responsible for the declining incidence of HL.

Available data reported that lymphoma can occur in all age group (Cohen, 2000; Shome *et al.*, 2004; Serraino *et al.*, 2005; Haddadin, 2005; Pattle and Farrell, 2006). Present data indicated that a clear difference observed in the incidence of HL among different age groups with the majority of cases (80%) occurring in the

Table 3: Frequency of expression of latent membrane protein (LMP-1) of EBV in Hodgkin's lymphoma (HL) and non-Hodgkin's lymphoma (NHL) patients according to gender and age at diagnosis

Parameters	Type of lymphoma			
	HL		NHL	
	N	%	N	%
Age groups				
<50 years	15/24	62.50	2/12	16.66
≥50 years	3/6	50.00	6/13	46.15
Gender				
Male	12/20	60.00	3/12	25.00
Female	6/10	60.00	5/13	38.46

Table 4: Frequency of expression of latent membrane protein (LMP-1) of EBV in Hodgkin's lymphoma (HL) and non-Hodgkin's lymphoma (NHL) patients according to subtype and grade

Type of lymphoma	Subtype	Positive	
		N	%
HL mixed	Cellularity	11/18	61.11
	Nodular sclerosis	2/7	28.57
	Lymphocyte depletion	2/2	100.00
	Lymphocyte rich	1/1	100.00
	Unclassified	2/2	100.00
	High	2/8	25.00
NHL	Intermediate	3/5	60.00
	Low	3/12	25.00




Fig. 2: Expression of latent membrane protein-1 of EBV in Hodgkin's lymphoma (HL). (a) HL patient with mixed cellularity. (b) HL patient with nodular sclerosis. (c) HL patient with mixed cellularity (Negative control). For all slides magnifications were 40X. Arrows indicate the position of RS cell




Fig. 3: Expression of latent membrane protein-1 of EBV in non-Hodgkin's lymphoma (NHL) patients. (a) NHL patient with low grade. (b) NHL patient with intermediate grade. (c) NHL patient with high grade. For all slides, magnifications were 100X. Arrows indicate the tumor cell

younger age groups below 50 years versus 20% of cases occurring above or equal that age. Similar findings were also reported in certain areas such as Cyprus with 84.3% of HL below 50 years and Egypt with 84.4% below 50 years (Soliman and Boffetta, 2004). In addition, 33.3% of the HL cases fell within the age interval 10 to 19 years which represented an early major peak of incidence. An observation in this studies correlates very well with the previous research findings, in that HL age specific incidence with one peak between 11 and 20 years was reported in other studies (Almasri, 2004; Haddadin, 2005). The slightly difference between these study and our study are more likely due to small size of our sample. Moreover, it has been reported that the overall incidence of HL has decreased in parallel with a slight increase in young adults during the past decades world wide.

Unlike HL, 52% of NHL cases fill above age of 50 years and no peak of incidence was noted among NHL cases at different age groups. This result is in agreement with the results of study in Egypt with 49.5% NHL cases below 50 years (Soliman and Boffetta, 2004). Our and these data clearly suggest that NHL cases seem to be distributed somewhat regularly within different age groups. By contrast, HL seems to occur more frequently in male than in female, with a male to female ratio: 2:1. The overall frequency of HL in male and female is entirely consistent with the published literature, minor differences almost certainly reflecting differences in sample size. However, no sex difference was observed in NHL where the sex ratio was approximately equal.

Among the various subtypes of HL, the MC was found to be the commonest subtype in present study (60%) followed by the NS (23.3%). Present results are similar to recent estimate from Iraq with predominance of MC (67.5%) followed by NS (12%) (Pileri *et al.*, 2002). However, these results are in contrast with western countries where the NS is the most frequent subtype. For

example, in Italy and USA, NS was estimated to be about 75-80% of all HL cases (Pileri *et al.*, 2002). Moreover, the most frequent grades of NHL were the high grade (48%), followed by low grade (32%). This result differed from previous study in Saudi Arabia which reported that low, intermediate and high grades of NHL constituted about 18, 37 and 29%, respectively (Thomas and Abdullah, 1995). These unexpected findings, however, should be taken cautiously in that it difficult to explain. In addition, these small differences may be a coincident or required another study with larger sample size. Taken together, the variations in the prevalence of NHL in various studies could be due to different study methodologies of classifications, difference in sample sizes and/or misclassification due to the unspecialized personnel used to classify lymphoma. It could also be due to some environmental factors or other unknown causes.

It has been proposed that LMP-1 is expressed in many EBV-associated cancers and is responsible for most of the altered cellular growth properties that are induced by EBV infection (Cohen, 2000; Thorley-Lawson and Gross, 2004; Kim *et al.*, 2006; Kapatai and Murray, 2007; Zhao *et al.*, 2008). In this work, staining results of the LMP-1 were used to sub-classify the NH and NHL into positive EBV or negative EBV. Our staining data revealed that the frequency of expression of LMP-1 of EBV was higher in HL patients (60%) than in NHL patients (32%). These data are in agreement with the above data which revealed that the HL occurred more frequent than NHL in Jordanian patients with lymphoma. Also, our current result correlates very well with the previous research findings (Vassallo *et al.*, 2001; Ridha, 2004). These observations further support the notion that expression of LMP-1 of EBV might involved in development and progression of HL. However, other study showed that incidence of EBV in developed country was 50% in their HL cases (Cohen, 2000; Thorley-Lawson and Gross, 2004;

Serraino *et al.*, 2005; Kim *et al.*, 2006; Pattle and Farrell, 2006; Kapatai and Murray, 2007), whereas, others reported that EBV expression was more than 90% in underdeveloped countries (Vassallo *et al.*, 2001; Pileri *et al.*, 2002; Zhao *et al.*, 2008; Ridha, 2004). Taken together, these findings also suggest that efforts to maintain low lymphomas must be considered for people with EBV infection.

With regard to histological subtypes and their association with EBV, our findings indicated that EBV expression was higher in the MC (61.11%), followed by NS (28.57%). These results were in consistent with results in previous study which showed that the EBV expression was higher in MC (66.6%), followed by NS (46.1%) (Vassallo *et al.*, 2001). It is also worth to note that all cases of the LD, LP and UC subtypes were found to be positive for EBV. The high prevalence of EBV in these subtypes should be interpreted cautiously due to the small number of cases in each category.

The presence of EBV seems to occur more frequent in HL (60.0%) than NHL cases (32.0%). These results were in agreement with some previously studies which also found 30% EBV positive in their NHL cases (Bossolasco *et al.*, 2002; Ridha, 2004). On other hand, different studies reported higher association of the EBV with NHL as high as 80% in other developing countries (Cohen, 2000; Rama Rao *et al.*, 2000). It is also worth mentioning that Burkitt's lymphoma is a high-grade malignant NHL that is most commonly associated with EBV infection (Thorley-Lawson and Gross, 2004; Ridha, 2004; Rama Rao *et al.*, 2000). Furthermore, in Africa, Burkitt's lymphoma is associated with *Plasmodium falciparum* malaria and tumors usually present in the jaw, over 90% of these cases are associated with EBV. In the United States, patients with Burkitt's lymphoma usually present with abdominal tumors, only 20% of which are associated with EBV (Cohen, 2000). Therefore, the low rate of LMP-1 of EBV positive in NHL cases that has been seen in present study may be due to little cases of Burkitt's lymphoma in our sample. This was clearly evident from our study which reported only one case in our sample. It is important to mention that the mode of transmission in our population is still uncertain. Therefore, it is worth to investigate the mode of transmission of EBV in our population with larger sample sizes of lymphoma patients.

Furthermore, the frequency of EBV in intermediate grade was much higher with approximately 60% of cases with EBV positive compared to the high or low grade cases, where about 25% were EBV positive. The overall frequency of expression of LMP-1 of EBV in these different grades is somewhat consistent with the published literature with minor differences. Taken

together, the above studies confirm that the association of EBV with lymphoma is strongly but variably linked to various environmental factors prevailing in different geographic regions. These include the economic status, geographical location, ethnic variations and health conditions of studied areas as well as the great diversity of lymphoma subtypes.

In conclusion, present results indicated for the first time that the presence of HL cases seems to occur more frequent in our population. Similarly, frequency of expression of LMP-1 of EBV was higher in HL patients than in NHL patients. The data also showed that the age, sex and clinical characteristics of Jordanian patients with HL and NHL included in this study are slightly different from those of neighbor countries. Furthermore, we suggest that a high sensitive technique must be carried out to verifying our observations.

The present study also provided evidence supporting the previous findings that people with EBV might develop lymphoma. Thus, there is an urgent need to educate our people at different ages about the potential threat that EBV infection might cause to them. These findings also provided an additional reason for doctors to take EBV infection very seriously in patients with lymphomas. In addition, our results should encourage physicians to achieve complete viral suppression treatment for those patients with persisting EBV infection in an attempt to minimize the risk of this life-threatening disease.

ACKNOWLEDGMENTS

This study has been supported in part by the Deanship of Scientific Research at Al Al-Bayt University. The financial and technical supports from Al Al-Bayt University are gratefully acknowledged. We would also like to thanks Dr. Ali Shouter, medical staff in forensic laboratory in Irbid and medical research staff in Jordan University of Science and Technology for their valuable advice, technical support and assistance.

REFERENCES

- Almasri, N.M., 2004. Hodgkin's lymphoma in north Jordan. Does it have different pattern. Saudi Med. J., 25: 1971-1981.
- Asano, N., K. Yamamoto, J.I. Tamaru, T. Oyama and F. Ishida *et al.*, 2009. Age-related Epstein-Barr virus (EBV)-associated B-cell lymphoproliferative disorders: Comparison with EBV-positive classic Hodgkin lymphoma in elderly patients. Blood, 113: 2629-2636.

- Bamanikar, S., S. Thunold, K.R. Devi and A. Bamanikar, 1995. The pattern of malignant lymphoma in Oman. *Trop. Med. Hyg. J.*, 98: 351-354.
- Bossolasco, S., P. Cinque, M. Ponzoni, M.G. Vigano, A. Lazzarin, A. Linde and K.I. Falk, 2002. Epstein-Barr virus DNA load in cerebrospinal fluid and plasma of patients with AIDS-related lymphoma. *J. NeuroVirol.*, 8: 432-438.
- Castella, A., S. Joshi, T. Raaschou and N. Mason, 2001. Pattern of malignant lymphoma in the United Arab Emirates. *Acta Oncol.*, 40: 660-664.
- Cohen, J.I., 2000. Epstein-Barr virus infection. *N. Engl. J. Med.*, 343: 481-492.
- Gribben, J.G., 2007. How I treat indolent lymphoma. *Blood*, 109: 4617-4626.
- Haddadin, W.J., 2005. Malignant lymphoma in Jordan: A retrospective analysis of 347 cases according to the World Health Organization classification. *Ann. Saudi Med.*, 25: 398-403.
- Harris, N.L., E.S. Jaffe and J.O. Armitage and M. Shipp, 1999. Lymphoma classification: From R.E.A.L. to W.H.O. and beyond. *Cancer: Principles Practice Oncol.*, 13: 1-14.
- Hemminki, K., P. Lenner, J. Sundquist and J.L. Bermejo, 2008. Risk of subsequent solid tumors after non-Hodgkins lymphoma: Effect of diagnostic age and time since diagnosis. *J. Clin. Oncol.*, 26: 1850-1857.
- Jemal, A., A. Thomas, T. Murray and M. Thun, 2002. Cancer statistics. *Cancer J. Clin.*, 52: 23-47.
- Kapatai, G. and P. Murray, 2007. Contribution of the Epstein Barr virus to the molecular pathogenesis of Hodgkin lymphoma. *J. Clin. Pathol.*, 60: 1342-1349.
- Kim, L.H., S.C. Peh and S. Poppema, 2006. Expression of retinoblastoma protein and p16 proteins in classic Hodgkins lymphoma: Relationship with expression of p53 and presence of Epstein-Barr virus in the regulation of cell growth and death. *J. Human Pathol.*, 37: 92-100.
- Knight, J.S., A. Tsodikov, D.M. Cibrik, C.W. Ross, M.S. Kaminski and D.W. Blayney, 2009. Lymphoma after solid organ transplantation: Risk, response to therapy and survival at a transplantation center. *J. Clin. Oncol.*, 27: 3354-3362.
- Lu, P., 2005. Staging classification of lymphoma. *Seminars Nulc. Med.*, 35: 160-164.
- Pattle, S.B. and P.J. Farrell, 2006. The role of Epstein-Barr virus in cancer. *Expert Opin. Biol. Ther.*, 6: 1193-1205.
- Pileri, S.A., S. Ascani, L. Leoncini, E. Sabattini and P.L. Zinzani *et al.*, 2002. Hodgkin's lymphoma: The pathologist's viewpoint. *J. Clin.*, 55: 162-176.
- Rama Rao C., M.I. Gutierrez, K. Bhatia, F. Fend and J. Franklin *et al.*, 2000. Association of Burkitt's lymphoma with the Epstein-Barr virus in two developing countries. *Leukemia Lymphoma*, 39: 329-337.
- Ridha, W.K., 2004. Epstein Barr virus-mediated deregulation of cell cycle pathways in malignant lymphomas: Implication of NFkB, p53, p27 and the mutant p21 Ras using in situ technique. Ph.D. Thesis, University of AL-Nahrain, Iraq.
- Serraino, D., P. Piselli, C. Angeletti, M. Scuderi and M.R. Ippolito, 2005. Infection with Epstein-Barr virus and cancer: An epidemiological review. *J. Biol. Regul. Homeost. Agents*, 19: 63-70.
- Shome, D.K., S.M. George, F. Al-Hilli and A.A. Satir, 2004. Spectrum of malignant lymphomas in Bahrain. Leitmotif of a regional pattern. *Saudi Med. J.*, 25: 146-167.
- Soliman, A.S. and P. Boffetta, 2004. Lymphoma and leukemia. Middle East Cancer Consortium (MECC) Registries. http://seer.cancer.gov/publications/mecc/mecc_lymphoma.pdf.
- Staudt, L.M., 2000. The molecular and cellular origins of Hodgkin's disease. *J. Exp. Med.*, 191: 207-212.
- Stjernsward, J., F. Ferris, S. Khleif, W. Jamous and I. Treish *et al.*, 2007. Jordan palliative care initiative: A WHO demonstration project. *J. Pain Symptom Manage.*, 33: 628-633.
- Thomas, J.O. and M.A. Abdullah, 1995. Malignant lymphoma in western province of Saudi Arabia. *East Afr. Med. J.*, 72: 355-358.
- Thorley-Lawson, D.A. and A. Gross, 2004. Persistence of the Epstein-Barr virus and the origins of associated lymphomas. *N. Engl. J. Med.*, 350: 1328-1337.
- Travis, L.B., 2007. Evaluation of the risk of therapy-associated complications in survivors of Hodgkin Lymphoma. *Hematology*, 1: 192-196.
- Vassallo, J., K. Metze, F. Traina, C. de Souza and I. Lorand-Metze, 2001. Expression of Epstein-Barr virus in classical Hodgkin's lymphomas in Brazilian adult patients. *Haematologica*, 86: 1227-1228.
- Zhao, P., Y. Lu, L. Liu and M. Zhong, 2008. Aberrant expression of ID2 protein and its correlation with EBV-LMP1 and p16 (INK4A) in Classical Hodgkins Lymphoma in China. *Bio. Med. Central J.*, 8: 379-379.