



Journal of Environmental Science and Technology

ISSN 1994-7887

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Evaluation of a New Egyptian Probiotic by African Catfish Fingerlings

¹A.M. Abdelhamid, ¹A.I. Mehrim, ²M.I. El-Barbary,

³S.M. Ibrahim and ¹A.I. Abd El-Wahab

¹Department of Animal Production, Faculty of Agriculture,
Al-Mansoura University, Al-Mansoura, Egypt

²National Institute of Fisheries and Oceanography, Cairo, Egypt

³Salah Ibrahim Fish Hatchery, Kafr El-Sheikh, Egypt

Abstract: A preliminary study (120 days) was conducted on African catfish (initial body weight 90 g) to evaluate the beneficial effects of a new patent local probiotic (T-Protophyt 2000) when added to their diet (25% crude protein) at graded levels (0, 1, 2 and 3 g kg⁻¹ diet). The diet containing 1 g kg⁻¹ (T₂) reflected the best growth and feed utilization parameters. Increasing the probiotic level increased fish carcass protein, fat and energy contents, as well as RBCs, WBCs, platelets and A/G ratio but decreased blood proteins. Also, T₂ treatment led to improvement of most histometric characteristics of the dorsal muscles of African catfish compared with the control (T₁) and other treatments (T₃ and T₄). The bacterial activity of this probiotic was tested *in vitro* against nine of pathogenic strains of Gram-negative bacteria (*Aeromonas hydrophilla*, *Pseudomonas aeruginosa*, *Pseudomonas fluorescent*, *Vibrio* sp., *Klebsiella* sp., *Shigella* sp., *Salmonella* sp., *Proteus* sp. and *Escherichia coli*) at two concentrations (120 and 240 µg) compared with oxytetracycline (OTC 30 and 60 µg). The results showed positive effect of the probiotic at the two concentrations against all the tested bacteria.

Key words: Probiotic, catfish, growth, blood, histometric parameters, pathogenic bacteria

INTRODUCTION

Dietary live yeast improved ($p < 0.01$) fish body weight, reduced muscular fat and serum triglycerides and cholesterol but increased RBC's, Hb, PCV and serum glucose. So, dietary live yeast may improve growth performance and hematological picture in fish (Kobeisy and Hussein, 1995). Moreover, Abdelhamid *et al.* (2000) reported significant positive effects of the combination of dried live yeast and lacto-sacc on tilapia growth, feed conversion and nutrients utilization.

The highest level of these separate or combined additives (20 g kg⁻¹) was the best for improving fish body weight and feed conversion. Also, Magouz *et al.* (2002) and Abou Zied *et al.* (2003) came to the same conclusion that dietary supplementation of lacto-sacc (0.1% of the diet) produced high growth rate, survival rate and feed and protein utilization of Nile tilapia. Lacto-sacc had also positive effect on economical efficiency of tilapia production. However, Olvera-Novoa *et al.* (2002) reported that fish fed 25-30% (of the dietary

Corresponding Author: Dr. Ahmed Ismail Mehrim, Department of Animal Production,
Faculty of Agriculture, Al-Mansoura University, Al-Mansoura,
P.O. Box 35516, Egypt

protein) yeast diets showed the best growth performance, feed conversion, protein efficiency ratio, nitrogen utilization, incidence cost and profit index.

Additionally, Khattab *et al.* (2004) found that Biogen^(R) (0.1% of the diet), as a feed supplement, had improved growth performance, feed conversion, chemical composition of the whole fish body as well as the blood profile of Nile tilapia fish. Recently, Allam (2007) found that Pronifer^(R) (lactic acid bacteria and its fermentation metabolites) as feed additive at 3% level improved tilapia growth and their blood profile. It reduced fish fat content and alleviated the hazard effects of *A. hydrophila* on mortality rate. The African catfish *Clarias gariepinus* is distributed throughout Africa. It is of growing economic value in the African aquaculture industry (Goda *et al.*, 2007; Osman *et al.*, 2007; Abdelhamid, 2009a).

Probiotics are pure cultures of one or more living microorganisms given in feed that proliferates in the host gastrointestinal (GI) tract. They ensure that the host maintains a beneficial microbial population in the GI tract (Linge, 2005). They confer a healthy effect on the host as significant microbial food supplements in the field of prophylaxis (Geovanny *et al.*, 2007). The research of probiotics for aquatic animals is increasing with the demand for environment-friendly aquaculture. Some probiotics were designed to treat the rearing medium, like biocontrol when the treatment is antagonistic to pathogens or bioremediation when water quality is improved. Most probiotics have been undertaken by isolating and selecting strains from aquatic environment (Gatesoupe, 1999). Also, probiotics have found use in aquaculture as a means of disease control, supplementing or even in some cases replacing the use of antimicrobial compounds (Irianto and Austin, 2002; Sahu *et al.*, 2008). Since, the use of expensive chemotherapeutants for controlling diseases have been widely criticized for their negative impacts (Sahu *et al.*, 2008). Therefore, the objectives of the present study were to evaluate effects of graded levels of a new-local probiotic on African catfish concerning their growth performance, feed utilization, carcass composition, histometric characteristics of the dorsal muscles, blood profile as well as effect of this probiotic on some pathogenic bacteria of fish.

MATERIALS AND METHODS

Experimental Management

A field study was conducted to evaluate the effects of dietary inclusion of graded levels of a newly local produced probiotic (T-Protyphyt 2000: It is a local product with a patent No. 23593. It consists of 15% zinc salts, 10% inorganic phosphorus, 5% dried fermentation products of *Aspergillus oryzae* growth and starch as carrier up to 1 kg. Each gram of this product contains 100 unit of phytase, 75 unit of protease, 25 unit of lipase and 15 unit of amylase) on African catfish (*Clarias gariepinus*). For this reason, 4 net Hapas (1×3×1.5 m diameters) were constructed and implanted in 1 Feddan (Egyptian area unit = 4200 m²) earthen pond in a private fish farm at Tolompat 7, Alriad, Kafr El-Sheikh Governorate, Egypt. The Hapas' net was 1 cm opens, from Tailand. The pond was irrigated via a pump from an agricultural drain. Eighty similar-size catfish were purchased from a private neighbor farm, regardless to their sex, with an average body weight of 90 g.

The feeding trial started on the 20th of July 2008 and ended on the 20th of November 2008 (120 days) using a commercial diet (25% crude protein, from Almorshedy for Trading and Development, Meet Ghamr-Dakhalia-Zagazig Road, Egypt). This commercial diet contained yellow corn, soybean meal (44%), wheat bran, fish meal (65%), corn gluten (60%), lime stone, common salt, dicalcium phosphate and molasses and had not less than 25% crude protein, 3% crude lipids, 3935 Kcal gross energy kg⁻¹ diet and not more than 5.30% crude

fiber, according to the manufacture's formula. Table 1 shows the proximate analysis of the basal diet which was carried out according to AOAC (1995). The diet was ground to add the tested probiotic (at levels of 0, 1, 2 and 3 g kg⁻¹ diet, referred to treatments No. T₁, T₂, T₃ and T₄, respectively). Molasses at 5% of each diet was used to spread the probiotic and then all diets were repelleted. The tested diets were offered once daily (10 am) at 5% of the fish biomass at each Hapa. The feed quantity was adjusted periodically according to the actual body weight changes.

Fish Performance and Quality

Measurements used for the evaluation included fish weight and length and water quality parameters (i.e., temperature using centigrade thermometer, salinity using conductivity TDS meter model 470, pH using pH meter model 340, dissolved oxygen using oxygen meter model 970, all instruments were from Jenway-England).

At the start and at the end of the experiment, fish samples were collected and kept frozen till the proximate analysis of the whole fish body according to AOAC (1995). Their gross energy contents were calculated according to NRC (1993).

At the end of the experiment some fishes from each treatment were sacrificed and fish dorsal muscles were sampled. Samples were fixed in 10% neutralized formalin solution to histometric examination according to Pearse (1968).

Blood Parameters

At the end of the experiment blood samples were collected from the fish caudal peduncle of the different groups. Adequate amounts of whole blood in small plastic vials containing heparin were used for the determination of hemoglobin (Hb) using commercial colorimetric kits (Diamond Diagnostic, Egypt). Also, total erythrocytes (RBCs), platelets and total leucocytes (WBCs) were counted according to Dacie and Lewis (1995) on an Ao Bright-Line Haemocytometer model (Neubauer improved, Precicolor HBG, Germany). Other blood samples were collected and transferred for centrifugation at 3500 rpm for 15 min to obtain blood plasma for determination of plasma total protein according to Gornall *et al.* (1949), albumin according to Weichsebum (1946) and globulin by difference according to Doumas and Biggs (1972).

Bacterial Strains

Nine of Gram-negative bacteria, namely *Aeromonas hydrophilla*, *Pseudomonas aeruginosa*, *Pseudomonas fluorescens* and *Vibrio* sp., in addition to the *Enterobacteriaceae* (*Klebsiella*, *Shigella*, *Salmonella*, *Proteus* and *Escherichia coli*) were used in this study, these bacteria were isolated from Nile tilapia fish and identified by specific media and biochemical testes, except *Aeromonas hydrophilla* that was identified by primed Polymerase Chain Reaction (PCR).

Assessment of Antibacterial Activity of the Probiotic

Powder samples of probiotic and (OTC) antibiotic (because of its wide antibacterial spectrum and high potency, OCT is the most commonly used antibiotic against various diseases caused by Gram-negative and Gram-positive bacteria in fish farming) were suspended in sterile water and used at two concentrations each (120 and 240 µg of probiotic and 30 and 60 µg of OTC). Four wells were punched with a cork borer (6 mm in diameter) in plates of nutrient agar (NA) freshly seeded with 0.1 mL of 24 h old of each tested bacterial cultures. Different concentrations of probiotic and antibiotic were put into the wells, left

1 h to allow diffusion, plates were incubated for 24 h at 37°C. The diameter of clear zones surrounding the wells were measured and recorded expressing the antibacterial activity.

Statistical Analysis

The obtained data were statistically analyzed using SAS (1996) procedures for personal computer. When F-test was positive, least significant difference (Duncan, 1955) was calculated for the comparisons among means.

RESULTS

Experimental Diet

The chemical analysis of the basal diet used in the present experiment is given in Table 1. Calculated gross energy based on factors of 5.65, 9.45 and 4.11 Kcal g⁻¹ protein, fat and carbohydrate, respectively according to NRC (1993) was 395.5 Kcal 100 g⁻¹, protein/energy (P/E) ratio was 58.76 mg Kcal⁻¹.

Water Quality Criteria

Water parameters were measured twice daily (10 am and 10 pm). There were no significant differences among Hapas' water throughout the experimental period because all Hapas were in the same pond. Therefore, the means of the whole period are given in the following Table 2.

Growth Performance

Growth performance parameters shows in Table 3 shows that T₂ (1 g T-Protophyt 2000 kg⁻¹ diet) was the best among various dietary treatments concerning body weight gain, daily body weight gain, RGR, SGR and condition factor as well as PER and PPV (even feed conversion ratio which not given in the Table 3). Yet, all data in this Table 3 are lower than those in literature (El-Haroun, 2007) probably for low dietary protein and fat determined (23.24 and 2.87%, respectively) as well as for poor experimental rearing conditions, e.g., stocking rate, feeding rate and frequency and using Hapas which restricted the fish growth and negatively affected feed conversion and nutrients utilization. Therefore, the evaluation will continue in a serial paper under other good ambient conditions. T₂ gave significantly ($p \leq 0.05$) the highest final body weight and condition factor. However, all supplemental diets with the tested probiotic (T₂, T₃ and T₄) reflected better results than the control (T₁). T₄ did not differ significantly ($p \geq 0.05$) than either T₂ or T₃.

Table 1: Proximate chemical analysis of the basal diet (% as fed)

Composition	Values
Moisture	9.01
Crude protein	23.24
Ether extract	2.87
Total carbohydrates	57.67
Ash	7.21

Table 2: Water quality parameters (Means±SE) of the whole experimental period, regardless to the treatments

Parameters	Values
Temperature (°C)	20.61±1.078
Salinity (‰)	2.778±0.038
pH value	8.370±0.186
Dissolved oxygen (mg L ⁻¹)	7.475±0.225

Table 3: Growth performance of catfish after 120 days of feeding the experimental diets (Means±SE)

Items	Treatment			
	T ₁	T ₂	T ₃	T ₄
Final body weight (g)	144	162	150	148
Body weight gain (g)	54	72	60.0	58
Daily weight gain (mg)	450	600	500	483
Relative growth rate* (%)	0.351±0.046 ^{NS}	1.131±0.242 ^{NS}	1.017±0.135 ^{NS}	0.793±0.083 ^{NS}
Specific growth rate** (% day ⁻¹)	0.250±0.029 ^{NS}	0.619±0.095 ^{NS}	0.582±0.054 ^{NS}	0.485±0.040 ^{NS}
Final body length (cm)	26.67±0.876 ^b	29.90±0.850 ^{ab}	30.30±0.569 ^a	29.33±0.333 ^a
Condition factor	0.647±0.047 ^c	0.710±0.023 ^a	0.651±0.006 ^b	0.653±0.024 ^{abc}
Survival (%)	100	100	100	100
Protein efficiency ratio***	0.314	0.387	0.342	0.328
Protein productive value (% ****)	5.300	7.238	6.360	6.322
Energy utilization (%*****)	3.443	4.282	3.887	4.294

a-c: Means in the same row with different superscripts are significantly ($p < 0.05$) different; NS: Not significant at $p \geq 0.05$; *RGR: (Final body weight-Initial body weight)/Initial body weight; **SGR: (ln final weight-ln initial weight)×100/experimental period; ***PER: Body weight gain/consumed feed protein; ****PPV: Retained protein×100/consumed feed protein; *****EU: Retained energy×100/consumed feed energy

Carcass Composition

Proximate chemical analysis of the whole fish body at the start and at the end of the 120 day experimental period is summarized in Table 4. These data indicated that moisture content was higher and ether extract as well as energy content were lower at start than at the experimental end; otherwise, no remarkable changes were recorded. Concerning dietary treatments, there were slight increases in crude protein and ether extract percentages but lower ash content due to the dietary inclusion of T-Protophyt 2000.

Table 4: Proximate chemical analysis (% fresh basis) of whole fish body at the start and at the end of the feeding period

Composition	Treatment				
	At start	T ₁	T ₂	T ₃	T ₄
Moisture	75.21	71.28	71.62	72.02	70.60
Crude protein	18.57	17.93	18.36	18.58	18.86
Ether extract	0.76	4.10	4.48	4.21	5.13
Ash	4.72	5.60	4.66	4.63	5.32
Growth energy (KCal 100 g ⁻¹)	112.10	140.00	146.00	144.80	155.00

Histometric Examination of Fish Dorsal Muscles

There were no significant ($p \geq 0.05$) differences of all histometric parameters (smallest diameter (µm), mean diameter (µm), smallest/largest ratio, intensity of muscular bundles mm⁻², the percentage of muscular bundles area mm⁻² and the percentage of connective tissue mm⁻²) of African catfish dorsal muscles among all treatments. However, fish fed diet supplemented with commercial probiotic T-Protophyt 2000 at level of 1 g kg⁻¹ diet (T₂) realized slight improvement of these histometric characteristics of fish dorsal muscles compared with the control (T₁) or other treatments (T₃ and T₄). It is of interest to note that, T₂ treatment realized the best growth performance (Table 3) and carcass composition (Table 4) of fish compared with the control and other treatments. This means that supplementation of commercial probiotic T-Protophyt 2000 at level of 0.1% to fish diets led to improvement of most histometric characteristics of the dorsal muscles of African catfish (Table 5, Fig. 1a-d).

Blood Profile

Data of hematological and biochemical parameters are given in Table 6, which clears that there were no significant ($p \geq 0.05$) effects of the tested probiotic on hemoglobin (Hb) content and white blood cells count (WBCs) comparing with the control. Yet, there were significant




Fig. 1: Cross-section of muscular bundles and interstitial connective tissue of the dorsal muscles of catfish in the (a) 1st, (b) 2nd, (c) 3rd and (d) 4th treatments respectively. (X 400, H and E stains)

Table 5: Effect of dietary supplementation of T-Protyph 2000 on histometric characteristics of dorsal muscles of catfish (Means±SE)

Parameters	Treatment			
	T ₁	T ₂	T ₃	T ₄
Smallest diameter (µm)	30.40±2.03 ^{NS}	38.40±1.59 ^{NS}	31.20±2.57 ^{NS}	34.00±1.78 ^{NS}
Largest diameter (µm)	36.40±1.72 ^{NS}	43.60±2.03 ^{NS}	40.00±3.09 ^{NS}	41.20±1.85 ^{NS}
Mean diameter (µm)	33.40±1.50 ^{NS}	41.00±1.64 ^{NS}	35.60±2.69 ^{NS}	37.60±1.63 ^{NS}
Smallest/Largest ratio	0.84±0.06 ^{NS}	0.88±0.03 ^{NS}	0.78±0.04 ^{NS}	0.83±0.03 ^{NS}
Intensity of muscular bundles (mm ⁻²)	586.67±92.66 ^{NS}	586.67±92.66 ^{NS}	679.33±92.66 ^{NS}	679.33±92.66 ^{NS}
% of muscular bundles area (* mm ⁻²)	58.77±19.07 ^{NS}	69.63±4.80 ^{NS}	57.66±8.67 ^{NS}	69.63±4.80 ^{NS}
% of connective tissue (** mm ⁻²)	41.23±19.07 ^{NS}	30.37±4.80 ^{NS}	42.34±8.67 ^{NS}	30.37±4.80 ^{NS}

^{NS}: Not significant at p≥0.05; * % of muscular bundles area mm⁻² = ([3.14 X (mean diameter/2)²]×Intensity of muscular bundles mm⁻²)×100, whereas: the muscular bundles were considered in approximately circular shape; ** % of connective tissue mm⁻² = (1- muscular bundles area, mm⁻²)×100

Table 6: Hematological and biochemical analysis of catfish blood after 120 days experimental feeding (Means±SE)

Items	Treatment			
	T ₁	T ₂	T ₃	T ₄
Hemoglobin, g dL ⁻¹	6.80±0.40 ^{NS}	6.60±0.10 ^{NS}	6.95±0.05 ^{NS}	7.50±0.20 ^{NS}
RBCs, x 10 ⁶ µL ⁻¹	2.60±0.10 ^{bc}	2.10±0.10 ^c	2.75±0.15 ^a	2.80±0.20 ^{ab}
WBCs, x 10 ³ µL ⁻¹	32.50±2.50 ^{NS}	40.00±5.00 ^{NS}	42.50±7.50 ^{NS}	47.50±7.50 ^{NS}
Platelets µL ⁻¹	52.50±7.50 ^b	85.00±10.0 ^{ab}	67.50±7.50 ^{ab}	100.00±5.00 ^a
Total protein, g dL ⁻¹	6.15±0.15 ^a	5.60±0.10 ^b	4.20±0.30 ^c	5.70±0.20 ^{abc}
Albumin (A), g dL ⁻¹	1.70±0.10 ^b	1.65±0.05 ^c	1.45±0.05 ^d	1.85±0.05 ^{cd}
Globulin (G), g dL ⁻¹	4.45±0.05 ^a	3.90±0.20 ^{ab}	2.75±0.25 ^b	3.85±0.15 ^{ab}
A/G ratio*	0.382±0.018 ^b	0.425±0.035 ^a	0.530±0.030 ^a	0.481±0.005 ^a

^{a-d}: Means in the same row with different superscripts are significantly (p<0.05) different; RBCs: Red blood cells (Erythrocytes); WBCs: White blood cells (Leucocytes); *A/G ratio: Albumin/Globulin

(p<0.05) positive effects on platelets count and A/G ratio and negative effects on total protein and globulin contents; otherwise, no clear trend was recorded for the effect of

probiotic inclusion levels. T₄ did not differ significantly ($p \geq 0.05$) with T₃ in most parameters, except A/G ratio, but differ ($p \leq 0.05$) with T₂ in both red blood cells count (RBCs) and A/G ratio. T₃ did not differ significantly ($p \geq 0.05$) with T₂ in platelets count and A/G ratio. The increased A/G ratio is related to the decrease in globulin values.

Antagonism to Pathogens

The results in Table 7 and Fig. 2 A-D showed the positive effect of the probiotic at the two concentrations against all the tested bacteria and showed nearly no clear difference between the two concentrations of probiotic. Also, it has a similar effect of the antibiotic (OTC) especially with the pathogenic bacteria, *Aeromonas* and *Pseudomonas*, which showed sensitivity towards probiotic, while the *Vibrio* sp., showed resistance to OTC at the two concentration (Fig. 2D).

Table 7: The antibacterial activity of aqueous probiotic compared with OTC

Tested bacterial strains	Inhibition zone (mm) of probiotic		Inhibition zone (mm) of OTC	
	120 ----- (µg) -----	240	30 ----- (µg) -----	60
<i>A. hydrophilla</i>	32	37	36	36
<i>P. aeruginose</i>	35	38	30	35
<i>P. fluorescens</i>	45	44	42	42
<i>Vibrio</i> sp.	27	36	-	-
<i>Salmonella</i> sp.	30	30	35	42
<i>Shigella</i> sp.	35	35	37	38
<i>Klebsiella</i> sp.	21	20	30	30
<i>Proteus</i> sp.	20	22	30	40
<i>E. coli</i>	34	36	38	38




Fig. 2: (a-d) The inhibition zones of some bacteria by the action of probiotic compared with OTC. (A) *A. hydrophilla*, (B) *P. aeruginose*, (C) *P. fluorescens*, (D) *Vibrio* sp. 1 = 120 µg probiotic, 2 = 240 µg probiotic, 3 = 30 µg OTC, 4 = 60 µg OTC

DISCUSSION

The values of water parameters are within the acceptable ranges recommended for pisciculture (Abdelhamid, 1996, 2009b; Abdelhakim *et al.*, 2002). However, the optimum growth of African catfish requires 28-30°C, <5 ppt salinity, <15 mg L⁻¹ dissolved oxygen, 6.5-9.0 pH and 50-100 mg L⁻¹ hardness in the rearing water (Chapman, 2000).

Craig and Helfric (2002) reported that protein levels in aquaculture feeds generally have an average of 28-32% for catfish. Protein requirements are lower for omnivorous fish and for larger fish than carnivorous and smaller ones. Feeding rate affects also fish requirements of protein (Jauncey, 1998). However, Machiels and Henken (1985) reported 40% crude protein, 19.2 KJ DE g⁻¹ and 13 mg protein/KJ as optimal requirements for *C. gariepinus* (40-120 g). Recently, Eid (2007) recommended a diet containing 25% protein, 6% fat and 72 mg protein Kcal⁻¹ for adult catfish. Also, El-Gendy (2009) found that the best dietary crude protein and fat levels were 35.9 and 11.7% (465.88 Kcal 100 g⁻¹), respectively and 77.1 P/GE ratio for the African catfish fingerlings (13 g initial body weight).

Robinson *et al.* (2009) reported that even though catfish have been cultured for many years, there is still considerable variation in feeding practices on commercial catfish farms. Catfish are generally fed once daily to what is commonly called saturation. Catfish (27 g) feed is generally recommended to contain 28-32% protein, starting with a 32% protein feed in spring and change to a 28% feed as the temperature increases.

Abdelhamid *et al.* (1996) and Attia *et al.* (2007) confirmed the economical-environmental benefits of using the dietary supplemental microbial-phytase. It increases (p<0.05) body weight, feed efficiency, carcass % and serum contents of P, Ca, Mg and Zn. Also, El-Dakar and Gohar (2004) found that growth and survival of *Peneaus japonicus* post larvae fed probiotic diet was higher than those fed the basal diet. The *in vitro* study revealed that the probiotic (*Bacillus subtilis*) decreased the proteolyses activity of the bacterial pathogens (*Aeromonas hydrophila*, *Edwardsiella tarda* and *Vibrio proteolyticus*). *Bacillus subtilis* had a positive effect against the pathogens and on reducing the antibiotic susceptibility when presented in culture water or in feed of shrimp. Moreover, Salem *et al.* (2004) reported that bacterial and yeast probiotics (*Bacillus subtilis* and *Saccharomyces cerevisiae*) improve the activity of lactic acid bacteria which could inhibit the pathogenic bacteria (*E. coli*). Castillo (2008) also concluded that certain *Bacillus* strains act as probiotic bacteria and block the communication system of Enter pathogens such as *Yersinia* and *Salmonella* sp. Yet, Hidalgo *et al.* (2006) concluded that no significant effects on growth and survival were found following the addition of *Bacillus toyaoi*, T and *B. cereus*, E as probiotics to dentex diets.

However, some lactic acid bacteria isolated from the gastrointestinal tract of fish can act as probiotics. These candidates are able to colonize the gut and act antagonistic against Gram negative fish pathogens. These harmless bacteriocin-producing strains may reduce the need to use antibiotics in future aquaculture (Ringo and Gastesoupe, 1998). Generally, the probiotics actively inhibit the colonization of potential pathogens in the digestive tract by antibiosis or by competition for nutrients and/or space, alteration of microbial metabolism and/or by the stimulation of host immunity. Probiotics may stimulate appetite and improve nutrition by the production of vitamins, detoxification of compounds in the diet and by breakdown of indigestible components (Irianto and Austin, 2002).

Additionally, Abd El-Rahman and El-Bana (2006) used *Micrococcus luteus* as a bacterial probiotic which presented *in vitro* and *in vivo* antagonistic effects against the pathogenic bacteria *Aeromonas hydrophila*. The inhibition zone to *A. hydrophila* was 40 mm in diameter due to *M. luteus*. The dietary inclusion of this probiotic improved significantly the final

weight, weight gain, specific growth rate, feed conversion, protein efficiency ratio erythrocytic counts, hemoglobin content and survival rate of *O. niloticus* fish. Rollo *et al.* (2006) reported an improvement in tolerance to acute stress of sea bream fry fed with probiotics. Also, Taoka *et al.* (2006) indicated that probiotics treatment is promising as an alternative method to antibiotics for disease prevention in aquaculture. Additionally, Attalla (2007) found that supplementation of dietary bacteria or yeast produced significantly better weight gain, specific growth rate and feed utilization in tilapia cultivation.

Biobuds[®] and yeast or Biogen[®] as probiotics led to improving the growth performance, feed conversion, protein efficiency ratio and feed costs for tilapia fingerlings (Mohamed, 2007; Mohamed *et al.*, 2007). Moreover, El-Ashram *et al.* (2008) concluded that, super Biobuds[®] can improve body gain, survival and enhance resistance to challenge infection. Yet, Abdelhamid and Elkatan (2006) found that dietary supplementation of Biobuds[®] slightly improved body weight gain but reduced the survival rate of tilapia fingerlings. El-Haroun *et al.* (2006) and El-Haroun (2007) reported that Biogen[®] dietary supplementation improved growth performance and feed utilization, carcass protein and fat percentages as well as economical profit in Nile tilapia and catfish culture, respectively. Wongsu and Werukhamkul (2008) came to the same conclusion, since they found that catfish fed diets containing probiotics plus phytase enzyme showed 35% higher weight gain and better feed conversion by more than 25% in comparison to catfish fed control diets in a three-month trial.

However, during the past two decades, the use of probiotics as an alternative to the use of antibiotics has shown to be promising in aquaculture, particularly in fish and shellfish larviculture (Tinh *et al.*, 2008). Recently, Aly *et al.* (2008a) found that some *Bacillus* and *Citrobacter* strains isolated from Nile tilapia (*B. pumilus*, *B. firmus* and *C. freundii*) showed inhibitory effects against *A. hydrophila*. Also, Aly *et al.* (2008b) reported that the probiotic activity of two bacteria (*Bacillus subtilis* and *Lactobacillus acidophilus*) was evaluated by its effect on the immune response of Nile tilapia (*Oreochromis niloticus*), beside its protective effect against challenge infection. Furthermore, their *in-vitro* inhibitory activity was evaluated. The *in vitro* antimicrobial assay showed that *Bacillus subtilis* and *Lactobacillus acidophilus* inhibited the growth of *A. hydrophila*. The *B. subtilis* inhibited the development of *P. fluorescens* while *L. acidophilus* inhibited the growth of *Strept. iniae*. The *B. subtilis* and *L. acidophilus* proved harmless when injected in the *O. niloticus*. The feed, containing a mixture of *B. subtilis* and *L. acidophilus* or *B. subtilis* alone, showed significantly greater numbers of viable cells than feed containing *L. acidophilus* only after 1, 2, 3 and 4 weeks of storage at 4 and 25°C. The survival rate and the body-weight gain were significantly increased in the fish given *B. subtilis* and *L. acidophilus* for one and two months after application. The hematocrit values showed a significant increase in the group that received the mixture of *B. subtilis* and *L. acidophilus* compared with the control group. The nitroblue tetrazolium (NBT) assay, neutrophil adherence and lysozyme activity, showed a significant increase in all the probiotic-treated groups after 1 and 2 months of feeding, when compared with the untreated control group. The serum bactericidal activity was high in the group that was given a mixture of the two bacteria.

Also, Marzouk *et al.* (2008) reported that probiotics (*B. subtilis* and *Saccharomyces cerevisiae*) revealed significant improvement in growth parameters and showed failure in re-isolation of some pathogens. Varley (2008) cited also that probiotics show real benefits in the synergistic effects with the beneficial bacteria in making inroads into improving gut health. So, probiotics may improve the growth performance and immune response of fish (Wang *et al.*, 2008).

The increased count of WBCs may be caused by protein resorption (Merck, 1976). Hypoproteinemia may be due to protein loss, decreased albumin, or to increased globulin

(Merck, 1974). An insufficient amount of protein in the diet may lead to a low total protein of blood. This is the case in starvation. In other cases, although adequate protein is being taken in the feed, absorption from the alimentary tract may be defective. This may occur in vitamin deficiencies particularly of the B-vitamins (Varley, 1978).

In the field of physical structure of tilapia muscles, Abdelhamid *et al.* (2004) found that probiotics (Betafin[®] and Biopolym[®]) not only increased body weight, growth rates and total productivity, but also improved muscular protein percentage, radius of the muscular bundles, total surface area occupied by the muscular bundles mm⁻² (least thickness of connective tissues between muscular bundles and thickness of skin and subcutaneous layer) and net return.

CONCLUSION

In conclusion, this preliminary study revealed that this probiotic under study is useful for enhancing fish growth, feed and nutrients utilization, fish chemical composition and muscular structure, besides fish resistance for pathogenic bacteria, i.e. it may be useful also from the economic point of view. Yet, it is to recommend also more research work on this probiotic under various conditions, e.g., different fish species, initial weight, stocking rates, feeding rates, dietary protein levels and sources, etc.

REFERENCES

- Abdelhakim, N.F., M.N. Baker and M.A. Soltan, 2002. Aquatic Environment for Fish Culture. Faber and Faber, Cairo.
- Abdelhamid, A.M., 1996. Field and Laboratorial Analysis in Animal Production. 1st Edn., Dar Alnashr for Universities, Cairo, Egypt.
- Abdelhamid, A.M., 2009a. Recent Trends in Fish Culture. New Universal Office, Alexandria.
- Abdelhamid, A.M., 2009b. Fundamentals of Fish Production and Culture. New Universal Office, Alexandria.
- Abdelhamid, A.M. and M.S.A. Elkatan, 2006. A study on Nile tilapia fingerlings during wintering using dietary addition of Bio-Buds[®]-2X. *J. Agric. Sci. Mansoura Univ. Egypt*, 31: 5705-5711.
- Abdelhamid, A.M., A.E. Abd El-Khalek, M.A.A. Mostafa, S.A.A. Gomaah and F.F. Khalil, 2004. Effect of using Betafin and/or Biopolym as natural additives in producing Nile tilapia fish in poly-culture semi-intensive system in earthen ponds. *J. Agric. Sci. Mansoura Univ.*, 29: 3149-3162.
- Abdelhamid, A.M., F.F.M. Khalil and M.A.A. Seden, 2000. Possibility of using dried live yeast and lacto-sacc in Nile tilapia fingerlings diets. *J. Agric. Sci. Mansoura Univ.*, 25: 4905-4911.
- Abdelhamid, A.M., F.I. El-Hawary, M.M. El-Sawah and W.I. Saber, 1996. Benefits of dietary supplemental microbial-phytase in the broilers production. Proceedings of Conference Food Borne Contamination and Egyptian's Health, Nov. 26-27, Mansoura Univ., Egypt, pp: 345-353.
- Abd El-Rahman, A.M.M. and L.F.A. El-Bana, 2006. Using of *Micrococcus luteus* as a probiotic bacteria among cultured *Oreochromis niloticus*. *SCVMJ*, 10: 73-82.
- Abou-Zied, R.M., A.M. Abd El-Maksoud and A.A. Ali, 2003. Effect of virginiomycin or lacto-sacc on the growth performance of Nile tilapia fish during the nursing period. *Egypt. J. Nutr. Feeds*, 6: 481-489.

- Allam, H.Y.H., 2007. Physiological effects of some additives on growth, blood constituents and immunity in Nile tilapia (*Oreochromis niloticus*). Ph.D. Thesis, Fac. of Agric., Assiut Univ., Egypt.
- Aly, S.M., A.M. Abd-El-Rahman, G. John and M.F. Mohamed, 2008a. Characterization of some bacteria isolated from *Oreochromis niloticus* and their potential use as probiotics. *Aquaculture*, 277: 1-6.
- Aly, S.M., Y.A. Ahmed, A.A. Ghareeb and M.F. Mohamed, 2008b. Studies on *Bacillus subtilis* and *Lactobacillus acidophilus*, as potential probiotics, on the immune response and resistance of tilapia nilotica (*Oreochromis niloticus*) to challenge infections. *Fish Shellfish Immunol.*, 25: 128-136.
- AOAC., 1995. Association of Official Analytical Chemists. Official Methods of Analysis. 16th Edn., AOAC., Arlington, Virginia, USA.
- Attalla, R.F., 2007. The use of some microorganisms species as growth promoters for *Sarotherodon galilacus*. *Pak. J. Oceanography*, 3: 25-36.
- Attia, F.A.M., H.S. Abd Elhalim and K.A. Ibrahim, 2007. Improving phytate phosphorus utilization by adding phytase to low protein, phosphorus and calcium corn-soybean broiler diets. *Agric. Res. J. Suez Canal Univ. Egypt*, 7: 1-10.
- Castillo, M., 2008. Blocking pathogens with *Bacillus* strains. *Feed-Mix*, 16: 19-19.
- Chapman, F.A., 2000. Farm-Raised Channel Catfish. Department of Fisheries and Aquatic Sciences, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville.
- Craig, S. and L.A. Helfric, 2002. Understanding Fish Nutrition, Feeds, and Feeding. Department of Fisheries and Wildlife Sciences, College of Vet. Med., Pub. No. 420-256, Virginia State, Petersburg.
- Dacie, J.V. and S.M. Lewis, 1995. Practical Haematology. 8th Edn., Churchill Livingstone, Edinburgh, Scotland.
- Doumas, B.T. and H.G. Biggs, 1972. Determination of Serum Albumin. In: Standard Methods of Clinical Chemistry, Cooper, G.A. (Ed.). Vol. 7, Academy Press, New York.
- Duncan, D.B., 1955. Multiple range and multiple F-tests. *Biometrics*, 11: 1-42.
- Eid, A.M.S., 2007. Nutritional Requirements for Fish. Academy of Scientific Research and Technology, Cairo, Egypt.
- El-Ashram, A.M.M., M.F. Mohammed and S.M. Aly, 2008. Effect of Biobuds as a commercial probiotic product in cultured tilapia. Proceedings of 8th International Symposium on Tilapia in Aquaculture, Oct. 12-14, Cairo, Egypt, pp: 1089-1096.
- El-Dakar, A.Y. and Y.M. Gohar, 2004. Use of *Bacillus subtilis* in microparticulate diets for producing biosecure *Peneaus japonicus* postlarvae. *J. Agric. Sci. Mansoura Univ. Egypt*, 29: 6853-6871.
- El-Gendy, E.H.I., 2009. Effect of protein and oil levels on growth performance of catfish (*Clarias gariepinus*). M.Sc. Thesis, Faculty of Agriculture, Suez Canal University, Ismailia, Egypt.
- EL-Haroun, E.R., 2007. Improved growth rate and feed utilization in farmed African catfish *Clarias gariepinus* (Burchell 1822) through a growth promoter Biogen® supplementation. *J. Fish. Aquatic Sci.*, 2: 319-327.
- EL-Haroun, E.R., A.M.A.S. Goda and M.A.K. Chowdhury, 2006. Effect of dietary probiotic Biogen® supplementation as a growth promoter on growth performance and feed utilization of Nile tilapia *Oreochromis niloticus* (L.). *Aquac. Res.*, 37: 1473-1480.
- Gatesoupe, F.J., 1999. Review, the use of probiotics in aquaculture. *Aquaculture*, 180: 147-165.

- Geovanny, G.R., B.J. Luis and M.A. Shen, 2007. Probiotics as control agents in aquaculture. *J. Ocean Univ. China*, 6: 76-79.
- Goda, A.M.A.S., E.R. El-Haroun and M.A.K. Chowdhury, 2007. Effect of totally or partially replacing of fish meal by alternative protein sources on growth of African catfish *Clarias gariepinus* (Burchell, 1822) reared in concrete tanks. *Aquacult. Res.*, 38: 279-287.
- Gornall, A.G., G.J. Bardawill and M.M. Parid, 1949. Method of determination protein in serum blood. *J. Biol. Chem.*, 177: 751-751.
- Hidalgo, M.C., A. Skalli, E. Abellan, M. Arizcun and G. Cardenete, 2006. Dietary intake of probiotics and maslinic acid in juvenile dentex (*Dentex dentex* L.): Effects on growth performance, survival and liver proteolytic activities. *Aquac. Nutr.*, 12: 256-266.
- Irianto, A. and B. Austin, 2002. Review, probiotics in aquaculture. *J. Fish Dis.*, 25: 633-642.
- Jauncey, K., 1998. *Tilapia Feeds and Feeding*. Pisces Press Ltd., Stirling, Scotland, pp: 240.
- Khattab, Y.A.E., A.M.E. Shalaby, S.M. Sharaf, H.I. El-Marakby and E.H. Rizkalla, 2004. The physiological changes and growth performance of the Nile tilapia *Oreochromis niloticus* after feeding with Biogen® as growth promoter. *Egypt, J. Aquat. Biol. Fish.*, 8: 145-158.
- Kobeisy, M.A. and S.Y. Hussein, 1995. Influence of dietary live yeast on growth performance and some blood constituents in *Oreochromis niloticus*. *Proceedings of 5th Science Conference Animal Nutrition*, Dec. 12-13, Ismailia, Egypt, pp: 417-425.
- Linge, P., 2005. The use of probiotics and yeast derivatives in India. *World Poultry*, 21: 12-15.
- Machiels, M.A.M. and A.M. Henken, 1985. Growth rate, feed utilization and energy metabolism of the African catfish, *Clarias gariepinus* (Burchell, 1822), as affected by dietary protein and energy content. *Aquaculture*, 44: 271-284.
- Magouz, F.I., M.K. Mohsen and A.H. Gooda, 2002. Effect of including some biological feed additives in the diet on growth performance and feed efficiency of Nile tilapia (*Oreochromis niloticus*). *Proceedings of 2nd Conference Foodborne Contamination and Egyptian's Health*, Apr. 23-24, Mansoura Univ., Egypt, pp: 329-339.
- Marzouk, M.S., M.M. Moustafa and N.M. Mohamed, 2008. The influence of some probiotics on the growth performance and intestinal microbial flora of *Oreochromis niloticus*. *Proceedings of the 8th International Symposium on Tilapia in Aquaculture*, 2008, Cairo, Egypt, pp: 1059-1071.
- Merck, E., 1974. *Klinisches Labor*. 12. Auflage Merck, Darmstadt.
- Merck, E., 1976. *Labordiagnostik in der Tiermedizin*. Diagnostica Merck, Darmstadt.
- Mohamed, K.A., 2007. Effect of using probiotic and yeast as growth promoters in commercial diet of tilapia (*Oreochromis niloticus*) fingerlings. *Agric. Res. J. Suez Canal Univ.*, 7: 41-47.
- Mohamed, K.A., B. Abdel Fattah and A.M.S. Eid, 2007. Evaluation of using some feed additives on growth performance and feed utilization of monosex Nile tilapia (*Oreochromis niloticus*) fingerlings. *Agric. Res. J. Suez Canal Univ.*, 7: 49-54.
- NRC (National Research Council), 1993. *Nutrient Requirements of Fish*. National Academy Press, Washington DC, USA., pp: 114.
- Olvera-Novoa, M.A., C.A. Martinez-Palacios and L. Olivera-Castillo, 2002. Utilization of torula yeast (*Candida utilis*) as a protein source in diets for tilapia (*Oreochromis mossambicus* Peters) fry. *Aquac. Nutr.*, 8: 257-264.
- Osman, A.G.M., S. Wuertz, I.A.A. Mekkawy, H. Exner and F. Kirschbaum, 2007. Embryo-toxic effects of lead nitrate of the African catfish *Clarias gariepinus* (Burchell, 1822). *J. Ichthyol.*, 23: 48-58.
- Pearse, G.W., 1968. Histological effects and diagnostic problems of mycotoxins in poultry. *Proceedings of the 25th West States Poultry Disease Conference, WSPDC'1968*, IEEE Xplore, London, pp: 76-79.

- Ringo, E. and F.J. Gatesoupe, 1998. Review, Lactic acid bacteria in fish: A review. *Aquaculture*, 160: 177-203.
- Robinson, E.H., M.H. Li and M.W. Brunson, 2009. Feeding Catfish in Commercial Ponds. Oklahoma State University Cooperative Extension Fact Sheets (SRAC-181), Division of Agricultural Sciences and Natural Resources, Oklahoma.
- Rollo, A., R. Sulpizio, M. Nardi, S. Silvi and C. Orpianesi *et al.*, 2006. Live microbial feed supplement in aquaculture for improvement of stress tolerance. *Fish Physiol. Biochem.*, 32: 167-177.
- Sahu, M.K., N.S. Swarnakumar, K. Sivakumar, T. Thangaradjou and L. Kannan, 2008. Probiotics in aquaculture: Importance and future perspectives. *Indian J. Microbiol.*, 48: 299-308.
- Salem, A.Z., M.M.I. El-Adawy, M.S.M. Salem and A.A. Hasan, 2004. Effect of probiotics feed as additives on the activity of isolated and characterized lactic acid intestinal bacteria against *Escherichia coli*-10 in sheep. *Egypt. J. Nutr. Feeds*, 7: 167-184.
- SAS, 1996. SAS/STAT Guide For Personal Computers. Version 6.12, SAS Institute Cary, NC.
- Taoka, Y., H. Meeda, T.Y. Jo, S.M. Kim, S.I. Park, T. Yoshikawa and T. Sakata, 2006. Use of live and dead probiotic cells in tilapia *Oreochromis niloticus*. *Fish. Sci.*, 72: 755-766.
- Tinh, N.T.N., K. Direckens, P. Sorgeloos, and P. Bossier, 2008. A review of the functionality of probiotics in the larviculture food chain. *Marine Biotechnol.*, 10: 1-12.
- Varley, H., 1978. *Practical Clinical Biochemistry*. 4th Edn., Arnold-Heinemann Publishers (India) Private Limited, New Delhi, pp: 28.
- Varley, M., 2008. Managing gut health without antibiotics. *Pig Progress*, 24: 27-28.
- Wang, Y.B., Z. Tian, J. Yao and W. Li, 2008. Effect of probiotics, *Enterococcus faecium*, on tilapia (*Oreochromis niloticus*) growth performance and immune response. *Aquaculture*, 277: 203-207.
- Weichsebum, T.E., 1946. Method for determination of albumin in serum blood. *Am. J. Clin. Pathol.*, 16: 40-40.
- Wongsa, P. and P. Werukhamkul, 2008. Improving growth performance and FCR in pangasius catfish with a combination of probiotics and phytase enzyme. *AQUA Culture Asia Pacific Magazine*, Sept./Oct., pp: 22-23.