



Journal of Environmental Science and Technology

ISSN 1994-7887

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>



Research Article

Observation of Changes in Sediment Nature by Environmental Impacts of Abu-Makhadeg Area, Red Sea, Egypt

¹Hashem A. Madkour, ²Abbas M. Mansour, ³Mohamed A. Sebak, ⁴Amani Badawai and ⁵Atef El-Taher

¹National Institute of Oceanography and Fisheries, Red Sea Branch, 84511 Hurghada, Egypt

²Department of Geology, Faculty of Science, Qena, South Valley University, 83523 Qena, Egypt

³Department of Physics, College of Science and Arts, Jof University, Al-Qurayat Branch, P.O. Box 756, Saudi Arabia

⁴National Institute of Oceanography and Fisheries, Alexandria Branch, Alexandria, Egypt

⁵Department of Physics, Faculty of Science, Azhar University, Assuit Branch, Egypt

Abstract

Background and Objectives: More than 15 tourist projects occupy the Red Sea coast of the Sharm Abu-Makhadeg area (Makadai area) Red Sea, Egypt. During the 90's, landscape destruction, landfills, shoreline change, dredging rocky tidal flat and brine water discharge were the main environmental impacts in Abu-Makhadeg area. The scope of the present work were monitoring of changes of sediment nature by environmental impacts of Sharm Abu-Makhadeg area and record all developmental activities during the past 20 years.

Materials and Methods: Grain size and geochemical analyses had been carried out on 32 surface marine sediment samples collected from Sharm Abu-Makhadeg area along the Egyptian Red Sea coast. **Results:** The sediments were characterized by the abundance of sand with minor amounts of mud and gravel. Generally, sand fraction is the main category among the three constituents. Geochemically, the factor controlling the carbonate content of studied sediments includes material supply of biogenic and terrigenous components. Organic matter recorded high values in some samples with increase depth. The high organic matter in surface marine sediments is primarily due to the high supply from primary productivity, terrestrial and reworked sediments. Texture was the main controlling factor for the organic matter enrichment. **Conclusion:** Grain size distribution and geochemical aspects of surface marine sediments of Sharm Abu-Makhadeg area using some statistical methods revealed the dominance of muddy sediments of most samples with high values of organic matter than background areas reflected the negative impact of the biogenic life and humans.

Key words: Monitoring, observation sediments, sand fraction, impacts, Sharm Abu-Makhadeg area, Red sea, Egypt, grain size distribution and geochemical aspects

Citation: Hashem A. Madkour, Abbas M. Mansour, Mohamed A. Sebak, Amani Badawai and Atef El-Taher, 2019. Observation of changes in sediment nature by environmental impacts of Abu-Makhadeg area, Red sea, Egypt. J. Environ. Sci. Technol., 12: 55-64.

Corresponding Author: Atef El-Taher, Department of Physics, Faculty of Science, Azhar University, Assuit Branch, Egypt Tel: 00201015687527

Copyright: © 2019 Hashem A. Madkour *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Sharm Abu-Makhadeg between Hurghada and Safaga along the Egyptian Red Sea coast occupies a small area with a beach about 4.5 km. Sharm Abu-Makhadeg area and its surrounding regions are a major recreational area sea bird and marine-mammal rookery and important commercial fishing ground¹. Now several of hotels and tourist villages are being built around the sharm. This tourism projects including what is working and the ones under construction and tourism projects total area of Sharm Abu-Makhadeg more than 50 tourist projects.

The rest of the beach is already sold for developers and destruction clearly appears in the land and offshore. For example because the rocky tidal flat surrounds the sharm, the needs for swimming pools or marinas entail dredging zones in-front of resorts during the nineties¹.

In the present work, 32 samples were collected of surface marine sediments from Sharm Abu Makhadeg area and to follow up on those natural variations sedimentary environmental influences after the passage of more than 15 years of study by Mansour¹.

Several investigations on the surface marine sediments were carried out on the Egyptian Red Sea coast²⁻²⁰. However, investigations including the impact of development activities along the Egyptian Red Sea coast are rare. The scope of the present work was observation of changes of sediment nature by environmental impacts of Sharm Abu-Makhadeg area and record all development activities during the past 20 years. The impact of development activities in Sharm Abu-Makhadeg area was measured by surveying and by analyzing the surface marine sediment²¹⁻²⁷. In the same manner, this paper deals with the changes that have occurred in marine sediments as a result of these activities through the mechanical analysis of sediment samples and the study of the geochemistry of these sediments.

MATERIALS AND METHODS

Study area: Sharm Abu Makhadeg area, the famous Egyptian coast of the Red Sea tourist areas and are called Madinat Makadi Tourist. Sharm Abu Makhadeg has a small and very narrow fringing reef occupying its northern border, a wide and fascinating coral reef covers its southern part (Fig. 1, 2). South of the Sharm, a crescent tidal flat area with a maximum width of 800 m is developed' decreasing southward to reach to ab. 75 m at Sharm El-Arab, with a few pools (lagoons) (Fig. 3a, b).

The inland geomorphology of Wadi Abu-Makhadeg and its accumulations of alluvial fans north and south, which Quaternary coral reef terraces occur. Vegetated coastal dunes and sabkhas occupy the lowland areas at the wadi mouth (Fig. 1, 2). Westwards, Miocene and younger sediments extend to the high Basement rocks mountains. Wadi Abu Makhadeg fan-out down slope of hills, being dry most of the time and is characterized by sporadic and abrupt fluvial activity. Quaternary sand and gravels cover the study area. Some Pliocene marine outcrops formed of inter-bedded sandstone, marls and clayey rocks, while the outcropping Miocene rocks are formed of clastic base with gypsum and carbonate beds¹. The majority of the area westward is covered by high rugged outcropping basement rocks, which mainly composed of granitoid rocks, volcanics and metavolcanics and gabbroic basic rocks. Generally, these outcropping basement rocks is traversed by N-E and N-W trending faults and fractures. Along these faults, the basement rocks are brecciated, sheared and milonitized.

Field works: In this study, 32 surface marine sediment samples have been collected from Sharm Abu-Makhadeg area (Fig. 1, 2), the location, depth and description of bottom characteristics of the collected samples are given in Table 1 and Fig. 2 and 3. Surface marine sediments samples collected from the study area represent three different environmental feature; beach, intertidal zone and offshore zone until 25 m water depth. The sampling was carried out by a grab sampler and Scuba diving. The later was used in areas rich with corals where grab sampler failed to collect samples. After anchoring the boat at each station, the sampler was lowered to the sea floor and left a few seconds before being pulled back to the surface. The sediment caught in the sampler or collected by Scuba diving was placed in labeled plastic bags and returned to the laboratory.

Laboratory methods and treatment of data: All samples were washed several times to remove soluble salts. Grain size analysis were performed using the sieving technique according to Folk and Ward²⁸. The resulting data were processed on a personal computer using the "BASIC" program "SEDPAC"²¹. The carbonate content and the organic matter of all sediment samples were determined in the National Institute of Oceanography and Fisheries, Red Sea Branch. All geochemical analyses were carried out in duplicates and the average of data was determined. Carbonate content was determined by treating the samples with (1 N HCL acid). The




Fig. 1: Study area along the Red Sea coast, Egypt (After authors)

insoluble residue remaining after acid washing was determined and the carbonate percentage was calculated. Determination of organic matter was made by sequential weight loss²⁹ at 550°C. All analysis of the sediment samples carried out of the laboratories of the National Institute of Oceanography and Fisheries, Red Sea Branch, Egypt in the period between March and August, 2018.

The obtained data of the granulometric and geochemical analyses were dealt statistically in order to exclude the characteristic parameters. The statistical treatment includes the average, correlation coefficient and cluster analysis.

Analyses are carried out on the data using the computer programs of the SPSS system available in National Institute of Oceanography and Fisheries, Red Sea Branch, Egypt.

RESULTS

Grain size analysis: The marine sediments in the study area were mainly composed of sand (68.85-96.8% with average 84.13%), mud constitutes (0.001-30.24% with average 10.97%) and gravel was very rare (0.14-30.11% with average 4.9%) (Table 2). Generally, the correlation coefficients between




Fig. 2: Distribution of surface marine sediment samples at Sharm Abu-Makhadeg area (After authors)




Fig. 3(a-d): General view of Abu-Makhadeg Bay: (a-b) Southern part of beach and intertidal area of Abu-Makhadeg area, (c) Nature of bottom sediments near the beach of the northern part of Abu-Makhadeg Bay and (d) Rehabilitation operation of coral reefs and marine organisms of Abu-Makhadeg Bay by this project (After authors)

sediments types, grain size parameters, carbonate content and total organic matter was weak (Table 3). With the exception of these relations gravel with Mean Size (MZ)

($r = -0.82$), gravel with carbonate content ($r = -0.52$), sand with mud ($r = -0.7$), sand with skewness (Sk_s) ($r = -0.56$), mud with mean size (MZ) ($r = 0.77$), mud with sorting (6_s) ($r = 0.51$), mud

Table 1: Sample location, depth and bottom facies at Sharm Abu-Makhadeg area

| Sa. No. | Position | | Depth (m) | Bottom facies |
|---------|----------------------|-----------------------|-----------|--------------------------------|
| | Latitude o / // N | Longitude o / // E | | |
| M1 | 26 59 02 | 33 54 53 | Beach | Coarse sand |
| M2 | 26 59 04 | 33 54 53 | 4 | Muddy sand with seagrass |
| M3 | 26 59 03 | 33 54 53 | 4 | Biogenic sand with seagrass |
| M4 | 26 59 02 | 33 54 48 | Beach | Muddy sand |
| M5 | 26 59 02 | 33 54 50 | 2 | Sandy gravel |
| M6 | 26 59 02 | 33 54 50 | 2.2 | Sand with seagrasses |
| M7 | 26 59 09 | 33 54 53 | 1.5 | Sand with seagrasses |
| M8 | 26 59 13 | 33 54 53 | 2.2 | Muddy sand with seagrass |
| M9 | 26 59 16 | 33 54 54 | 2.9 | Muddy sand with seagrass |
| M10 | 26 59 17 | 33 54 53 | 1.4 | Biogenic sand |
| M11 | 26 59 11 | 33 54 48 | 2.6 | Muddy sand |
| M12 | 26 59 08 | 33 54 47 | 2.1 | Muddy sand |
| M13 | 26 59 07 | 33 54 46 | 2.1 | Sand |
| M14 | 26 59 06 | 33 54 45 | Beach | Coarse sand |
| M15 | 26 59 09 | 33 54 42 | Beach | Coarse sand |
| M16 | 26 59 13 | 33 54 45 | 1.25 | Sandy mud with seagrasses |
| M17 | 26 59 17 | 33 54 43 | 2.4 | Muddy sand with seagrass |
| M18 | 26 59 20 | 33 54 41 | 1.2 | Biogenic sand |
| M19 | 26 59 24 | 33 54 40 | 9 | Biogenic sand |
| M20 | 26 59 25 | 33 54 39 | 17 | Very fine sand |
| M21 | 26 59 27 | 33 54 31 | 19 | Very fine sand |
| M22 | 26 59 27 | 33 54 22 | 21 | Sandy mud with seagrasses |
| M23 | 26 59 27 | 33 54 15 | 8 | Very fine sand with seagrasses |
| M24 | 26 59 27 | 33 54 13 | 3.5 | Very fine sand with seagrasses |
| M25 | 26 59 25 | 33 54 11 | 2.5 | Sandy mud with seagrasses |
| M26 | 26 59 21 | 33 54 10 | 3.2 | Sandy mud with seagrasses |
| M27 | 26 59 28 | 33 54 09 | Beach | Sandy gravel |
| M28 | 26 59 33 | 33 54 09 | 10 | Muddy sand with seagrasses |
| M29 | 26 59 38 | 33 54 07 | 17 | Biogenic sand |
| M30 | 26 59 39 | 33 54 05 | 10 | Muddy sand with seagrasses |
| M31 | 26 59 42 | 33 54 03 | 6 | Muddy sand with seagrasses |
| M32 | 26 59 38 | 33 54 00 | 25 | Muddy sand with seagrasses |

with skewness (Sk_i) ($r = 0.55$), mud with total organic matter (TOM) ($r = 0.47$), Mean Size (MZ) with kurtosis (K_G) ($r = 0.58$), Mean Size (MZ) with TOM ($r = 0.47$), carbonate content with total organic matter (TOM) ($r = 0.52$) (Table 3). The marine sediments collected from the study area had mean size (Mz) values ranging from -0.32 – 4.06Φ , averaging 2.6Φ (Table 2). Sorting (ϕ) of sediments varies between 0.65 and 1.95Φ averaging 1.28Φ (Table 2). The inclusive graphic skewness (SK_i) of the collected sediments ranges from -0.54 to 0.5Φ , averaging -0.06Φ (Table 2). The graphic kurtosis (KG) of sediments sampled from Sharm Abu-Makhadeg area ranges from 0.68 – 5.09Φ , averaging 2.13Φ (Table 2).

Cluster analysis (using Ward's method) includes gravel, sand and mud content separates all samples (32 samples) of Sharm Abu-Makhadeg area into 6 main clusters according to the abundance of size fraction (Fig. 4). Of these clusters only three had a high number of samples. Clusters 1, 2 and 6

constitute 72% of the total samples and are characterized by very high sand fraction 88.3, 94.4 and 79.9%, respectively. Most samples of clusters 1, 2 and 6 fall in shallow marine environment and offshore area (Fig. 2). Cluster 3, 4 and 5 represented 28% of the total samples and were distinguished by the lowest content of mud except cluster 5. Sediment samples of clusters 3 and 4 were belonging to the supratidal and beach areas and some samples from offshore area (Fig. 2). On the other hand, cluster 5 recorded the highest content of mud compared with the other clusters. Sediment samples of clusters 5 were belonging offshore area (Fig. 4).

Geochemistry

Carbonate content: In general, the total carbonate in the Sharm sediments was between 2.94 and 84.17% with an average 43.18% (Table 2).

Table 2: Results of grain size and geochemical analysis of surface marine sediments at Sharm Abu-Makhadeg area

| Sa. No. | Sediment types | | | Grain size parameters | | | | Geochemical analysis | | | Depth (m) |
|---------|----------------|-------|-------|-----------------------|----------|-----------------|----------------|----------------------|--------|---------|-----------|
| | Gravel | Sand | Mud | Mz | ϕ_i | Sk _i | K _G | Carb. (%) | OC (%) | TOM (%) | |
| M1 | 12.80 | 86.20 | 1.00 | 1.53 | 1.68 | -0.37 | 0.77 | 27.71 | 1.22 | 1.53 | Beach |
| M2 | 1.61 | 86.42 | 11.97 | 3.24 | 1.02 | 0.05 | 2.84 | 44.90 | 2.66 | 3.33 | 4 |
| M3 | 4.85 | 90.20 | 4.96 | 2.53 | 1.35 | -0.54 | 1.90 | 43.78 | 1.78 | 2.22 | 4 |
| M4 | 5.73 | 81.05 | 13.22 | 3.11 | 1.57 | -0.22 | 3.69 | 55.05 | 1.85 | 2.31 | Beach |
| M5 | 30.11 | 69.89 | 0.002 | -0.32 | 0.79 | 0.12 | 0.89 | 2.94 | 0.54 | 0.67 | 2 |
| M6 | 4.58 | 85.42 | 10.00 | 3.10 | 1.38 | -0.16 | 3.33 | 37.47 | 2.14 | 2.68 | 2.2 |
| M7 | 1.64 | 89.32 | 9.05 | 3.15 | 1.20 | -0.16 | 3.43 | 25.00 | 2.02 | 2.52 | 1.5 |
| M8 | 2.92 | 81.08 | 16.00 | 3.36 | 1.31 | 0.04 | 4.38 | 51.55 | 1.56 | 1.95 | 2.2 |
| M9 | 3.43 | 79.05 | 17.53 | 2.93 | 1.82 | -0.13 | 1.93 | 69.17 | 4.98 | 6.22 | 2.9 |
| M10 | 0.14 | 96.80 | 3.06 | 2.06 | 1.18 | -0.06 | 0.85 | 84.17 | 2.20 | 2.75 | 1.4 |
| M11 | 4.14 | 77.54 | 18.32 | 3.69 | 1.46 | -0.02 | 5.09 | 37.91 | 3.02 | 3.78 | 2.6 |
| M12 | 6.62 | 79.95 | 13.44 | 2.78 | 1.79 | -0.31 | 3.32 | 54.22 | 2.66 | 3.33 | 2.1 |
| M13 | 6.10 | 86.51 | 7.39 | 2.84 | 1.47 | -0.41 | 3.51 | 45.99 | 1.66 | 2.08 | 2.1 |
| M14 | 12.61 | 87.34 | 0.05 | -0.20 | 0.65 | -0.05 | 1.09 | 3.68 | 0.15 | 0.19 | Beach |
| M15 | 11.50 | 87.97 | 0.53 | 0.54 | 1.42 | 0.43 | 0.68 | 28.98 | 1.80 | 2.25 | Beach |
| M16 | 0.54 | 69.77 | 29.69 | 3.84 | 1.51 | 0.50 | 1.24 | 17.31 | 1.20 | 1.50 | 1.25 |
| M17 | 6.56 | 75.49 | 17.95 | 2.95 | 1.95 | -0.20 | 2.56 | 59.32 | 2.92 | 3.65 | 2.4 |
| M18 | 0.96 | 96.08 | 2.97 | 1.94 | 1.33 | -0.13 | 0.87 | 79.64 | 3.06 | 3.82 | 1.2 |
| M19 | 1.21 | 96.70 | 2.09 | 2.23 | 1.14 | -0.33 | 1.27 | 56.64 | 2.22 | 2.78 | 9 |
| M20 | 0.78 | 90.34 | 8.89 | 2.91 | 0.86 | -0.21 | 1.47 | 66.73 | 2.12 | 2.65 | 17 |
| M21 | 0.73 | 90.86 | 8.41 | 3.04 | 1.07 | -0.11 | 2.23 | 72.72 | 2.75 | 3.44 | 19 |
| M22 | 1.80 | 70.03 | 28.17 | 3.79 | 1.70 | 0.08 | 3.01 | 58.68 | 2.79 | 3.49 | 21 |
| M23 | 0.70 | 94.30 | 5.00 | 3.04 | 0.75 | -0.31 | 1.36 | 20.09 | 0.98 | 1.23 | 8 |
| M24 | 2.11 | 94.50 | 3.39 | 2.71 | 0.83 | -0.20 | 1.81 | 26.20 | 1.37 | 1.71 | 3.5 |
| M25 | 0.92 | 68.85 | 30.24 | 4.06 | 1.39 | 0.44 | 1.78 | 37.36 | 4.17 | 5.21 | 2.5 |
| M26 | 1.16 | 70.26 | 28.58 | 3.94 | 1.44 | 0.43 | 1.80 | 22.47 | 3.76 | 4.70 | 3.2 |
| M27 | 24.35 | 75.56 | 0.09 | -0.31 | 0.69 | -0.10 | 0.72 | 17.94 | 2.20 | 2.75 | Beach |
| M28 | 2.59 | 83.21 | 14.20 | 2.68 | 1.72 | -0.19 | 1.64 | 46.69 | 3.83 | 4.79 | 10 |
| M29 | 1.32 | 92.54 | 6.14 | 2.57 | 1.24 | -0.25 | 1.33 | 62.45 | 4.34 | 5.42 | 17 |
| M30 | 1.80 | 89.84 | 8.36 | 2.91 | 1.19 | -0.14 | 1.82 | 53.86 | 3.44 | 4.30 | 10 |
| M31 | 0.23 | 87.40 | 12.36 | 3.17 | 1.13 | 0.05 | 2.53 | 42.50 | 3.48 | 4.35 | 6 |
| M32 | 0.21 | 81.72 | 18.07 | 3.62 | 1.03 | 0.39 | 3.10 | 28.57 | 2.90 | 3.62 | 25 |
| Min. | 0.14 | 68.85 | 0.001 | -0.32 | 0.65 | -0.54 | 0.68 | 2.94 | 0.15 | 0.19 | 0.00 |
| Max. | 30.11 | 96.80 | 30.24 | 4.06 | 1.95 | 0.50 | 5.09 | 84.17 | 4.98 | 6.22 | 25.00 |
| Avg. | 4.90 | 84.13 | 10.97 | 2.61 | 1.28 | -0.06 | 2.13 | 43.18 | 2.43 | 3.04 | 5.72 |

Mz: mean size, ϕ_i : Sorting, Sk_i: Skewness, K_G: Kurtosis, Carb.: Carbonate content, *Values ppm, OC: Organic carbon. Source: Folk and Ward²⁸

Table 3: Correlation coefficients between sediment types, grain size parameters, carbonate content, total organic matter, depth of surface marine sediments at Sharm Abu-Makhadeg area

| Parameters | Gravel | Sand | Mud | Mz | ϕ_i | Sk _i | K _G | Carb. (%) | TOM (%) | Depth |
|-----------------|--------|-------|------|------|----------|-----------------|----------------|-----------|---------|-------|
| Gravel | 1 | | | | | | | | | |
| Sand | -0.33 | 1 | | | | | | | | |
| Mud | -0.45 | -0.70 | 1 | | | | | | | |
| Mz | -0.82 | -0.15 | 0.77 | 1 | | | | | | |
| ϕ_i | -0.23 | -0.35 | 0.51 | 0.42 | 1 | | | | | |
| Sk _i | -0.02 | -0.56 | 0.55 | 0.15 | -0.03 | 1 | | | | |
| K _G | -0.29 | -0.21 | 0.42 | 0.58 | 0.31 | -0.06 | 1 | | | |
| Carb.% | -0.52 | 0.36 | 0.05 | 0.33 | 0.38 | -0.33 | 0.12 | 1 | | |
| TOM% | -0.44 | -0.14 | 0.47 | 0.47 | 0.44 | 0.14 | 0.12 | 0.52 | 1 | |
| Depth | -0.38 | 0.11 | 0.19 | 0.33 | -0.13 | 0.04 | 0.06 | 0.29 | 0.31 | 1 |

Organic Carbon (OC) and Total Organic Matter (TOM): In general, the sediments had values of organic Carbon 0.15-4.98% with average 2.43% and total organic matter content varying from 0.19-6.22% with average 3.04%

(Table 2). The results of correlation coefficient illustrated in Table 3, weak positive correlation between TOM and depth and high positive correlation coefficient between TOM and carbonate (Table 3).




Fig. 4: Dendrogram from cluster analysis (ward's method) and histogram exhibiting cluster of grain size texture

DISCUSSION

Sediments in Sharm Abu-Makhadeg as well as in the beach, intertidal and lagoons are composed of a mixture of carbonates and siliciclastics. However, biogenic activity is regarded as the major source of sediments few meters away from the wadi mouth. Intertidal sediments are mainly sands compared to beach sediments, gravel content is low and mud is a little higher. Sediments of lagoons are finer than those of beach and intertidal area. Generally, the sediments of the investigated area were found to consist of a wide variety of

texture classes, from coarse sand to sandy mud. The marine sediments in the study area are mainly composed of sand. Generally, the mud contents seem to be increase with increasing water depth and distance from the shoreline. The reason for this high mud content is due to the landfill operation in the past.

Cluster analysis (using Ward's method) includes gravel, sand and mud content separates all samples (32 samples) of Sharm Abu-Makhadeg area into 6 main clusters. According to cluster analysis fine grains are transported by sea waves to the offshore. According to Mansour¹ waves and currents

redistribute terrigenous debris carried into the sea either via wadi or NW winds on the tidal flat and most likely also sweep some of the fine terrigenous sediments from the submarine slopes into the deeps. The distribution of the coarser sediments may reflect the abundance of terrigenous sediments in the beach and biogenic fragments in the reef areas. Generally, the sediments of Sharm Abu-Makhadeg are characterized by a fine texture. Sand is the main category among the three constituents as showed (Fig. 4).

The distribution of mean grain size (M_Z) is included here for comparison with those of the various geochemical parameters. The influence of these sources is more or less localized resulting in a particular textural and compositional distribution, especially in the nearshore area. The marine sediments collected from the study area have mean size (M_Z) values fine sands. Sorting (ϕ_1) of sediments are poorly sorted. The sediments in the study area show change from strongly coarse skewed and fine skewed near the beaches to nearly symmetrical and coarse skewed distribution with increasing water depth. The kurtosis (K_G) values of this area do not show any trend for increase or decrease with water depth.

Most samples of beach and intertidal have low carbonates content resulted from very high landfilling and dredging, as a result of human activities carried out in the coastal region during the 1990s. High carbonate content is recorded in samples away from offshore areas. The biogenic constituents are relatively high in the seaward sediments less influenced by terrigenous constituent. Mansour *et al.*²⁴ found that the carbonate content of shallow marine sediments between Hamata and Gemsha along the Egyptian Red sea coast varies from 39.98-84.40% with an average of 77.56. This average value is higher than that recorded in Sharm Abu-Makhadeg. This is attributed to the increasing in landfilling and dredging income to the tidal flat in Sharm Abu-Makhadeg area.

The investigated area is subjected to occasional torrents rich with fine sediments with high organic content. Therefore, sediments with darkish gray color cover most area. However, samples with high organic content occur mainly in the Sharm itself and in the dredged areas of tidal flat in the past. In general, the average organic carbon and total organic matter contents of the sediments are relatively higher than those of other areas in the Red sea³⁰. This is probably due to this high organic productivity. The terrestrial input of organic matter by wadi might be very high compared with that from the plant debris and animals inhabiting its waters and living on its bottom. The high content of organic matter is probably responsible for metal enrichment.

Sharm Abu Makhadeg is a small embayment and construction marinas in front of each hotel will close the Sharm and destroy the environment of the area, a practice

that has been widespread in Hurghada and Safaga. The other problem is that the construction of private embankments used as jetties should be halted and jetties should only be allowed if they are floating or constructed on pillars, which allow the natural flow of sedimentation. Human activities may increase turbidity, alter water circulation and sediment distribution patterns and may destroy entire reef systems. The unnatural obstruction in the water disrupted tidal flows and the natural deposit of sediments, so that after a certain time hundreds of m² of land were ebbed away from the property, ruining the plans an architect had designed for land. The fine grained sediment resulting from the development activity increases the water turbidity and now the visibility of water is very poor.

CONCLUSION

Abu Makhadeg area are rich with organic matter. Fortunately, sediments on both sides of the sharm are covered with rocky limestone o tidal flat. Dredging of this rocky cover reveals fine sediments and concentrate organic matter and trace metals in confined pools. The contribution of land-derived materials is significant high in bottom sediments of Sharm Abu-Makhadeg. Cooperation between the investors and the EEAA is very important in order to centralize decisions and responsibilities. Sharing the developers of the area in constructing new marina, cornish in front of their resorts and round road behind their properties will be very faithful and decrease the environmental problems.

SIGNIFICANCE STATEMENT

The scope of the present work are monitoring of changes of sediment nature by environmental impacts of Sharm Abu-Makhadeg area and record all development activities during the past 20 years. The impact of development activities in Sharm Abu-Makhadeg area was measured by surveying and by analyzing the surface marine sediment. These study Reasonable recommendations have been outlined to offer guidelines to mitigate the impact on the marine environment. These recommendations are something of a compromise between the economic interests and environmental concerns in the area.

ACKNOWLEDGMENT

The samples were collected through the National Institute of Oceanography and Fisheries plan, Red Sea Branch, Hurghada, Egypt.

REFERENCES

1. Mansour, A.M., 1999. Changes of sediment nature by environmental impacts of Sharm Abu Makhadeg area, Red Sea, Egypt. *Sedimentol. Egypt*, 7: 25-36.
2. Ahmed, N.A. and H.A. Madkour, 2006. The environmental impacts of the Red Sea coast at Quseir District, Red Sea, Egypt. *Proceedings of the 3rd International Conference for Development and the Environment in the Arab World*, March 21-23, 2006, Assiut, Egypt, pp: 733-757.
3. Dar, A.M., 2002. Geological basis to study the environmental defect in the marine ecosystem as a result of tourist activities in Hurghada area and surroundings, red sea, Egypt. Ph.D. Thesis, Suez Canal University, Egypt.
4. El-Askary, M.A., S.M. Nasr, A.A. Moussa and M.H. El-Mamony, 1988. Geochemical approach to the beach and bottom sediments of the Jubal area at the entrance of the Gulf of Suez. *Bull. Inst. Oceanogr. Fish.*, 14/1: 105-121.
5. El-Mamoney, M.H., 1995. Evaluation of terrestrial contribution to the red sea sediments, Egypt. Ph.D. Thesis, Faculty of Science, Alexandria University, Egypt.
6. El-Sayed, M.K., 1984. Reefal sediments of Al-Ghardaqa, Northern Red Sea, Egypt. *Mar. Geol.*, 56: 259-271.
7. El-Taher, A. and H.A. Madkour, 2011. Distribution and environmental impacts of metals and natural radionuclides in marine sediments in-front of different wadies mouth along the Egyptian Red Sea Coast. *Applied Radiat. Isot.*, 69: 550-558.
8. El-Taher, A. and H.A. Madkour, 2013. Texture and environmental radioactivity measurements of Safaga sand dunes. *Indian J. Geo-Mar. Sci.*, 42: 35-41.
9. El-Taher, A. and H.A. Madkour, 2014. Environmental and radio-ecological studies on shallow marine sediments from harbour areas along the Red Sea coast of Egypt for identification of anthropogenic impacts. *Isotopes Environ. Health Stud.*, 50: 120-133.
10. El-Taher, A., S. Alashrah, H.A. Madkour, A. Al-Sayed and T. El-Erian, 2018. Radionuclides distribution in marine sediment from Abu Soma Bay, Egyptian Red Sea Coast. *J. Environ. Sci. Technol.*, 11: 95-103.
11. Frihy, O.E., A.M. Fanos, A.A. Khafagy and K.A. Abu Aesha, 1996. Human impacts on the coastal zone of Hurghada, Northern Red Sea, Egypt. *Geo-Mar. Lett.*, 16: 324-329.
12. Hamouda, A.Z. and M.A. El-Wahhab, 2009. Sediment characteristics and water circulation of Big Jemsa Bay, Red Sea, Egypt. *Mar. Geophys. Res.*, 30: 95-104.
13. Hamouda, A.Z. and M.A. El-Wahhab, 2012. Detection of the bottom facies characteristics at El Zeit Bay, Red Sea, by using a single-beam acoustic sound. *Oceanol. J.*, 52: 60-71.
14. Madkour, H.A. and M.A. Dar, 2007. The anthropogenic effluents of the human activities on the red sea Coast at Hurghada Harbour (Case study). *Egypt. J. Aquat. Res.*, 33: 43-58.
15. Madkour, H.A., A. El-Taher, A.N. El-Hagag Ahmed, A.W. Mohamed and T.M. El-Erian, 2012. Contamination of coastal sediments in El-Hamrawein Harbour, Red Sea, Egypt. *J. Environ. Sci. Technol.*, 5: 210-221.
16. Madkour, H.A., A.M. Mansour, A.E.H.N. Ahmed and A. El-Taher, 2014. Environmental texture and geochemistry of the sediments of a subtropical mangrove ecosystem and surrounding areas, Red Sea Coast, Egypt. *Arab J. Geol. Sci.*, 7: 3427-3440.
17. Nawar, A.H., A.M. Mohamed, A.W. Mohamed and H.A. Madkour, 1997. Landfilling as a controlling factors of bottom sediments physicochemical characteristics. *Proceedings of the 1st International Conference and Trade Fair on Environmental Management and Technologies*, February 16-18, 1997, Cairo, Egypt.
18. Madkour, H.A., M.A.K. Abdelhalim and A. El-Taher, 2013. Assessment of heavy metals concentrations resulting natural inputs in Wadi El-Gemal surface sediments, Red Sea coast. *Life Sci. J.*, 10: 686-694.
19. Piller, W.E. and A.M. Mansour, 1990. The Northern Bay of Safaga (Red Sea, Egypt) An actioupalaeontological approach II Sediment analysis and sedimentary facies. *Beitr. Palaont. Osterr.*, 16: 1-102.
20. Madkour, H.A., M.A.K. Abdelhalim, K.A. Obirikorang, A.W. Mohamed, A.E.H.N. Ahmed and A. El-Taher, 2015. Environmental implications of surface sediments from coastal lagoons in the Red Sea coast. *J. Environ. Biol.*, 36: 1421-1427.
21. Malecki, G., 1986. SEDPAK-charakterisierung von sedimenten aufgrund der korngr. Analyse. *Ber. Geol. B.-A., Blg. 12*, Wien.
22. Mansour, A.M., 1995. Sedimentary facies and carbonate-siliciclastic transition of Sharm El Bahari and Sharm El Qibli, Red Sea, Egypt. *Egypt. J. Geol.*, 39: 57-76.
23. Mansour, A.M., A.H. Nawar and A.M. Mohamed, 1997. Recent intertidal sediments and negative impact of human activities, Red Sea coast, Egypt. *Egypt. J. Geol.*, 41: 239-272.
24. Mansour, A.M., A.H. Nawar and A.W. Mohamed, 2000. Geochemistry of coastal marine sediments and their contaminant metals, Red Sea, Egypt: A legacy for the future and a tracer to modern sediment dynamics. *Sedimentol. Egypt*, 8: 231-242.
25. Mansour, A.M., A.H. Nawar and H.M. Madkour, 2005. Metals concentration of recent invertebrates along the Red Sea Coast of Egypt: A Tool for monitoring environmental hazards. *Sedimentol. Egypt*, 3: 171-185.

26. Maxwell, W.G.H., 1968. Atlas of the Great Barrier Reef. Elsevier, Amsterdam, Pages: 268.
27. Mohamed, M.A.E. H.A. Madkour and M.I. El-Saman, 2011. Impact of anthropogenic activities and natural inputs on oceanographic characteristics of water and geochemistry of surface sediments in different sites along the Egyptian Red Sea coast. *Afr. J. Environ. Sci. Technol.*, 5: 494-511.
28. Folk, R.L. and W.C. Ward, 1957. Brazos river bar: A study in the significance of grain size. *J. Sediment. Petrol.*, 27: 3-26.
29. Dean, Jr. W.E., 1974. Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition: Comparison with other methods. *J. Sediment. Res.*, 44: 242-248.
30. Madkour, H.A.M., 2015. Detection of damaged areas due to tourism development along the Egyptian Red Sea coast using GIS, remote sensing and foraminifera. State of the Art, National Institute of Oceanography and Fisheries (NIOF), Red Sea Branch, pp: 1-135.