



Journal of Medical Sciences

ISSN 1682-4474

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

JMS (ISSN 1682-4474) is an International, peer-reviewed scientific journal that publishes original article in experimental & clinical medicine and related disciplines such as molecular biology, biochemistry, genetics, biophysics, bio-and medical technology. JMS is issued six times per year on paper and in electronic format.

For further information about this article or if you need reprints, please contact:

Wilson Abrão Saad
Department of Physiology and Pathology, School of Dentistry
Paulista State University, UNESP
Rua Humaitá, 1680
14801-903-Araraquara, SP, Brazil

Tel: +55(16) 3301-6488
Fax: +55(16) 3322-4118

Nitregic Pathways and L-Type Calcium Channel of MnPO Influencing Cardiovascular Homeostasis

^{1,5}Wilson Abrão Saad, ⁴Ismael Francisco Motta Siqueira Guarda, ³Luiz Antonio de Arruda, Camargo, ¹Talmir Augusto Faria Brizola dos Santos, ¹Sylvio Simões and ⁶William Abrão Saad

The median preoptic nucleus (MnPO) is one of most important site of the lamina terminalis implicated in the regulation of hydro electrolytic and cardiovascular balance. The purpose of this study was to determine the effect of L-Type calcium channel antagonist, nifedipine, on the increase of median arterial blood pressure (MAP) induce by angiotensin II (ANG II) injected into the MnPO. The influence of nitric oxide (NO) on nifedipine antipressor action has also been studied by utilizing N^w-nitro-L-arginine methyl ester (L-NAME) (40 µg 0.2 µL⁻¹) a NO synthase inhibitor (NOSI), 7-nitroindazole (7-NIT) (40 µg 0.2 µL⁻¹), a specific neuronal NO synthase inhibitor (nNOSI) and sodium nitroprusside (SNP) (20 µg 0.2 µL⁻¹) a NO donor agent. We have also investigated the central role of losartan and PD123349 (20 nmol 0.2 µL⁻¹), AT₁ and AT₂, respectively (selective non peptide ANG II receptor antagonists), in the pressor effect of ANG II (25 pmol 0.2 µL⁻¹) injected into the MnPO. Male Wistar rats weighting 200-250 g, with cannulae implanted into the MnPO were utilized. Losartan injected into the MnPO, prior to ANG II, blocked the pressor effect of ANGII. PD 123319 only decreased the pressor effect of ANG II. Rats pre-treated with either 50 µg 0.2 µL⁻¹ or 100 µg 0.2 µL⁻¹ of nifedipine, followed by 25 pmol 0.2 µL⁻¹ of ANG II, decreased ANG II-pressor effect. L-NAME potentiated the pressor effect of ANG II. 7-NIT injected prior to ANG II into the MnPO also potentiated the pressor effect of ANGII but with less intensity than that of L-NAME. SNP injected prior to ANG II blocked the pressor effect of ANG II. The potentiation action of L-NAME and 7-NIT on ANG II-pressor effect was blocked by prior injection of nifedipine. The results described in this study provide evidence that calcium channels play important roles in central ANG II-induced pressor effect. The structures containing NO in the brain, such as MnPO, include both endothelial and neuronal cells, which might be responsible for the influence of nifedipine on the pressor effect of ANG II. These data have shown the functional relationship between L-Type calcium channel and a free radical gas NO in the MnPO, on the control of ANG II-induced pressor effect acting in AT₁ and AT₂ receptors.

Key words: Angiotensin II, rats, calcium channel, nitric oxide, blood pressure, MnPO

¹Basic Institute of Biosciences-UNITAU, Taubaté, SP, Brazil

²Department of Exact and Natural Science UNIARA Araraquara SP Brazil

³Department of Physiology and Pathology, School of Dentistry, Paulista State University, UNESP Araraquara, S Brazil

⁴Department of Anesthesiology Clinic Hospital State of São Paulo, São Paulo, Brazil

⁵Department of Physiology, Federal University of São Carlos SP-Brazil

⁶Department of Gastroenterology, Clinic Hospital of University of São Paulo USP São Paulo Brazil

INTRODUCTION

Recent experiments of our laboratory demonstrated that the NO has interrelationship with AT₁ and AT₂ receptors of the CNS in the pressor effect of ANG II. The influence of vasopressin receptor and nitric oxide on the water, sodium intake and arterial blood pressure induced by angiotensin injected into the third ventricle of the rat brain (Saad *et al.*, 2006).

The influence of calcium channel in the pressor effect of ANG II has not yet been studied. Centrally ANG II elicits responses such as increased blood pressure, thirst, sodium appetite and the release of vasopressin (Reid, 1988; Saad *et al.*, 2002a). Nitric oxide synthase such as neuronal NOS, are expressed in the MnPO (Summy-Long *et al.*, 2002). N^w-nitro-L-arginine methyl ester (L-NAME) reduces renal blood flow, urine flow rate and urinary sodium excretion (Naees *et al.*, 1992). The influence of NO of the Central Nervous System (CNS) on several physiological parameters has been demonstrated (Saad *et al.*, 2002b; Saad *et al.*, 2003; Saad *et al.*, 2004a). NO plays an important role in the hydromineral and cardiovascular regulation induced by ANG II (Saad *et al.*, 1999). NO may facilitate the release of excitatory transmitters, possibly through a presynaptic cyclic GMP-dependent mechanism (Wu *et al.*, 1997). L-NAME increases blood pressure which is at least in part salt sensitive (Hodge *et al.*, 2002). Paraventricular neurons of the hypothalamus (PVN) have an important role in central regulatory mechanisms such as the temperature regulation, by releasing NO. Pharmacodynamic and pharmacokinetic studies of the 7-NIT also proved the pharmacological participation of NO in the CNS (Bush and Pollak, 2001).

It has been widely accepted that calcium ions are critically important for fast synaptic transmission, including the processes of neurotransmitter release from pre-synaptic terminals and post-synaptic receptor-mediated events (Ghosh and Greenberg, 1995). A major factor determining a neuronal Ca²⁺-dependent signal is the opening of permeability pathways for Ca²⁺ in the cell membrane (Tsien *et al.*, 1988). NMDA-type glutamate receptors seem to be concerned with the rapid behavioral actions of ANG II (Zhu and Herbert, 1997a).

Several data indicate that the MnPO is indeed the target of afferent from chemosensitive, osmosensitive and barosensitive systems concerned with fluid homeostasis and cardiovascular regulation (Gardiner and Stricker, 1985; O'Neill and Brody, 1987; Cunningham *et al.*, 2002; Yashuda *et al.*, 2002). The neuronal circuit involving both the MnPO and the organ vasculosum of the lamina terminalis (OVLT) has been reported to play an

important role in vasopressin secretion and thirst (Mangiapani *et al.*, 1983). The injection of L-NAME a nitric oxide synthase inhibitor, into the MnPO, increased the Mean Arterial Pressure (MAP) (Saad *et al.*, 2004a).

The influence of nifedipine L-Type calcium channel antagonist and NO in the CNS on the cardiovascular regulation has been previously demonstrated in our laboratory utilizing the Lateral Ventricle (LV) as the site of ANG II, nifedipine and L-NAME injection. The antidipsogenic actions of two L-type calcium channel blockers, verapamil and diltiazem injected into LV (Zhu and Herbert, 1997b) which bind to the phenylalkylamine and the benzothiazepine sites respectively did not significantly suppress ANG II-induced drinking behavior, however nifedipine injected prior to ANG II into LV suppress the dipsogenic effect of ANG II, suggesting that ANG II acts through a dihydropyridine-type calcium channel. These experiments verify whether the calcium channels has interrelationship with NO and ANG II receptors, of the MnPO, in the blood pressure regulation. The objective of this study determined the role of voltage-sensitive L-Type calcium channels in ANG II-induced pressor response when injected into the MnPO and the influence of NO, AT₁ and AT₂ ANG II receptors was also explored.

MATERIALS AND METHODS

This study was conducted in the Basic Institute of Biosciences-UNITAU, Taubaté, SP, BRAZIL and in the Department of Biology and Health Science UNIARA Araraquara SP BRAZIL in the year of 2004/2005.

All experiments were performed in adult male Wistar rats weighing 200-250 g. The animals were housed in individual metabolic cages. Food (Purina Rat Chow) and tap water were available *ad libitum*. The temperature was maintained at 22±2°C. The light cycle was held at 12:12 with lights on 06:00 h. All experiments were conducted during the light period, between 09:00 AM and 03:00 PM.

The Medical Ethics Committee of the Universidade Estadual Paulista UNESP approved all protocols in this study.

Experimental protocol

Experiment 1: There were six groups of rats: Rats were injected with the following: saline+saline 0.15 M NaCl (Group 1, n = 10). Saline+ANG II 25 pmol (Group 2 n = 9). Saline+Losartan (Group 3 n = 9). Saline+PD 123319 (Group 4 n = 8). Losartan+ANG II (Group 5 n = 8). PD 123319+ANG II (Group 6 n = 7). After the second injection, MAP was recorded during 30 min.

Experiment 2: There were six groups of rats. Rats were injected with the following: saline 0.15 M NaCl, (Group 1, n = 9). Saline+ANGII 25 pmol (Group 2, n = 9). Saline+Nifedipine 50 µg (Group 3; n = 8). Nifedipine 50 µg followed 10 min later by 25 pmol ANGII (Group 4, n = 7). Nifedipine 100 µg followed 10 min later by 25 pmol ANGII (Group 5, n = 7) and Sodium nitroprusside 20 µg followed 10 min later by 25 pmol ANGII (Group 6, n = 7). After the second injection, MAP was recorded during 30 min.

Experiment 3: There were seven groups of rats. Rats were injected with the following: (saline 0.15 M NaCl, (Group 1, n = 10). Saline+ANGII 25 pmol (Group 2, n = 10). Nifedipine 100 µg followed 10 min later by 25 pmol ANGII (Group 3, n = 9). L-NAME 40 µg followed 10 min later by 25 pmol ANGII (Group 4, n = 9). 7-NIT 40 µg followed 10 min later by 25 pmol ANGII (Group 5, n = 8). L-NAME 40 µg+Nifedipine 100 µg followed 10 min later by 25 pmol ANGII (Group 6, n = 7). 7-NIT 40 µg+Nifedipine 100 µg followed 10 min later by 25 pmol ANGII (Group 7, n = 6). After the second injection, MAP was recorded during 30 min.

Drugs

- Saline 0.15M NaCl (control)
- PD123319 and losartan (20 nmol 0.2 µL⁻¹) purchased from DuPont, Merck, (Wilmington, DE USA)
- Saline 0.15M NaCl (control)
- Angiotensin II (25 pmol 0.2 µL⁻¹) purchased from Sigma (Chemical Co., St. Louis, MO)
- Nifedipine (50 µg 0.2 µL⁻¹ or 100 µg 0.2 µL⁻¹) purchased from Sigma (Chemical Co., St. Louis, MO)
- L-NAME (40 µg 0.2 µL⁻¹) purchased from Sigma (Chemical Co., St. Louis, MO)
- 7-Nitroindazole (7-NIT) (40 µg 0.2 µL⁻¹) purchased from Sigma (Chemical Co., St. Louis, MO)
- Sodium Nitroprusside (20 µg 0.2 µL⁻¹) purchased from Sigma (Chemical Co., St. Louis, MO)

Cerebral surgery: Male Holtzman rats weighing 250-300 g were anesthetized with zoletil 50 mg kg⁻¹ (tiletamine chloridrate 125 mg and zolazepan chloridrate 125 mg) into quadriceps muscle. A stainless steel cannula with 10 and 12 mm long and 0.7 mm OD was implanted into the MnPO according to the coordinates of Paxinos and Watson (1986) atlas rat brain. The cannulae were fixed to the skull with the aid of jeweler screws and dental acrylic resin and protected with a stiletto. The experiments started after 5 days of the brain surgery.

Vascular catheter: After the animals recovery from brain surgery (5 days) PE-10 polyethylene tubing connected to PE-50 tubing was inserted into the abdominal aorta




Fig. 1: Photomicrograph of a hematoxylin stained transverse section of the rat brain showing site of injection into the MnPO (arrow). CC Corpus Callosum; LV Lateral Ventricle; AC Anterior commissure

through the femoral artery under zoletil 50 mg kg⁻¹ (tiletamine chloridrate 125 mg and zolazepan chloridrate 125 mg) anaesthesia. The polyethylene tube was tunneled subcutaneously to the back of the rat and externalized at the dorsal cervical region. Catheters were fill with heparinized saline and plugged with 23-G obturators. Rats recovered from surgery (vascular catheter) for a minimum of 24 h before beginning of testing.

Drug injection: The drugs were injected into the MnPO by using a Hamilton micro syringe (5 µL) connected by a PE-10 polyethylene tubing (25 cm) to a needle (0.3 mm o.d.), which was introduced into the brain through the cannula previously fixed to the animals' head. The volume of injection was always 0.2 µL injected over a period of 30-60 sec.

Arterial blood pressure recordings: Direct mean arterial blood pressure (MAP) was record in anesthetized and unrestrained rats. The animal was remove from the home cage and placed in a test cage, without access to food or water. The previously implanted catheter was connected to a Statham (P23 Db) pressure transducer (Statham-Gould, Valley View, OH) coupled to a multi channel recorded (PowerLab multirecord). This program permits the acquisition of cardiovascular data by computer.

Histology: At the end of the experiments, the rats were anesthetized with ether and perfused with saline and buffered formalin. The brains were removed, fixed in 10% formalin, frozen to -25°C and cut into 20-30 µm coronal sections. Subjects were included in the analysis only if their cannula was in the lateral medial portion of MnPO (Fig. 1). We examined every section throughout the

cannula path and eliminated all animals with cannula that perforated the ependyma, detected by histology and Evans blue.

Statistical analysis: The results are reported as mean±SEM. The ANOVA and Newman-Keuls post-hoc test were used to determine the significance. The values were considered statistically significant with 5% level ($p < 0.05$).

RESULTS

Effect of AT₁ and AT₂ ANG II receptor antagonists on the pressor effect of ANG II: Microinjections of ANG II into the MnPO caused an increase in MAP compared to control (29±3 mmHg vs. 5±1 mmHg, $p < 0.05$). The injection of saline+losartan and saline+PD 123319 into MnPO produced no change in MAP (5±1 mmHg and 4±1 mmHg, $p > 0.05$). The micro-injection of losartan into MnPO prior to ANGII abolished the increase in the MAP as it was previously seen when ANG II was micro-injected alone (5±1 mmHg, $p < 0.05$). The micro-injection of PD 123319 in to MnPO prior to ANG II decreased the pressor effect of ANG II with less intensity than that of losartan (8±1 mmHg, $p < 0.05$) Table 1.

Effects of nifedipine and sodium nitroprusside on the increase of MAP induced by the injection of ANG II into the MnPO: Microinjections of ANG II into the MnPO cause an increase in MAP compared to control (28±2 mmHg vs. control 5±1 mmHg, $p < 0.05$). The micro-injection of saline+nifedipine into the MnPO caused no change in the MAP (7±1 mmHg). Nifedipine 50 µg injected into the MnPO followed by ANG II decreased the pressor effect (8±1 mmHg, $p < 0.01$). Nifedipine 100 µg injected into the MnPO followed by ANG II blocked the pressor effect (3±0.5 mmHg, $p < 0.01$). The injection of SNP also blocked the pressor effect of ANG II (8±1 mmHg, $p < 0.01$). SNP injected alone into the MnPO produced no change in the MAP Table 2.

Effect of nifedipine, L-NAME and 7-NIT on ANG II pressor effects: The MAP of rats injected with ANG II into MnPO produced an increase in the MAP 29±1 vs control 6±2 mmHg. Nifedipine injected prior to ANG II produced a decrease in the pressor effect 4±1 mmHg ($p < 0.001$). The MAP of rats injected with L-NAME prior to ANGII was 35±1 mmHg ($p < 0.001$). The MAP of rats injected with 7-NIT prior to ANGII was 32±1 mmHg ($p < 0.001$). Nifedipine 100 µg prior to L-NAME followed by ANG II blocked the pressor effect (7±1 mmHg, $p < 0.05$). Rats injected with nifedipine 100 µg prior to 7-NIT followed by ANG II decreased the pressor effect (5±2 mmHg, $p < 0.001$) (Table 3).

DISCUSSION

These results showed the important participation of L-Type calcium channel and NO in the regulation of arterial pressure induced by ANG II injected into MnPO. These results also showed that the MnPO is more sensitive to the action of these drugs than when injected into the LV as previously demonstrated in our laboratory.

In the MnPO ANG II via AT₁ receptors mediates cardiovascular responses to an acute increase in cerebrospinal fluid sodium as well as the chronic pressor responses to high sodium intake in spontaneously hypertensive rats (Budzikowski and Leenen, 2001). The micro-injection of losartan into MnPO prior to ANGII abolished the MAP increase. The micro-injection of PD 123319 into MnPO prior to ANG II partially reduced the pressor effect of ANG II. There is functional evidence that ANG II exerts excitatory effects on neurons of the PVN and that these effects are antagonized by CGP 42112A (AT₂ receptor antagonist) (Felix *et al.*, 1991). These results showed that AT₁ and AT₂ receptors of the MnPO are important for the role of ANG II on central regulation of MAP. These findings are supported by the results of McKinley *et al.* (2003) showing that angiotensin AT₁, AT₂ and AT₄ receptors are also plentiful in the brain. In the MnPO ANG II via AT₁ receptors mediated cardiovascular

Table 1: Effect of nifedipine, L-NAME and 7-NIT on ANGII pressor effects

	SAL + SAL	SAL + ANGII	SAL + LOS	SAL + PD	LOS + ANGII	PD + ANGII
(mmHg)	5±1 ⁺	29±3*	5±1 ⁺	4±1 ⁺	5±1 ⁺	8±1**

⁺ $p < 0.05$ vs SAL+ANGII; * $p < 0.05$ vs SAL+SAL

Table 2: Effects of nifedipine and sodium nitroprusside on the increase of MAP induced by the injection of ANGII into the MnPO

	SAL + SAL	SAL + ANGII	SAL + NIF	NIF (50 µg) + ANGII	NIF (100 µg) + ANGII	SAL + SNP	SNP + ANGII
(mmHg)	5±1 ⁺	28±2*	7±1 ⁺	8±1 ⁺	3±1 ⁺	4±1 ⁺	8±1**

⁺ $p < 0.05$ vs SAL+ANGII; * $p < 0.05$ vs SAL+SAL

Table 3: Effect of nifedipine, L-NAME and 7-NIT on ANGII pressor effects

	SAL + SAL	SAL + ANGII	NIF + ANGII	LNA + ANGII	7NI + ANGII	NIF (100 µg) + LNA + ANGII	NIF (100 µg) + 7NI + ANGII
(mmHg)	6±2 ⁺	29±1*	4±1 ⁺	35±1**	32±1**	7±1 ⁺	5±2 ⁺

⁺ $p < 0.05$ vs SAL+ANGII; * $p < 0.05$ vs SAL+SAL

responses to an acute increase in CSF sodium as well as the chronic pressor responses to high sodium intake in SHR. Furthermore, the salt-sensitive component appears to be ANGII-dependent, as it was associated with increase of plasma ANG II levels and could be reversed by the treatment with ANG II receptor antagonist (Thunhorst and Johnson, 1994).

It has been demonstrated that NO plays an important role in the cardiovascular regulation (Saad *et al.*, 2002a). FK409, a NO donor agent, combined with losartan, abolished the increase in the MAP induced by ANG II (Saad *et al.*, 2002b). L-NAME induced an increase in blood pressure, which is at least in part salt sensitive. L-NAME produced an increase in the pressor effect of ANGII and this effect may be due to local vasoconstriction and all at once by neuronal NOS inhibition. We have also investigated the role of 7-NIT (nNOS) injected into the MnPO in the pressor effect of ANG II. 7-NIT has less effective potentiation effect on ANG II pressor effect than L-NAME when injected into the MnPO. Pharmacodynamic and pharmacokinetic studies of the 7-NIT proved the neural pharmacological participation of NO in the CNS in many physiological functions (Bush and Pollack, 2001). This let us to postulate that the eNOS and nNOS in the MnPO are involved in the pressor effect of ANG II.

Angiotensinergic neural pathways and calcium channels are important in neural function and may have important homeostatic roles, particularly related to cardiovascular function by involving NO. The activation of glutamate-NMDA receptors can increase intracellular Ca^{2+} concentration, which in turn acts on calmodulin to evoke nitric oxide syntheses (Garthwaite, 1991; Dawson and Snyder, 1994). In most neurons of the central nervous system, there are at least two major classes of calcium channel: voltage sensitive calcium channels (VSCCs) and receptor-operated calcium channels (ROC). Nifedipine may have interfered with Ca^{2+} influx in the presynaptic terminals, where L-type calcium channels play important roles in the modulating presynaptic neurotransmitter release (Miller, 1998; Ghosh and Greenberg, 1995). It may also have altered Ca^{2+} -dependent signal events in postsynaptic neurons since previous studies demonstrated the permissive effects of VSCCs on NMDA receptor-mediated Ca^{2+} influx (Bumashev, 1996). ANG II into MnPO may act primarily on AT_1 receptors that increase intracellular Ca^{2+} concentration, which in turn acts on calmodulin to evoke nitric oxide synthesis. Central NO-cGMP pathway is involved in ANGII-induced drinking behavior (Zhu and Herbeert, 1997b). The NOS and your RNAm is plentiful found into MnPO (Summy-Long *et al.*, 2002). In the MnPO, the influence of

L-Type calcium channel and NO utilizing cGMP pathways on ANG II pressor effect can explain part of these results as has been demonstrated for the water intake (Zhu and Herbert, 1997a). An interaction between L-Type calcium channel and a free radical gas (NO) in the MnPO on the control of ANG II-induced pressor effect acting primarily in AT_1 and secondary in AT_2 receptors, respectively may be postulated. Also we may infer that ANGII act in AT_1 receptor opening calcium channels and inhibit the NO release.

ACKNOWLEDGEMENTS

This study was supported by CNPq (520408/96.9), FAPESP (99/06582-2 and 02/03806-1), FUNDUNESP (01/06756-3), PRONEX and FUNADESP-UNIARA.

REFERENCES

- Budzikowski, A.S. and F.H. Leenen, 2001. ANG II in median preoptic nucleus and pressor responses to CSF sodium and high sodium intake in SHR. *Am. J. Physiol.*, 281: H1210-H1216.
- Bumashev, N., 1996. Calcium permeability of glutamate-gated channels in the central nervous system. *Curr. Opin. Neurobiol.*, 6: 311-317.
- Bush, M.A. and G.M. Pollack, 2001. Pharmacokinetics and pharmacodynamics of 7-nitroindazole, a selective nitric oxide synthase inhibitor, in the rat hippocampus. *Pharm. Res.*, 18: 1607-1601.
- Cunningham, J.T., S.B. Bruno, R.R. Grindstaff, K.H. Higgs, D. Mazzella and M.J. Sullivan, 2002. Cardiovascular regulation of supraoptic vasopressin neurons. *Progress Brain Res.*, 139: 257-273.
- Dawson, T.M. and S.K. Snyder, 1994. Gases as biological messengers: Nitric oxide and carbon monoxide in the brain. *J. Neurosci.*, 14: 5147-5159.
- Felix, D., M.C. Khosla, K.L. Barnes, H. Imboden, B. Montani and C.M. Ferrario, 1991. Neurophysiological responses to angiotensin (1-7). *Hypertension*, 17: 1111-1114.
- Gardiner, T.W. and E.M. Stricker, 1985. Impaired drinking responses of rats with lesions of nucleus medianus: Circadian dependence. *Am. J. Physiol.*, 248: R224-R230.
- Garthwaite, J., 1991. Glutamate, nitric oxide and cell-cell signaling in the nervous system. *Trends Neurosci.*, 14: 60-67.
- Ghosh, A. and M.E. Greenberg, 1995. Calcium signaling in neurons: Molecular mechanisms and cellular consequences. *Science*, 268: 239-247.

- Hodge, G., V.Z. Ye and K.A. Duggan, 2002. Salt-sensitive hypertension resulting from nitric oxide synthase inhibition is associated with loss of regulation of angiotensin II in the rat. *Exp. Physiol.*, 87: 1-8.
- Mangiapane, M.L., T.N. Trasher, L.C. Keil, J.B. Simpson and W.F. Ganong, 1983. Deficits in drinking and vasopressin secretion after lesions of the nucleus medianus. *Neuroendocrinol.*, 37: 73-77.
- McKinley, M.J., A.L. Albiston, A.M. Allen, C.N. May, R.M. McAllen, B.J. Oldfield, F.A. Mendelsohn and S.Y. Chai, 2003. The brain rennin-angiotensin system: Location and Physiological roles. *Intl. J. Biochem. Cellular Biol.*, 35: 901-918.
- Miller, R.J., 1998. Calcium signaling in neurons. *Trends Neurosci.*, 11: 415-419.
- Naees, P.A., K.A. Kirkebeen, G. Cristensen and F. Kiil, 1992. Inhibition of renal nitric oxide synthesis with MG-monomethyl-L-arginine and NG-nitro-L-arginine. *Am. J. Physiol.*, 262: F939-F945.
- O'neill, T.P. and M.J. Brody, 1987. Role of the median preoptic nucleus in centrally evoked pressor responses. *Am. J. Physiol.*, 252: R1165-R1172.
- Paxinos, G. and C. Watson, 1986. *The Rat Brain in Stereotaxic Coordinates*, New York: Academic Press.
- Reid, J.A., 1988. Actions of angiotensin II on the brain: Mechanisms and physiological role. *Am. J. Physiol.*, 246: F533-F543.
- Saad, W.A., L.A.A. Camargo, A.F. Pereira and S. Simões, 1999. Effect of injection of L-NAME on drinking response. *Braz. J. Med. Biol. Res.*, 32: P1413-P1416.
- Saad, W.A., L.A.A. Camargo, I.F.M.S. Guarda, T.A.F. Santos, R.S. Guarda, W.A. Saad, S. Simões and R.J. Antunes, 2002a. Interaction between supraoptic nucleus and septal area in the control of water, sodium intake and arterial blood pressure induced by injection of angiotensin II. *Pharmacol. Biochem. Beh.*, 77: 667-674.
- Saad, W.A., I.F.M.S. Guarda, R.S. Guarda, L.A.A. Camargo, T.A.F.B. Santos, W.A. Saad and S. Simões, 2002b. Role of nitric oxide and beta receptors of the central nervous system on the salivary flow induced by pilocarpine injection into the lateral ventricle. *Pharmacol. Biochem. Beh.*, 72: 229-235.
- Saad, W.A., I.F.M.S. Guarda, L.A.A. Camargo, T.A.F.B. Santos, R.S. Guarda, S. Simões, W.A. Saad and R.J. Antunes, 2003. Role of nitric oxide of the median preoptic nucleus (MnPO) on the alterations of salivary flow, arterial pressure and heart rate induced by injection of pilocarpine into MnPO and intraperitoneally. *Braz. J. Med. Biol. Res.*, 36: 897-905.
- Saad, W.A., L.I. Gutierrez, I.F.M.S. Guarda, T.A.F.B. Santos, W.A. Saad, S. Simões and R.S. Guarda, 2004a. Nitric oxide of the supraoptic nucleus influences the salivary secretion, sodium renal excretion, urinary volume and arterial blood pressure. *Life Sci.*, 74: 1593-1603.
- Saad, W.A., L.I. Gutierrez, I.F.S.M. Guarda, L.A.A. Camargo, T.A.F.B. Santos, W.A. Saad, S. Simões and R.S. Guarda, 2004b. Lateral hypothalamus lesions influences water and salt intake and sodium and urine excretion, arterial blood pressure induced by L-NAME and FK 409 injections into median preoptic nucleus in conscious rats. *Life Sci.*, 75: 685-697.
- Saad, W.A., I.F.M.S. Guarda, L.A.A. Camargo, R.S. Guarda, W.A. Saad and T.A.F.B. Santos, 2006. Influence of vasopressin receptor and nitric oxide on the water, sodium intake and arterial blood pressure induced by angiotensin injected into the third ventricle of the rat brain. *J. Biol. Sci.*, 6: 182-86.
- Summy-Long, J.Y., V. Bui, S. Gest and M. Kadekaro, 2002. Nitric oxide, interleukin and prostaglandin interactions affecting the magnocellular system. *Brain Res.*, 940: 10-20.
- Thunhorst, R.T. and A.K. Johnson, 1994. Renin-angiotensin, arterial pressure and salt appetite in rats. *Am. J. Physiol.*, 266: R458-R465.
- Tsien, R.W., D. Lipscombe, D.V. Madison, K.R. Bley and A.P. Fox, 1988. Multiple types of neuronal calcium channels and their selective modulation. *Trends Neurosci.*, 11: 431-38.
- Wu, S.Y., S.L. Dun, V. Forstmann and N.J. Dun, 1997. Nitric oxide and excitatory post-synaptic currents in immature rat sympathetic preganglionic neurons *in vitro*. *Neuroscience*, 79: 237-245.
- Yashuda, Y., K. Honda, H. Negoro, T. Higuchi, Y. Goto and S. Fukuda, 2000. The contribution of the median preoptic nucleus to renal sympathetic nerve activity increased by intracerebroventricular injection of hypertonic saline in rat. *Brain Res.*, 867: 107-114.
- Zhu, B. and J. Herbert, 1997a. Angiotensin II interacts with nitric oxide cyclic Gmp pathway in the central control of drinking behavior: Mapping with *c-fos* and NADPH-diaphorase. *Neuroscience*, 79: 543-553.
- Zhu, B. and J. Herbert, 1997b. Calcium channels mediate angiotensin II-induced drinking behaviour and *c-fos* expression in the brain. *Brain Res.*, 778: 206-214.