



Microbiology

Journal

ISSN 2153-0696



Academic
Journals Inc.

www.academicjournals.com

Bacteria Help Infants Digest Milk More Effectively Than Adults

Infants are more efficient at digesting and utilizing nutritional components of milk than adults due to a difference in the strains of bacteria that dominate their digestive tracts.

Researchers from the University of California, Davis, and Utah State University report on genomic analysis of these strains, in the November 2010 issue of the journal *Applied and Environmental Microbiology*, identifying the genes that are most likely responsible for this difference.

“Human milk *oligosaccharides* (HMOs) are the third-largest solid component of milk. Their structural complexity renders them non-digestible to the host,” say the researchers. “*Bifidobacterium longum* strains often predominate the colonic microbiota of exclusively breast-fed infants. Among the three recognized subspecies, *B. longum* subsp. *infantis* achieves high levels of cell growth on HMOs and is associated with early colonization of the infant gut.”

In the study, the researchers used whole-genome microarray comparisons to associate genotypic biomarkers among 15 *B. longum* strains exhibiting various HMO utilization patterns. They identified 5 distinct gene clusters on *B. longum* that were conserved (showed little or no variation) across all strains capable of growth on HMOs and have also diverged in strains incapable of growing on HMOs.

The results of this study suggest that *B. longum* has at least 2 distinct subspecies: *B. longum* subsp. *infantis*, adapted to utilize milk carbon and found primarily in the digestive tract of children, and *B. longum* subsp. *longum*, specialized for plant-derived carbon metabolism and associated with the adult digestive tract.

“Although early gut colonization is likely dependent on a multitude of dietary and nondietary factors, the delivery of complex *oligosaccharides* through milk creates an ideal and unique nutrient niche for the establishment of, and colonization by, *B. longum* subsp. *infantis* strains,” say the researchers. “During weaning, a gradual transitioning from milk-based to plant-based diets generates a shift in carbon availability in the gastrointestinal tract favorable for the expansion and formation of an adult-like *gastrointestinal tract microbiota*.”

Editor’s Note: This article is not intended to provide medical advice, diagnosis or treatment.