

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

Pakistan Journal of Biological Sciences

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Mutagenicity of Acridine and Ascorbic Acid in Rhizobia of Legume Trees

²M.A. Nassef, ¹ A.M. El-Adl, ¹K.A. Zaied, ²S.M. Abd El-Wahab and ²E.S. Ibrahim

¹ Department of Genetics, Faculty of Agriculture, Mansoura University, Egypt

² Institute of Water and Soil Research, Agricultural Research Center, Giza, Egypt

Abstract: The present study aimed to induce a great variation existing among *Rhizobium* spp. that nodulate *Leucaena leucocephala* and *Sesbania sesban*, through the treatment with acridine and ascorbic acid and selection of isolates for efficiency in the symbiotic association. Bacteria were isolated from nodules of both hosts growing in Egyptian soil, subcultured, and verified to be rhizobia. The isolates varied significantly in their resistance to antibiotics and NaCl, their growth on different carbohydrates, and their effect on the pH of culture media. Most isolates showed intermediate antibiotic resistance, the capacity to use numerous carbohydrates, and a neutral to acid pH response. The mean generation time of these isolates ranged from 4.813 to 6.437 and 4.600 to 6.523 for *Sesbania sesban* and *Leucaena leucocephala*, respectively. Both acridine orange and ascorbic acid had genotoxic hazards on all rhizobial strains examined here. They demonstrated a dose-response for decreasing cell survival at the levels that are not excessively toxic to bacteria. The standard assay with pre-incubation was quite toxic to the bacteria than plate incorporation test. Acridine orange and ascorbic acid shows an increase in number of auxotrophic mutants over the spontaneous value which is evidence for their mutagenicity caused by DNA damage. The biochemical mutants obtained in this study were identified using nine plates of minimal medium, each supplemented with different combinations of four growth factors. From the results reported here, it can be concluded that acridine mutagenesis is due to an enhancement of mismatch repair. In addition, ascorbic acid may be mutagenic and cytotoxic through the generation of hydrogen peroxide.

Key words: Acridine, antibiotic resistance, ascorbic acid, auxotrophic mutant, generation time, mutation frequency, mutation rate, *Rhizobium*, salt stress

Introduction

Successful nodulation and nitrogen fixation by legumes inoculated with *Rhizobium* strains depend on genetic and environmental factors. Once introduced, *Rhizobium* bacteria must multiply in the rhizosphere and infect their host plant to initiate the symbiotic process. Success depends on competitiveness, ability to survive in the soil, and incompatibility with the host (Dowling and Broughton, 1986). Many effective strains have been reported to be poor competitors and may, consequently, form very few nodules in the presence of ineffective strains (Trinick, 1982). Sensitivity to antibiotics varies between species and it has been suggested that such variation may be a useful taxonomic character (Graham, 1963). A potential problem with the use of antibiotic-resistant mutant strains for ecological studies is that they may vary in other important characteristics, including competitiveness for nodule formation and ability to fix nitrogen. Therefore, it is very important to compare resistant strains with wild-type strains if the genetically marked strains are to be used in experiments designed to draw inferences concerning the ecology of the parental strains.

Recently, Thomas and Macphée (1987) have reported that acridine-induced frame shift mutagenesis might be subject to glucose repression. Glucose indirectly controls the expression of a number of catabolic operons (e.g., Lac, gal, ara) by lowering the intracellular concentration of cyclic AMP (cAMP). Pons and Müller (1989) had shown that upon treatment with 9-aminoacridine (9AA) in non-growth medium of long-phase cells of *E. coli* K12/343/113 (Mohn and Ellenberger, 1977), there was an increase in *nad*-113 ~ *Nad*⁺ reversion of more than 5 orders of magnitude. Using this sensitive frame shift marker, Pons and Müller (1989) showed that the glucose effect in acridine-induced mutagenesis is actually an "energy effect" and that it is due to mismatch repair. *In vitro*, mutagenesis bioassays are used to identify environmental chemicals that are potential genotoxic hazards for man. Recent reports have indicated that ascorbic acid, at high concentrations, may be mutagenic (Galloway and Painter, 1979) and cytotoxic in these *in vitro* test systems. L-ascorbic acid (vitamin C) has been found to be capable of modifying the properties of DNA (McCarty, 1945). Ascorbate can inactivate several phages by induction of single-strand breaks in their DNA (Murata *et al.*, 1986), and can induce multiple breaks in the chromosomes of repair-deficient

mutants of *E. coli* (Van Sluys *et al.*, 1986). In this study, the genetic diversity of legume trees rhizobia concerning sensitivity to antibiotics and tolerance to salt, as well as, the mutagenic effect of acridine and ascorbic acid in different *Rhizobium* strains were determined.

Materials and Methods

Bacterial strains and cultures: *Rhizobium* strains are listed in Table 1. Tryptone-yeast extract medium (TY) was used as a complete medium to ensure the independence of mutations according to Beringer (1974). Liquid growth medium (VS) was used to allow expression of mutations according to Pain (1979). Minimal medium (MM) was used according to Balasa (1963) and Murphy and Elkan (1963) for mutants isolation and identification through supplemented with a suitable concentration of the appropriate requirements. Mineral medium was also used for carbon source utilization test according to Milnitsky *et al.* (1997) by replaced mannitol with the appropriate carbon source at a final concentration of 1% (w/v). Yeast extract mannitol medium (YEM) was used for culture maintenance according to Allen (1959). YEM with Congo-red containing 10 ml of Congo-red (0.25g of Congo-red in 100 ml DW) were added to each liter of YEM medium according to Vincent (1970). YEM with bromo-thymol blue (BTB) was incorporated into agar medium at the rate of 5 ml of 0.5% alcoholic solution per liter. BM₁ medium was used as a basal medium, in this medium phosphates were autoclaved separately. In addition, BM₁-mannitol medium was prepared by adding 1% mannitol (Lafreniere *et al.*, 1987). pH was adjusted to 6.7 and 2% agar was added with 1.5-2.0% to solidify the medium. So, BM₂ medium was similar to BM₁, but additionally buffered to pH 7 according to Lafreniere *et al.* (1987).

Mutagenic agents:

Acridines (phase-shift mutations): The effect of acridines would be the specific induction of A-T transversion (C-G to A-T, or A-T to T-A). The mode of action of acridine orange, used in this study was production of frame shifts and it was the product from Sigma Chemical Company, USA. Stock solution of acridine orange was prepared in phosphate buffer (pH 7.5; 10mmol/L) and filter sterilized according to Pandey and Kashyap (1992).

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

Table 1: *Rhizobium* isolate strains from different locations in Egypt

| Legume host (source of rhizobia) | Location | Designation |
|-------------------------------------|--------------------------------------|---------------------|
| <i>Sesbania sesban</i> | Far. Fac. of Science, Mansoura Univ. | FFSMU ₁ |
| <i>Sesbania sesban</i> | Far. Fac. of Agric., Mansoura Univ. | FAFMU ₂ |
| <i>Sesbania sesban</i> | Far. Fac. of Agric., Mansoura Univ. | FAFMU ₃ |
| <i>Sesbania sesban</i> | Far. Fac. of Agric., Mansoura Univ. | FAFMU ₄ |
| <i>Sesbania sesban</i> | Far. Fac. of Agric., Mansoura Univ. | FAFMU ₅ |
| <i>Sesbania sesban</i> | Far. Fac. of Agric., Mansoura Univ. | FAFMU ₆ |
| <i>Sesbania sesban</i> | Fac. of pharmacy., Mansoura Univ. | FPMU ₇ |
| <i>Sesbania sesban</i> | Fac. of Art., Minia Univ. | FFAMU ₈ |
| <i>Sesbania sesban</i> | Agric., Res. Center, Giza | ARCG ₉ |
| <i>Sesbania sesban</i> | Agric., Res. Center, Giza | ARCG ₁₀ |
| <i>Sesbania sesban</i> | Agric., Res. Center, Giza | ARCG ₁₁ |
| <i>Sesbania sesban</i> | Agric., Res. Center, Giza | ARCG ₁₂ |
| <i>Sesbania sesban</i> | Sakha Agric. Res. Station | SARS ₁₃ |
| <i>Sesbania sesban</i> | Sakha Agric. Res. Station | SARS ₁₄ |
| <i>Sesbania sesban</i> | Sakha Agric. Res. Station | SARS ₁₅ |
| <i>Leucaena leucocephala</i> , Lam. | National Res. Center, Dokki | NRC ₁₆ |
| <i>Leucaena leucocephala</i> , Lam. | National Res. Center, Dokki | NRC ₁₇ |
| <i>Leucaena leucocephala</i> , Lam. | National Res. Center, Dokki | NRC ₁₈ |
| <i>Leucaena leucocephala</i> , Lam. | National Res. Center, Dokki | NRC ₁₉ |
| <i>Leucaena leucocephala</i> , Lam. | Far. Fac. of Art, Minia Univ. | FFAMU ₂₀ |
| <i>Leucaena leucocephala</i> , Lam. | Far. Fac. of Art, Minia Univ. | FFAMU ₂₁ |
| <i>Leucaena leucocephala</i> , Lam. | Far. Fac. of Art, Minia Univ. | FFAMU ₂₂ |
| <i>Leucaena leucocephala</i> , Lam. | Far. Fac. of Art, Minia Univ. | FFAMU ₂₃ |
| <i>Leucaena leucocephala</i> , Lam. | Far. Fac. of Art, Minia Univ. | FFAMU ₂₄ |
| <i>Leucaena leucocephala</i> , Lam. | Ministry of Agric., Minia Govern. | MAMG ₂₅ |
| <i>Leucaena leucocephala</i> , Lam. | Ministry of Agric., Minia Govern. | MAMG ₂₆ |
| <i>Leucaena leucocephala</i> , Lam. | Hort. Res. Inst., Agric. Res. Center | HRI ₂₇ |
| <i>Leucaena leucocephala</i> , Lam. | Hort. Res. Inst., Agric. Res. Center | HRI ₂₈ |
| <i>Leucaena leucocephala</i> , Lam. | Hort. Res. Inst., Agric. Res. Center | HRI ₂₉ |
| <i>Leucaena leucocephala</i> , Lam. | Hort. Res. Inst., Agric. Res. Center | HRI ₃₀ |

Far. = Farm.

Table 2: Combination of supplements in minimal agar plates.

| | 1 | 2 | 3 | 4 | 5 |
|---|-----------------|-------------|-----------------|------------|---------------|
| 6 | Alanine | L-arginine | L-aspartic acid | L-cystein | Glycine |
| 7 | L-Glutamic acid | L-histidine | L-leucine | L-ysine | DL-methionine |
| 8 | L-Phenylalanine | Proline | L-tryptophan | L-tyrosine | DL-valine |
| 9 | Adenine | uracil | Biotin | Pyridoxine | Thiamine |

All supplements are at a final concentration of 20 µg/ml.

Table 3: Some phenotypic characteristics of *Rhizobium* isolates nodulating *Sesbania sesban*.

| Isolates | pH reaction | Growth at pH range | Mean generation time (h) | Opacity | Colony morphology | Size (mm) |
|----------|-------------|--------------------|--------------------------|------------------|-------------------|-----------|
| FFSMU-1 | Acidic | 5-11 | 6.090 | Opaque | High | 2.6 |
| FAFMU-2 | Acidic | 5-10 | 5.363 | Opaque | High | 5.3 |
| FAFMU-3 | Acidic | 5-10 | 5.467 | Opaque | High | 5.4 |
| FAFMU-4 | Acidic | 5-10 | 5.940 | Opaque | Medium | 4.8 |
| FAFMU-5 | Acidic | 5-11 | 5.277 | Opaque | Low | 4.4 |
| FAFMU-6 | Acidic | 5-10 | 6.437 | Translucent | Low | 4.2 |
| FPMU-7 | Acidic | 4-10 | 6.200 | Translucent | Low | 5.2 |
| FFAMU-8 | Acidic | 5-10 | 5.480 | Semi-translucent | Low | 2.2 |
| ARCG-9 | Neutral | 4-10 | 6.120 | Semi-translucent | Low | 6.8 |
| ARCG-10 | Acidic | 5-10 | 6.373 | Semi-translucent | Low | 4.1 |
| ARCG-11 | Acidic | 5-11 | 5.387 | Opaque | High | 4.6 |
| ARCG-12 | Acidic | 5-10 | 5.710 | Translucent | Medium | 4.8 |
| SARS-13 | Neutral | 5-11 | 4.813 | Translucent | High | 3.6 |
| SARS-14 | Neutral | 5-11 | 5.097 | Translucent | Low | 4.4 |
| SARS-15 | Acidic | 4-10 | 5.170 | Translucent | Medium | 3.0 |

Ascorbic acid (Vitamin C): L-ascorbic acid has the simplest chemical structure of all the vitamins. It is a reasonably strong reducing agent. The biochemical and physiological functions of ascorbic acid most likely derive from its reducing properties. It functions as an electron carrier. Loss of one electron due to interaction with oxygen or metal ions leads to semidehydro-L-ascorbate, a reactive free radical that can be reduced back to L-ascorbic acid by various enzymes in animals and plants. L-ascorbic acid, product from Sigma, Co was used in the present study to investigate the mutagenic activity in rhizobia.

Isolation and purification of *Rhizobium* isolates: The isolates of rhizobia listed in Table 1 were isolated from root nodules of *Sesbania sesban*, *Leucaena leucocephala* (Lam.) according to Chan

et al. (1988).

Cultural properties, colony morphology and pH reaction: Cultures were prepared by inoculating about 50 ml of yeast extract mannitol liquid medium (YEM) (Vincent, 1970) with a loopful of each of the stored cultures. The flasks were incubated at 28°C with 140 rev min⁻¹ for three days. During that time, growth rate was determined (Fitzsimon *et al.*, 1992). YEMA medium containing congo red was used for colony morphology characterization. YEMA medium containing 25 mg bromothymol blue (BTB) L⁻¹ was used for detecting pH changes. A yellow, blue and green colours indicated acidic, alkaline and neutral characters respectively, according to Norris (1965).

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

Table 4: Some phenotypic characteristics of *Rhizobium* isolates nodulating *Leucaena leucocephala*.

| Isolates | pH reaction | Growth at pH range | Mean generation time (h) | Opacity | Colony morphology | viscosity | Size (mm) |
|----------|-------------|--------------------|--------------------------|------------------|-------------------|-----------|-----------|
| NRC-16 | Acidic | 5-10 | 5.047 | Translucent | Low | | 4.10 |
| NRC-17 | Acidic | 5-10 | 5.357 | Translucent | Low | | 2.50 |
| NRC-18 | Acidic | 4-10 | 6.397 | Translucent | Low | | 5.00 |
| NRC-19 | Acidic | 5-11 | 5.287 | Semi Translucent | Low | | 5.20 |
| FFAMU-20 | Acidic | 5-10 | 5.043 | Semi Translucent | High | | 5.00 |
| FFAMU-21 | Acidic | 5-10 | 4.600 | Semi Translucent | Low | | 4.40 |
| FFAMU-22 | Acidic | 5-10 | 5.680 | Semi Translucent | High | | 5.20 |
| FFAMU-23 | Acidic | 4-11 | 5.267 | Opaque | Medium | | 4.30 |
| FFAMU-24 | Acidic | 4-10 | 5.040 | Opaque | Medium | | 4.10 |
| MAMG-25 | Acidic | 5-10 | 6.523 | Opaque | High | | 5.20 |
| MAMG-26 | Acidic | 5-10 | 5.457 | Opaque | High | | 6.20 |
| HRI-27 | Acidic | 5-10 | 5.097 | Translucent | High | | 5.90 |
| HRI-28 | Acidic | 5-11 | 5.570 | Translucent | High | | 6.90 |
| HRI-29 | Acidic | 5-11 | 6.053 | Translucent | Medium | | 5.80 |
| HRI-30 | Acidic | 4-10 | 5.340 | Translucent | Medium | | 3.80 |

Table 5: Mean generation time (h) of *Sesbania sesban* - rhizobia under salt stress at varying degrees.

| Isolates No. | Concentration of NaCl (%) | | | | | Relative sensitivity |
|--------------|---------------------------|--------|----|----|----|----------------------|
| | 0.01 | 3 | 4 | 5 | 6 | |
| FFSMU-1 | 6.090 | 9.366 | NG | NG | NG | 1.54 |
| FAFMU-2 | 5.363 | 12.606 | NG | NG | NG | 2.35 |
| FAFMU-3 | 5.466 | 9.346 | NG | NG | NG | 1.71 |
| FAFMU-4 | 5.940 | 8.346 | NG | NG | NG | 1.41 |
| FAFMU-5 | 5.276 | 7.380 | NG | NG | NG | 1.40 |
| FAFMU-6 | 6.436 | 9.86 | NG | NG | NG | 1.53 |
| FPMU-7 | 6.200 | 13.33 | NG | NG | NG | 2.15 |
| FFAMU-8 | 5.480 | 17.21 | NG | NG | NG | 3.14 |
| ARCG-9 | 6.120 | 12.05 | NG | NG | NG | 1.97 |
| ARCG-10 | 6.373 | 12.70 | NG | NG | NG | 1.99 |
| ARCG-11 | 5.386 | 14.446 | NG | NG | NG | 2.68 |
| ARCG-12 | 5.710 | 13.606 | NG | NG | NG | 2.38 |
| SARS-13 | 4.813 | NG | NG | NG | NG | 1.00 |
| SARS-14 | 5.096 | NG | NG | NG | NG | 1.00 |
| SARS-15 | 5.170 | NG | NG | NG | NG | 1.00 |
| F-test | ** | ** | | | | ** |
| L.S.D. 0.05 | 0.698 | 2.287 | | | | 0.38 |
| 0.01 | 0.940 | 3.079 | | | | 0.52 |

NG = No growth.

Table 6: Mean generation time (h) of *Leucaena leucocephala* - rhizobia under salt stress at varying degrees.

| Isolates No. | Concentration of NaCl (%) | | | | | Relative sensitivity |
|--------------|---------------------------|--------|--------|----|----|----------------------|
| | 0.01 | 3 | 4 | 5 | 6 | |
| NRC-16 | 5.046 | 12.936 | 40.336 | NG | NG | 7.99 |
| NRC-17 | 5.356 | 9.863 | 59.303 | NG | NG | 11.07 |
| NRC-18 | 6.396 | NG | NG | NG | NG | 1.00 |
| NRC-19 | 5.286 | 11.590 | NG | NG | NG | 2.19 |
| FFAMU-20 | 5.073 | 14.833 | NG | NG | NG | 2.923 |
| FFAMU-21 | 4.533 | 20.300 | NG | NG | NG | 4.02 |
| FFAMU-22 | 5.680 | 61.243 | NG | NG | NG | 10.43 |
| FFAMU-23 | 5.866 | 20.300 | 32.55 | NG | NG | 5.54 |
| FFAMU-24 | 5.040 | 19.240 | 26.72 | NG | NG | 5.30 |
| MAMG-25 | 6.523 | 27.543 | NG | NG | NG | 4.22 |
| MAMG-26 | 5.456 | 29.426 | NG | NG | NG | 5.39 |
| HRI-27 | 5.096 | 30.493 | NG | NG | NG | 5.24 |
| HRI-28 | 5.570 | 33.460 | NG | NG | NG | 5.95 |
| HRI-29 | 6.053 | 45.493 | NG | NG | NG | 7.52 |
| HRI-30 | 5.340 | 40.190 | NG | NG | NG | 6.86 |
| F-test | ** | ** | ** | | | ** |
| L.S.D. 0.05 | 0.522 | 8.100 | 6.574 | | | 1.89 |
| 0.01 | 0.702 | 10.906 | 8.850 | | | 2.55 |

NG = No growth.

Carbon source utilization: Cells were cultured in liquid mineral medium, collected by centrifugation at 7000 x g for 5 min at 4°C, washed twice with 20mM sodium phosphate buffer, pH 7.0, containing 1.50mM NaCl and 3mM KCl, and finally resuspended in mineral medium. After two hours, 20µL of liquid mineral medium were spotted on the surface of the same solid mineral medium plates and then incubated at 28°C for 4 days. Appropriate carbon source at a final concentration of 1% (w/v) was used according to

Milnitsky *et al.* (1997).

Tolerance to salinity of NaCl: Yeast extract agar (YEMA) plates were prepared for NaCl sensitivity. Plates, in three replications, were plated with 0.1 ml of a suitable dilution of an overnight broth culture amended with NaCl concentrations from 3 to 6% (w/v). Plates were incubated at 28°C for three days. All plates were visually scored for bacterial growth as separated colonies at zero

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

Table 7: Phenotypic evaluation of *Rhizobium* sp. isolated from *Sesbania sesban* nodules.

| Carbon sources | Isolates | | | | | | | | | | | | | | |
|--------------------------------|----------|---------|---------|---------|---------|---------|---------|---------|--------|---------|---------|---------|---------|---------|---------|
| | FFSMU 1 | FAFMU 2 | FAFMU 3 | FAFMU 4 | FAFMU 5 | FAFMU 6 | FAFMU 7 | FAFMU 8 | ARCG 9 | ARCG 10 | ARCG 11 | ARCG 12 | SARS 13 | SARS 14 | SARS 15 |
| Pentose: | | | | | | | | | | | | | | | |
| L- arabinose | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + |
| D- arabinose | + | + | + | + | + | + | - | + | + | + | + | + | + | + | + |
| Xylose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Hexoses: | | | | | | | | | | | | | | | |
| D- fructose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| D- galactose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| D- glucose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Mannose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Disaccharide: | | | | | | | | | | | | | | | |
| Lactose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Maltose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Sucrose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Sugar alcohol: | | | | | | | | | | | | | | | |
| Mannitol | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| C₃ compound: | | | | | | | | | | | | | | | |
| Glycerol | - | + | + | + | + | + | - | + | + | + | + | + | + | + | + |
| Polysaccharide: | | | | | | | | | | | | | | | |
| Dextrin | - | - | - | - | + | + | + | - | - | - | - | - | - | - | - |
| Dextrose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Diastase | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + |
| Inositol | + | + | + | - | + | + | + | + | + | + | + | + | + | + | + |

+ = Visible growth - = Invisible growth.

Table 8: Phenotypic evaluation of *Rhizobium* sp. isolated from the nodules of *Leucaena leucocephala*, Lam

| | Isolates | | | | | | | | | | | | | | | |
|---|----------|--------|--------|--------|----------|----------|----------|----------|----------|----------|---------|---------|--------|--------|--------|--|
| | NRC-16 | NRC-17 | NRC-18 | NRC-19 | FFAMU-20 | FFAMU-21 | FFAMU-22 | FFAMU-23 | FFAMU-24 | FFAMU-25 | MAMG-26 | MAMG-27 | HRI-28 | HRI-29 | HRI-30 | |
| Carbon sources | | | | | | | | | | | | | | | | |
| Pentose: | | | | | | | | | | | | | | | | |
| L- arabinose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| D- arabinose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Xylose | + | + | + | + | + | - | - | + | + | + | + | + | + | + | + | |
| Hexoses: | | | | | | | | | | | | | | | | |
| D-fructose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| D-galactose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| D-glucose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Mannose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Disaccharide: | | | | | | | | | | | | | | | | |
| Lactose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Maltose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sucrose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Sugar alcohol: | | | | | | | | | | | | | | | | |
| Mannitol | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| C₃ compound: | | | | | | | | | | | | | | | | |
| Glycerol | + | + | + | + | + | - | + | - | + | + | - | + | + | + | + | |
| Polysaccharide: | | | | | | | | | | | | | | | | |
| Dextrin | + | - | - | - | + | + | + | - | - | - | - | - | - | - | + | |
| Dextrose | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Diastase | + | + | + | + | + | + | + | + | + | + | + | + | + | + | + | |
| Inositol | + | + | + | + | + | - | + | + | + | + | - | + | + | + | + | |
| + = Visible growth - = Invisible growth. | | | | | | | | | | | | | | | | |

+ = Visible growth - = Invisible growth.

Table 9: Minimum inhibitory concentration ($\mu\text{g ml}^{-1}$) of nine antibiotics against *Rhizobium* isolates of *Sesbania sesban*.

| Isolates | Tetracycline | Ampicillin | Neomycin | Kanamycin | Spectomycin | Rifampcin | Pencillin-G | Chloramphenicol | Streptomycin |
|----------|--------------|------------|----------|-----------|-------------|-----------|-------------|-----------------|--------------|
| FFAMU-1 | 15 | 150 | 75 | 100 | 100 | 50 | 275 | 100 | 150 |
| FFAMU-2 | 15 | 100 | 75 | 45 | 100 | 100 | 460 | 75 | 75 |
| FFAMU-3 | 5 | 75 | 75 | 50 | 150 | 75 | 350 | 75 | 125 |
| FFAMU-4 | 20 | 100 | 150 | 150 | 175 | 75 | 400 | 125 | 100 |
| FFAMU-5 | 10 | 125 | 150 | 75 | 100 | 50 | 400 | 175 | 150 |
| FFAMU-6 | 5 | 75 | 75 | 75 | 100 | 50 | 400 | 300 | 150 |
| FFAMU-7 | 5 | 75 | 150 | 100 | 175 | 75 | 275 | 150 | 150 |
| FFAMU-8 | 15 | 75 | 75 | 100 | 100 | 100 | 200 | 50 | 100 |
| ARCG-9 | 20 | 75 | 75 | 125 | 100 | 75 | 125 | 100 | 150 |
| ARCG-10 | 10 | 125 | 100 | 75 | 100 | 50 | 400 | 275 | 200 |
| ARCG-11 | 10 | 100 | 50 | 100 | 100 | 50 | 150 | 225 | 75 |
| ARCG-12 | 25 | 75 | 150 | 125 | 100 | 50 | 300 | 225 | 125 |
| SARS-13 | 10 | 75 | 50 | 50 | 50 | 75 | 300 | 125 | 100 |
| SARS -14 | 15 | 50 | 100 | 100 | 100 | 75 | 300 | 250 | 175 |
| SARS -15 | 15 | 75 | 150 | 125 | 75 | 75 | 325 | 225 | 150 |

time and after 24, 36, 72 and 96 hour of incubation. The results were carefully recorded, keeping each part separate.

Growth characteristics: Growth characterization was done with mineral medium for salinity tolerance and VS medium in mutagenicity experiment, as the growth media. Exponential rates

were calculated from the bacterial concentrations x_0 , x_1 , x_2 , x_3 and x_4 at the following times, respectively t_0 (zero time), t_1 (24 h), t_2 (36 h), t_3 (72 h) and t_4 (96 h) with the equation:

$$\mu = (\log x_1 - \log x_0) / \log_e (t-t_0), \quad \text{MGT} = \frac{\ln_2}{\mu}$$

Table 10: Minimum inhibitory concentration ($\mu\text{g ml}^{-1}$) of nine antibiotics against *Rhizobium* isolates of *Leucaena leucocephala*.

| Isolates | Tetracycline | Ampicillin | Neomycin | Kanamycin | Spectomycin | Rifamycin | Pencillin-G | Chloramphenicol | Streptomycin |
|----------|--------------|------------|----------|-----------|-------------|-----------|-------------|-----------------|--------------|
| NRC-16 | 15 | 150 | 100 | 50 | 100 | 75 | 300 | 125 | 125 |
| NRC-17 | 20 | 75 | 50 | 75 | 100 | 100 | 300 | 150 | 150 |
| NRC-18 | 10 | 75 | 125 | 50 | 100 | 100 | 125 | 150 | 125 |
| NRC-19 | 10 | 100 | 125 | 75 | 100 | 150 | 375 | 75 | 125 |
| FFAMU-20 | 10 | 50 | 125 | 75 | 100 | 50 | 350 | 300 | 75 |
| FFAMU-21 | 15 | 50 | 50 | 50 | 100 | 75 | 325 | 225 | 100 |
| FFAMU-22 | 20 | 50 | 125 | 75 | 175 | 75 | 175 | 100 | 100 |
| FFAMU-23 | 10 | 75 | 100 | 50 | 125 | 75 | 250 | 100 | 100 |
| FFAMU-24 | 20 | 150 | 125 | 100 | 100 | 75 | 250 | 100 | 75 |
| MAMG-25 | 25 | 150 | 150 | 150 | 125 | 75 | 175 | 150 | 100 |
| MAMG-26 | 15 | 100 | 150 | 125 | 125 | 125 | 200 | 100 | 125 |
| HRI-27 | 25 | 100 | 100 | 100 | 200 | 150 | 400 | 100 | 125 |
| HRI-28 | 10 | 75 | 75 | 100 | 175 | 75 | 200 | 125 | 100 |
| HRI-29 | 10 | 25 | 75 | 100 | 100 | 125 | 325 | 125 | 75 |
| HRI-30 | 10 | 150 | 150 | 75 | 200 | 125 | 350 | 150 | 125 |

Where, $\text{Log}_e = 0.43429$, $\text{Ln}_2 = 0.69314$ (Schlegel, 1985).

The log phase was defined as the time between inoculation and establishment of the maximum growth rate (Schlegel, 1985) and was determined graphically by a plot of log cell number versus time.

Intrinsic antibiotic resistance (IAR): The ability of rhizobial strains to grow on nine different antibiotics at varying concentrations was assessed according to Chan *et al.* (1988). All antibiotics were obtained from Sigma Chemical Co. (St. Louis, Missouri, USA). Cultures were grown in VS broth to mid-log phase and then diluted to an optical density (OD) of 0.15 or 2×10^7 cells ml^{-1} (Turco *et al.*, 1986). Solutions of antibiotics were added to melted VS medium, which had been cooled to about 50°C . The minimum inhibitory concentration was determined by preparing plates with the previous series of antibiotic concentrations, spotting $10\mu\text{L}$ of cell suspension (2×10^7 cells ml^{-1}) on a sector of the plate, and incubating as described in Cole and Elkan (1973 & 1979). The minimum inhibitory concentration is defined as the lowest concentration of an antibiotic which completely inhibits the growth of an organism. The replica plates were incubated until colonies appeared on antibiotic-free plates.

Mutagenic action of acridine and ascorbic acid:

Starvation of bacteria: This method was used before the cells were treated with a mutagenic agents according to Thorne and Williams (1997).

Isolation of auxotrophic mutants: Mutagenesis of *Rhizobium* spp. was carried out by the treatment of the log phase cells grown on BM_1 -mannitol, with 60, 120 and 180 $\mu\text{g/ml}$ for both mutagens used in the present investigation and incubated for two hours at 28°C on a rotary shaker (160 rev. min^{-1}). The treated cells were washed twice with phosphate buffer saline (PBS; 3mM phosphate buffer in 0.7% NaCl, pH 6.8), suspended in liquid growth medium (VS) and incubated for 24 h at 28°C on a rotary shaker (160 rev. min^{-1}), to allow segregation and expression of mutation. For enrichment, the cells were washed twice in PBS and resuspended in BM_2 supplemented with the appropriate carbon source and 500 $\mu\text{g ml}^{-1}$ carbenicillin were added. After incubation at 28°C for 24 h on a rotary shaker, the enrichment cycle was repeated. Enriched cultures were washed and plated for single colonies on BM_2 medium modified (Lafreniere *et al.*, 1987) by using 0.6g L^{-1} peptone (Difco) as a nitrogen source and by adding filter sterilized 2, 3, 5-triphenyl tetrazolium chloride (TTC, Sigma) to a final concentration of 25mg L^{-1} . On this medium wild type colonies have a good growth and reduce TTC to triphenylformazan (TPF) (pink colonies) (Lenhard, 1956), while mutants show poor growth and TTC is not reduced (white colonies). Mutants were isolated and preserved at -69°C in yeast extract mannitol broth medium containing 10% glycerol (Lafontaine *et al.*, 1989). Survival was estimated by comparing with colony counts of untreated control according to Pandey and Kashyap (1992).

The mutation rate, shown previously, was derived from the formula:-

$$m = \log_e 2 (M_2 - M_1) / (N_2 - N_1)$$

Where, M_1 and M_2 are the number of mutant clones arising from plates sprayed at time 1 and 2, and N_1 and N_2 are the corresponding total bacterial counts (Beale, 1948). Mutation frequency was calculated according to Eckardt and Haynes (1977).

Identification of biochemical mutants of *Rhizobium*: Twenty possible requirements for single growth factors are tested. Nine plates of minimal medium, each supplemented with different combinations of six growth factors with the arrangement are shown in Table 2. A mutant strain with a single biochemical requirement when inoculated on all twelve plates will grow on two of them, one of plates 1-5 and one of plates 6-9 according to Holliday (1956).

Statistical analysis: Results were subjected to analysis of variance using the Statistical Analysis System (SAS). When analysis of variance showed significant treatment effects, the least significant difference (LSD) test was applied to make comparisons among the means at 0.05 and 0.01 levels of significance (Steel and Torrie, 1980).

Results and Discussion

Characterization of *Rhizobium* isolates: The isolates displayed a range of colony characteristics. The diameter of colonies of most isolates was 2.2 to 6.8 mm in *Rhizobium* of *Sesbania sesban* and 2.5 to 6.90 in *Rhizobium* of *Leucaena leucocephala*, and they were circular opaque to translucent. Colonies of a few isolates were greater than 5.2 mm in diameter, creamy white. Although, no quantitative data were collected, no correlation appeared to exist between colony form and isolate origin. The pH reaction of *Sesbania sesban* - *Rhizobium* isolates was ranged between acidic to neutral, in contrast, all the isolates of *Leucaena leucocephala* showed acidic pH reaction only. The collection of nodules and isolation of bacteria from many locations resulted in a collection of morphologically and biochemically diverse rhizobia that form symbiosis with leguminous trees. Differences among the rhizobial isolates shown here can not be ascribed to the genotype of the tree from which they were collected, because in some cases as much diversity was seen between isolates collected from one tree as between isolates collected from two different trees within one site. These differences were thought to be due to differences among rhizobial strains associated with each nodule (Batzli *et al.*, 1992). Variation also has been attributed to soil microsite characteristics such as aeration, nutrient availability, moisture content and competition (Postgate, 1982). These factors may explain variation among isolates in this study.

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

Table 11: Mutagenicity of acridine orange towards FFAMU-8 strain of *Sesbania sesban* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate ($\times 10^{-7}$) |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|---------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | |
| 0.0 | 6.6 | 100.0 | 0.0 | 0.0 | 0.0 | 55.0 | 100.00 | 0.003 | 0.006 | 0.40 | 0.00 |
| 60 | 4.3 | 24.66 | 0.28 | 6.49 | 6.43 | 25.0 | 45.45 | 1.20 | 4.80 | 4.20 | 3.07 |
| 120 | 2.7 | 48.36 | 0.50 | 20.93 | 20.54 | 5.40 | 9.81 | 1.48 | 27.47 | 26.86 | 19.31 |
| 180 | 1.5 | 30.71 | 0.23 | 15.0 | 15.00 | 2.30 | 4.30 | 0.70 | 24.73 | 29.13 | 41.07 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 0.53 | 8.70 | 0.13 | 9.09 | 8.44 | 1.20 | 1.89 | 0.17 | 6.14 | 11.32 |
| | 0.01 | 80.00 | 13.17 | 0.15 | 13.7 | 13.53 | 1.82 | 2.87 | 0.26 | 9.29 | 17.13 |

** Significant at 0.01 of probability level.

Table 12: Mutagenicity of acridine orange towards ARCG-10 strain of *Sesbania sesban* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate ($\times 10^{-7}$) |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|---------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | |
| 0.0 | 4.40 | 100.00 | 0.01 | 0.02 | 0.25 | 54.00 | 100.00 | 0.01 | 0.024 | 0.24 | 0.01 |
| 60 | 2.16 | 49.29 | 0.28 | 13.16 | 12.40 | 35.00 | 64.81 | 1.80 | 5.14 | 4.34 | 3.20 |
| 120 | 1.33 | 30.30 | 0.20 | 15.46 | 14.70 | 4.63 | 8.57 | 1.30 | 28.18 | 27.93 | 2.30 |
| 180 | 1.16 | 26.51 | 0.23 | 20.19 | 19.43 | 2.60 | 4.93 | 0.70 | 26.20 | 25.96 | 21.54 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 0.77 | 4.08 | 0.13 | 210.30 | 10.45 | 1.51 | 1.69 | 0.33 | 7.61 | 11.82 |
| | 0.01 | 1.17 | 6.18 | 0.19 | 15.60 | 15.83 | 2.27 | 2.56 | 0.50 | 11.52 | 17.90 |

** Significant at 0.01 of probability level.

Table 13: Mutagenicity of acridine orange towards HRI-27 strain of *Leucaena leucocephala* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate ($\times 10^{-7}$) |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|---------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | |
| 0.0 | 5.06 | 100.0 | 0.01 | 0.19 | 0.06 | 42.66 | 100.0 | 0.03 | 0.08 | 0.25 | 0.05 |
| 60 | 3.13 | 61.79 | 0.20 | 6.29 | 6.09 | 28.00 | 65.60 | 0.63 | 2.23 | 2.23 | 1.19 |
| 120 | 2.43 | 47.98 | 0.66 | 28.04 | 27.84 | 6.63 | 15.54 | 1.56 | 23.62 | 23.62 | 14.69 |
| 180 | 1.40 | 27.61 | 0.36 | 26.28 | 27.67 | 2.70 | 6.09 | 0.86 | 32.07 | 32.07 | 26.70 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 0.89 | 9.48 | 0.12 | 8.56 | 8.22 | 2.80 | 6.37 | 0.32 | 8.55 | 4.82 |
| | 0.01 | 1.36 | 14.35 | 0.67 | 12.95 | 12.44 | 4.24 | 9.65 | 0.49 | 12.95 | 7.31 |

** Significant at 0.01 of probability level.

Table 14: Mutagenicity of acridine orange towards NRC-19 strain of *Leucaena leucocephala* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate ($\times 10^{-7}$) |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|---------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | Total No. of colonies $\text{ml}^{-1} \times 10^6$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^6$ | Mutagenicity (%) | Mutation frequency $\times 10^6$ | |
| 0.0 | 4.66 | 100.0 | 0.003 | 0.073 | 0.05 | 55.67 | 100.00 | 0.03 | 0.05 | 0.01 | 0.04 |
| 60 | 3.50 | 74.99 | 0.50 | 13.26 | 13.14 | 12.33 | 22.45 | 1.26 | 10.15 | 10.09 | 6.19 |
| 120 | 2.40 | 51.43 | 0.63 | 26.69 | 26.22 | 6.63 | 10.11 | 1.60 | 28.56 | 29.23 | 23.50 |
| 180 | 1.030 | 22.14 | 0.23 | 23.33 | 23.26 | 2.10 | 3.77 | 0.70 | 33.31 | 33.25 | 30.66 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 0.49 | 7.53 | 0.22 | 11.46 | 11.45 | 1.97 | 1.26 | 0.13 | 5.74 | 11.46 |
| | 0.01 | 0.74 | 11.40 | 0.33 | 17.36 | 17.34 | 2.98 | 1.90 | 0.19 | 2.69 | 17.36 |

** Significant at 0.01 of probability level.

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

Table 15: Mutagenicity of ascorbic acid towards FFAMU-8 strain of *Sesbania sesban* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate $\times 10^{-7}$ |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|--------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | |
| 0.0 | 24.00 | 100.0 | 0.10 | 4.31 | 0.03 | 143.30 | 100.0 | 2.00 | 1.375 | 0.003 | 0.05 |
| 60 | 9.60 | 40.27 | 3.66 | 38.42 | 0.34 | 34.30 | 2.40 | 17.00 | 50.31 | 0.49 | 0.29 |
| 120 | 2.73 | 11.38 | 1.53 | 60.60 | 0.56 | 1.40 | 10.01 | 3.66 | 26.42 | 0.25 | 0.13 |
| 180 | 1.23 | 5.136 | 1.23 | 46.52 | 0.59 | 12.33 | 8.62 | 4.00 | 32.63 | 0.31 | 0.22 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 3.29 | 12.12 | 0.93 | 37.82 | 0.33 | 16.87 | 6.27 | 4.72 | 19.71 | 0.11 |
| | 0.01 | 4.99 | 18.35 | 1.41 | 57.27 | 0.42 | 25.53 | 9.49 | 7.15 | 29.84 | 0.17 |

** Significant at 0.01 of probability level.

Table 16: Mutagenicity of ascorbic acid towards ARCG-10 strain of *Sesbania sesban* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate $\times 10^{-7}$ |
|--|--|---------------|---|---------------------|--|--|---------------|--|---------------------|--|--------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutants $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | |
| 0.0 | 37.00 | 100.00 | 2.66 | 7.59 | 0.02 | 176.0 | 100.0 | 8.00 | 2.79 | 0.01 | 0.03 |
| 60 | 23.00 | 62.15 | 12.33 | 67.36 | 0.60 | 123.0 | 69.78 | 15.33 | 13.43 | 0.11 | 0.02 |
| 120 | 3.60 | 9.90 | 3.23 | 87.70 | 0.80 | 21.66 | 12.28 | 12.33 | 57.06 | 0.54 | 0.35 |
| 180 | 1.53 | 4.14 | 1.53 | 66.10 | 0.59 | 35.33 | 19.94 | 16.33 | 46.85 | 0.44 | 0.41 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 8.70 | 9.41 | 3.16 | 31.97 | 0.32 | 34.87 | 18.14 | 14.56 | 0.15 | 0.23 |
| | 0.01 | 13.18 | 14.25 | 4.78 | 48.42 | 0.49 | 52.80 | 27.47 | 22.05 | 0.22 | 0.34 |

** Significant at 0.01 of probability level.

Table 17: Mutagenicity of ascorbic acid towards NRC-19 strain of *Leucaena leucocephala* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate $\times 10^{-7}$ |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|--------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | |
| 0.0 | 33.66 | 100.00 | 0.00 | 0.00 | 0.00 | 133.33 | 100.0 | 0.00 | 0.00 | 0.00 | 0.00 |
| 60 | 11.00 | 32.67 | 6.00 | 53.70 | 0.53 | 25.66 | 19.25 | 3.33 | 11.42 | 0.53 | 0.38 |
| 120 | 5.63 | 16.72 | 3.53 | 61.54 | 0.61 | 17.66 | 13.25 | 12.33 | 70.77 | 0.70 | 0.53 |
| 180 | 1.47 | 5.24 | 1.48 | 82.18 | 0.81 | 2.66 | 7.25 | 5.66 | 59.84 | 0.59 | 0.74 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 4.28 | 6.16 | 2.05 | 15.18 | 0.15 | 18.83 | 5.18 | 14.84 | 0.15 | 0.19 |
| | 0.01 | 0.48 | 9.33 | 3.11 | 22.98 | 0.23 | 28.51 | 7.84 | 22.23 | 0.23 | 0.28 |

** Significant at 0.01 of probability level.

Table 18: The mutagenicity of ascorbic acid towards HRI-27 strain of *Leucaena leucocephala* rhizobia.

| Concentration $\mu\text{g ml}^{-1}$ | Preincubation | | | | | Plate-incorporation | | | | | Mutation rate $\times 10^{-7}$ |
|--|--|---------------|--|---------------------|--|--|---------------|--|---------------------|--|--------------------------------------|
| | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | Total No. of colonies $\text{ml}^{-1} \times 10^7$ | Survival % | Total No. of mutant $\text{ml}^{-1} \times 10^7$ | Mutagenicity (%) | Mutation frequency $\times 10^7$ | |
| 0.0 | 18.00 | 100.0 | 0.00 | 0.00 | 0.00 | 62.66 | 100.0 | 0.00 | 0.00 | 0.00 | 0.00 |
| 60 | 13.00 | 72.21 | 6.33 | 49.35 | 0.49 | 45.33 | 7.22 | 25.66 | 56.32 | 0.44 | 0.42 |
| 120 | 3.20 | 17.77 | 2.20 | 69.42 | 0.69 | 46.33 | 7.38 | 31.66 | 68.79 | 0.68 | 0.48 |
| 180 | 0.86 | 4.81 | 0.53 | 61.11 | 0.60 | 36.33 | 5.79 | 21.00 | 59.91 | 0.59 | 0.42 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** | ** |
| L.S.D. | 0.05 | 1.70 | 5.98 | 2.92 | 25.85 | 0.26 | 13.65 | 1.08 | 6.74 | 22.28 | 0.14 |
| | 0.01 | 2.58 | 9.06 | 4.42 | 39.15 | 0.39 | 20.67 | 1.64 | 10.21 | 33.74 | 0.22 |

** Significant at 0.01 of probability level.

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

Table 19: Identification of rhizobia biochemical mutants using nine plates of minimal medium each supplemented with different combinations of growth factors.

| Isolates | Plate No. | | | | | | | | |
|----------------|-----------|---|---|---|---|---|---|---|---|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| acr-1 FFAMU-8 | - | - | - | + | - | + | - | - | - |
| acr-2 FFAMU-8 | + | - | - | - | - | + | - | - | - |
| acr-3 FFAMU-8 | - | - | + | - | - | - | - | + | - |
| acr-4 FFAMU-8 | + | + | + | + | + | + | + | + | + |
| asc-1 FFAMU-8 | - | - | + | - | - | - | - | + | - |
| asc-2 FFAMU-8 | - | - | - | - | + | - | + | - | - |
| asc-3 FFAMU-8 | - | + | - | - | - | + | - | - | - |
| asc-4 FFAMU-8 | - | - | + | - | - | - | + | - | - |
| acr-1 ARCG -10 | + | - | - | - | - | - | - | + | - |
| acr-2 ARCG -10 | - | - | - | - | + | + | - | - | - |
| acr-3 ARCG -10 | - | - | - | - | - | - | - | - | - |
| acr-4 ARCG -10 | - | + | - | - | - | + | - | - | - |
| asc-1 ARCG -10 | - | - | - | - | + | - | + | - | - |
| asc-2 ARCG -10 | + | - | - | - | - | - | + | - | - |
| asc-3 ARCG -10 | - | - | - | + | - | - | - | - | + |
| asc-4 ARCG -10 | - | - | - | + | - | + | - | - | - |
| acr-1 - NRC-19 | - | - | + | - | - | - | - | + | - |
| acr-2 - NRC-19 | - | - | - | - | + | - | - | + | - |
| acr-3 - NRC-19 | - | + | - | - | - | + | - | - | - |
| acr-4 - NRC-19 | + | - | - | - | - | - | + | - | - |
| asc-1 - NRC-19 | - | - | + | - | - | - | - | + | - |
| asc-2 - NRC-19 | + | - | - | - | - | - | - | - | - |
| asc-3 - NRC-19 | + | - | - | - | - | - | - | - | + |
| asc-4 - NRC-19 | - | - | - | - | - | - | - | - | - |
| acr-1 - HRI-27 | - | - | - | - | + | - | - | + | + |
| acr-2 - HRI-27 | + | + | + | + | + | + | + | + | - |
| acr-3 - HRI-27 | - | - | - | + | - | - | + | - | - |
| acr-4 - HRI-27 | - | - | - | - | + | - | - | + | - |
| asc-1 - HRI-27 | - | - | - | - | - | - | - | - | - |
| asc-2 - HRI-27 | - | - | + | - | - | + | - | - | - |
| asc-3 - HRI-27 | - | - | - | - | - | - | - | - | - |
| asc-4 - HRI-27 | + | - | - | - | - | - | + | - | - |

++ = Positive growth.

- = Negative growth.

acr and asc = Auxotrophic mutant derived from acridine and ascorbic acid treatments, respectively.

Table 20: Genotype identification of *Rhizobium (Sesbania sesban)* auxotrophic mutants.

| Mutants | Source and/or reference | Mutagen | Genotype |
|------------------|-------------------------|-----------------|---------------------------|
| acr-1 - FFAMU-8 | FFAMU-8 | Acridine orange | <i>sys</i> ⁻ |
| acr-2 - FFAMU-8 | FFAMU-8 | Acridine orange | <i>ala</i> ⁻ |
| acr-3 - FFAMU-8 | FFAMU-8 | Acridine orange | <i>trp</i> ⁻ |
| acr-4 - FFAMU-8 | FFAMU-8 | Acridine orange | Revertant to wild type |
| asc-1 - FFAMU-8 | FFAMU-8 | Ascorbic acid | <i>trp</i> ⁻ |
| asc-2 - FFAMU-8 | FFAMU-8 | Ascorbic acid | <i>met</i> ⁻ |
| asc-3 - FFAMU-8 | FFAMU-8 | Ascorbic acid | <i>arg</i> ⁻ |
| asc-4 - FFAMU-8 | FFAMU-8 | Ascorbic acid | <i>leu</i> ⁻ |
| acr-1 - ARCG -10 | ARCG-10 | Acridine orange | <i>phe</i> ⁻ |
| acr-2 - ARCG -10 | ARCG-10 | Acridine orange | <i>gly</i> ⁻ |
| acr-3 - ARCG -10 | ARCG-10 | Acridine orange | Other requirements |
| acr-4 - ARCG -10 | ARCG-10 | Acridine orange | <i>arg</i> ⁻ |
| asc-1 - ARCG -10 | ARCG-10 | Ascorbic acid | <i>met</i> ⁻ |
| asc-2 - ARCG -10 | ARCG-10 | Ascorbic acid | <i>glu</i> ⁻ |
| asc-3 - ARCG -10 | ARCG-10 | Ascorbic acid | <i>pyrid</i> ⁻ |
| asc-4 - ARCG -10 | ARCG-10 | Ascorbic acid | <i>sys</i> ⁻ |

acr = Acridine orange

asc = Ascorbic acid.

Table 21: Genotype identification of *Rhizobium (Leucaena leucocephala)* auxotrophic mutants.

| Mutants | Source and/or reference | Mutagen | Genotype |
|----------------|-------------------------|-----------------|--|
| acr-1 - NRC-19 | NRC-19 | Acridine orange | <i>trp</i> ⁻ |
| acr-2 - NRC-19 | NRC-19 | Acridine orange | <i>val</i> ⁻ |
| acr-3 - NRC-19 | NRC-19 | Acridine orange | <i>arg</i> ⁻ |
| acr-4 - NRC-19 | NRC-19 | Acridine orange | <i>glu</i> ⁻ |
| asc-1 - NRC-19 | NRC-19 | Ascorbic acid | <i>trp</i> ⁻ |
| asc-2 - NRC-19 | NRC-19 | Ascorbic acid | <i>ade</i> ⁻ |
| asc-3 - NRC-19 | NRC-19 | Ascorbic acid | Revertant to wild type |
| asc-4 - NRC-19 | NRC-19 | Ascorbic acid | <i>val</i> ⁻ |
| acr-1 - HRI-27 | HRI-27 | Acridine orange | Revertant to wild type <i>lys</i> ⁻ |
| acr-2 - HRI-27 | HRI-27 | Acridine orange | <i>val</i> ⁻ |
| acr-3 - HRI-27 | HRI-27 | Acridine orange | Other requirements |
| acr-4 - HRI-27 | HRI-27 | Acridine orange | <i>asp</i> ⁻ |
| asc-1 - HRI-27 | HRI-27 | Ascorbic acid | <i>glu</i> ⁻ |
| asc-2 - HRI-27 | HRI-27 | Ascorbic acid | Other requirements |
| asc-3 - HRI-27 | HRI-27 | Ascorbic acid | <i>glu</i> ⁻ |
| asc-4 - HRI-27 | HRI-27 | Ascorbic acid | |

acr = Acridine orange,

asc = Ascorbic acid.

The results are in accord with those of Batzli *et al.* (1992), who found, so much rhizobial diversity was found between the two sites in Green Ridge State Forest (GRSF) in Allegany country, Maryland and even within individual trees at one site. Differences in rhizospheric microsite condition may be a significant source of variation among strains collected from the two sites. The results indicated that selection of rhizobia strains on the basis of the efficiency of the symbiotic association with leguminous trees is possible.

The mean generation time (h) of rhizobial isolates nodulating *Sesbania sesban* and *Leucaena leucocephala* rhizobia ranged from 4.813 to 6.437 and 4.600 to 6.523, respectively. These are in accordance with the results of Batzli *et al.* (1992), who found that the mean generation times of rhizobia (*Robinia pseudoacacia* L.) isolates used in their study ranged from 3 to 9 h. In addition, the results obtained here are in agreement with the results of the same authors, who found that 186 isolates of rhizobia from nodules of *Robinia pseudoacacia* L. varied significantly in their resistance to antibiotics and NaCl, their growth on different carbohydrates, and their effect on the pH of culture media. Most isolates, of the same authors, showed intermediate antibiotic resistance, the capacity to use numerous carbohydrates, and a neutral to acid pH response. The same authors also reported that isolates had greater similarity within sampling locations than among sampling locations.

Effect of salinity on mean generation time (MGTs) of *Rhizobium* sp. related to *Sesbania sesban* and *Leucaena leucocephala*:

The addition of NaCl to YEM broth increased the MGT for all isolates tested (Tables 5 and 6). Most of *Rhizobium* strains failed to grow in the highest levels of salt. The salt tolerance of each symbiont of the legume-*Rhizobium* symbiosis may differ (Parker *et al.*, 1977). Despite a large range in tolerance to salts among species of legumes, there are no agriculture legumes that can be considered to be highly salt tolerant. *Rhizobium* isolates of *Leucaena leucocephala* showed visible growth on 4% NaCl also appeared an increase in MGT. Comparing the sensitivity of the micro symbiont, *Rhizobium*, with that of the legume host will then indicate the relative value of efforts to increase rhizobial tolerance to salts in strain selection programs. The growth of all but some of the rhizobia (*Leucaena*) tested was slowed by the presence of NaCl (Table 6). These results are contrary to those obtained by Pillai and Sen (1973), who showed that the growth rate of *Rhizobium* spp. increased with 1% NaCl added to broth media (EC = 18.0mS cm⁻¹). The results obtained by Steinborne and Roughley (1975), on the other hand, are in agreement with the results obtained here, who shown that the growth of both *R. trifolii* and *R. meliloti* is slowed by the addition of salt. Large increases in MGT, or growth suppression, was not observed for any of the isolates of *Sesbania sesban* until the concentration of NaCl reached 4%. On the other hand, the growth suppression was observed for all isolates of *Leucaena leucocephala* at the concentration of 5% NaCl. The present results are in agreement with those of Hashem *et al.* (1998), who found that the addition of high concentrations of NaCl to YEMB increased the MGT of all *Rhizobium* (*Leucaena*) strains, and some strains failed to grow on 1.5 and 3% NaCl, however some isolates from highly saline soils (Swelim *et al.*, 1997) were consistently more tolerant to salt stress than other strain since it alone grew at > 3.0% NaCl.

The addition of high concentrations of NaCl to YEMB increased the MGT of all isolates from two- to five-fold, depending on the strain. This study identifies and characterizes some agronomically important strains of *Sesbania* and *Leucaena* nodulating bacteria from Egypt that are quite salt tolerant and resistant to 3.0% NaCl. The growth rates of these tolerant strains were slowed two- to five-fold by the elevation of salinity from 0 to 3% NaCl. The results obtained here are in accord with those of Saxena *et al.* (1996), who found that salinity-induced changes in the protein profiles in *Rhizobium* sp. exhibited alterations in the expression of as many as 19 proteins, which either showed an enhanced rate of synthesis or a decline in the levels as compared with controls. All these proteins were predominantly of low molecular mass (below 40 K Da) except for one (52 K Da), the difference in the protein profiles was more marked in salt-grown cells as compared with

salt-shocked cells.

Carbon source utilization: The results presented in Tables 7 and 8 indicate the utilization of monosaccharides (D-glucose, D-fructose and D-galactose), C₃ compounds (glycerol), sugar alcohol (D-mannitol), disaccharides (lactose, maltose and sucrose), pentoses (L-arabinose, D-arabinose and xylose) and polysaccharides (dextrin, dextrose and diastase), by different isolates of rhizobia-leguminous trees. The rhizobia isolate of *Sesbania sesban* named FAFMU₄ did not grow on the mineral medium with inositol, however, the isolate of the same host named FPMU₁ did not grow on the mineral medium with L-arabinose, D-arabinose and glycerol as a sole carbon source. In contrast, all isolates of *Sesbania sesban* rhizobia grew normally on the different carbon sources tested. On the other hand, the isolate of *Leucaena leucocephala* rhizobia named FFAMU₂₁ did not grow on the mineral medium with xylose, glycerol and inositol. However, the isolates named FFAMU₂₂ and FFAMU₂₃ did not grow on the mineral medium (MM) with xylose and glycerol, respectively. In addition, the isolate named MAMG₂₆ did not grow on the MM with glycerol and inositol. In contrast, all isolates grew normally on different carbon sources tested. The biochemical characteristics such as carbon sources utilization were similar to those described for fast-growing rhizobia (Arias *et al.*, 1979 and Jordan, 1984). The present results are in agreement with those reported by Milnitsky *et al.* (1997), who described some biochemical and genetic characteristics of seven rhizobial isolates that nodulate legume trees, native to Uruguay, they found that carbon utilization by the isolates could be assigned to the fast-growing group of rhizobia. The study of a greater number of isolates from our native legume trees and the use of other techniques will allow us to have a more complete picture about their taxonomy and evolutionary relationship.

Rhizobia from woody species have been described as fast, moderate and slow growing (Barnet and Catt, 1991), and many have yet to be assigned to a genus. Sutherland and Sprent (1993) concluded that rhizobia isolated from tree species are highly variable and that it is difficult to assign them to standard genera and species in terms of carbohydrate metabolism, pH change of the growth medium, optimal temperature of growth flagellation and growth rate. The results obtained in this study indicated that the fast-growing isolates of leguminous trees showed excellent growth within 3 days on most of the substrates tested and therefore resembled the fast-growing rhizobia and not the slow-growing species of *Rhizobium*. The slow-growing rhizobia were shown by Graham and Parker (1964) to be more fastidious.

Prevalence of intrinsic antibiotic resistance: The minimum inhibitory concentration (MIC) of nine antibiotics for wood legume *Rhizobium*, isolated from different locations are shown in Tables 9 and 10. The intrinsic antibiotic resistance (IAR) of wood legume rhizobia was extremely high against tetracycline (25 µg ml⁻¹) in the isolates ARCG₁₂, MAMG₂₅, and MAMG₂₆. The lowest minimum inhibitory concentration was occurred with tetracycline (≤ 25 µg ml⁻¹). Extremely high IAR against ampicillin (150 µg ml⁻¹) was shown in isolates, NRC₁₀, FFAMU₂₄, MAMG₂₅, and FFSMU₁. These results were comparable with those obtained from temperate fast-growing rhizobia reported by Kremer and Peterson (1982) and Stewers and Eaglesman (1984). Chan *et al.* (1988) reported that intrinsic antibiotic resistance (IAR) method is useful for identifying many *Rhizobium* spp., but its effectiveness for evaluating isolates from *A. sinicus* was inconsistent with certain antibiotic concentration combinations, similar to the results of Stein *et al.* (1982), who examined *Rhizobium phaseoli*. This variation of response may be attributed to spontaneous mutations of the organisms (Kingsley and Bohlool, 1983), or to differences in cell load upon inoculation. However, many antibiotic concentration did not provide definite patterns of resistance. Some isolates of rhizobia *Sesbania sesban*, *Leucaena leucocephala*, L were resistant to high concentration of neomycin (≤ 150 µg ml⁻¹), kanamycin (≤ 150 µg ml⁻¹), streptomycin (≤ 200 µg ml⁻¹), spectomycin (≤ 200 µg ml⁻¹), rifampicin (≤ 175 µg ml⁻¹), penicillin (≤ 400 µg ml⁻¹) and chloramphenicol (≤ 250 µg ml⁻¹). Soybean and non-soybean rhizobia resistant to high concentration of penicillin and other antibiotics have been reported by Chao (1987), who suggested

that the high antibiotic resistance among rhizobia isolated from subtropical-tropical Taiwan soils could be caused by the use of pesticides in the field.

In the present study, one isolate of *Sesbania sesban* rhizobia was extremely high against streptomycin ($200 \mu\text{g mL}^{-1}$). The resistance to streptomycin has proved to be a stable marker with high level of resistance available and little associated loss of symbiotic effectiveness (Levin and Montgomery, 1974). Two isolates of *Leucaena leucocophala* rhizobia were resistant to high concentrations of spectomycin ($\leq 200 \mu\text{g mL}^{-1}$). Spectomycin resistance in *Rhizobium* is also similar to that of streptomycin and provides a second potential marker for use in ecological studies (Schwinghamer and Dudman, 1973). Chan *et al.* (1988) found that both plasmid visualization and IAR showed similarities with the isolates of *Bradyrhizobium sp.* (*Astragalus sinicus* L.). Because antibiotic resistance of an organism is often coded by a plasmid, the smaller plasmid (80 Md) seen in some isolates may carry the gene that codes for tetracycline or neomycin resistance. These were the only isolates which both contained this plasmid and showed resistance to tetracycline and neomycin. MAMG₂₅₇, ampicillin resistant culture ($\leq 150 \mu\text{g mL}^{-1}$) was resistant to the same concentration of both neomycin and kanamycin. HRI₂₇, also showed the same trend for both ampicillin ($\leq 150 \mu\text{g mL}^{-1}$). Cultures of FAFMU₅ and FAFMU₆ resistant to neomycin ($\leq 150 \mu\text{g mL}^{-1}$) were resistant to penicillin ($\leq 400 \mu\text{g mL}^{-1}$). FAFMU₆ also exhibited a similar nature for both penicillin ($\leq 400 \mu\text{g mL}^{-1}$) and chloramphenicol ($\leq 300 \mu\text{g mL}^{-1}$). However, ARCG₁₀ streptomycin resistant culture ($\leq 200 \mu\text{g mL}^{-1}$) was resistant to penicillin ($\leq 400 \mu\text{g mL}^{-1}$). Date and Hurse (1992) reported that the reduction in symbiotic effectiveness of 15 of the 59 resistant strains of *Bradyrhizobium* for desmodium intortum is in agreement with Bromfield and Jones (1979) observation that antibiotic resistance can be associated with decreased effectiveness. However, their results offer little support for a general association between loss of effectiveness and resistance to particular groups of antibiotics, including streptomycin and rifampicin (Pankhurst, 1977).

Mutation frequencies of acridine orange in *Rhizobium* of legume trees: The results summarized in Tables 11, 12, 13 and 14 demonstrate a dose-response for decreasing cell survival at levels that are not excessively toxic to the bacteria at both pre-incubation and plate incorporation assays. The standard assay without pre-incubation involves adding, in order, the test compound, the bacterial tester strain, to soft agar, which is then briefly mixed and poured directly onto the BM₂ agar plate. This assay requires somewhat less time and fewer manipulations than the modification with pre-incubation, but is somewhat less sensitive for some mutagens. The standard assay with pre-incubation was quite toxic to the bacteria than plate incorporation test. This is in accordance with Adams (1959), who reported that acridines inhibit bacteriophage development at concentrations below those inhibiting the bacterial host.

Mutagenicity data for acridine orange appeared here revealed that in all cases the addition of plate incorporation resulted in equal or greater sensitivity than the standard assay of pre-incubation. A reproducible dose-response with an increase in the number of auxotrophic mutants as shown here over the spontaneous value is an evidence for mutagenicity.

Acridine hypersensitivity has also been found in T₄ phage with mutations in a variety of genes, many of which are DNA synthesis and repair genes, implying that some inhibitory effects of acridines act at this level (Woodworth and Kreuzer, 1996). An alternative view, by Brenner *et al.* (1961), is that acridine mutations and the majority of spontaneous mutations are not caused by base-pair substitutions at all, but by the deletion or addition of a base-pair. Obviously such mutations could not be reverted by any agent whose effect was simply to replace one base-pair by another, but only by the restoration of a deleted base or the removal of an extra one.

Mutagenic activity of ascorbic acid: The data presented in Tables 15, 16, 17 and 18 show that ascorbic acid had genotoxic hazards for *Rhizobium*. It demonstrates a dose-response for both cell survival and mutagenicity at levels that are not excessively toxic

to the bacteria. The standard assay without pre-incubation (plate incorporation) involves adding, in order, the test compound and the bacterial tester strain to agar plates. The mutagenic activity of ascorbic acid in all isolates tested resulted in equal or greater sensitivity in the addition of pre-incubation than without it (plate incorporation) at most concentrations used. The standard assay without pre-incubation is preferable for generating a dose-response curve and for general testing (Levin *et al.*, 1984). Recent reports have indicated that ascorbic acid, at high concentrations, may be mutagenic (Galloway and Painter, 1979) and cytotoxic (Stich *et al.*, 1979) in the *in vitro* mutagenesis bioassays. This effect was initially attributed to the generation of hydrogen peroxide (H₂O₂) and/or hydroxyl free radicals (HO[•]), but this does not seem to be clear.

These results reported here also show a dose-response for mutagenicity ratio in all rhizobial isolates tested. These are in agreement that oxidation products of ascorbic acid are mutagenic for microbial and mammalian cells; ascorbic acid in the presence of oxygen converts covalently-closed circular DNA to open circular DNA molecules, that causes single strand breaks in DNA (Bode, 1967).

The results obtained in this study are also in harmony with those of Amabile-Cuevas *et al.* (1991), who reported that treatment of *Staphylococcus aureus* with 1mM ascorbic acid for 6 h induced an increased loss of resistance markers in 4 of 6 strains tested, and agarose gel electrophoresis showed the disappearance of plasmid DNA in ascorbate-induced susceptible colonies. In addition, the presence of ascorbate induced a 50-75% decrease in minimal inhibitory concentrations of different antibiotics for resistant strains of *Staphylococcus aureus*.

Identification of biochemical mutants: As shown from the results presented in Table 19, nine plates of minimal medium are each supplemented with different combinations of growth-factors with the arrangement shown in the related Table 2. The number of requirements tested can be varied by altering the number of plates and the arrangement of growth-factors among them. Most of biochemical mutants as shown here in with a single biochemical requirement, because when inoculated on nine plates will grow on two of them, one of plates 1-5 and one of plates 6-9. Although, some of biochemical mutants such as acr-4 FAFMU-8 and acr-2-HRI-27 were reverted to wild type, because of their ability to grow on all of minimal medium plates. In contrast, some others such as acr-3 ARCG-10, asc-4-NRC-19, asc-1-HRI-27 and asc-3-HRI-27 failed to grow on any of minimal medium plates, indicating that these mutants have another requirements than that are listed in the nine plates recorded in this study. Mutants requiring for example, cysteine for growth, will only grow on plates 4 and 6. Mutants with alternative requirements will grow on more than two plates. When large number of mutants are to be identified, this technique is more efficient than previous methods in which the field of each is progressively reduced. With one or a few mutants auxanographic techniques are preferable. The present method for the identification of biochemical mutants of *Rhizobium* has been carried out according to Holliday (1956).

Leading to the results presented in Tables 19, 20 and 21 summarizes the identification of biochemical mutants induced. In this respect, once a mutant has been isolated, it is frequently desirable to characterize the nature of the genetic defect by further biochemical or genetic analyses. For example, a nutritional requirement can result from mutations in several genes. Thus, an auxotrophic mutant unable to synthesize a specific amino acid, for example, may be blocked in any one of the several sequential enzymatic reactions in that synthetic pathway.

The results obtained in this study are in agreement with those reported by Kerppola and Kahn (1988), who found that mutants in ornithine transcarbamylase, argininosuccinate synthetase or serine-glycine biosynthesis formed nitrogen-fixing (Fix⁺) nodules on the roots of alfalfa (*Medicago sativa*), as well as prototrophic revertants were always fix⁺. Malek (1974) found that most of ineffective auxotrophic mutants of *Rhizobium meliloti* required histidine, histidine + cysteine, cysteine, arginine or casamino acid + yeast extract showed resistance to five tested phages. The finding of Malek (1974) support observation of Kleczkowska

(1965) that mutation to phage resistance in *Rhizobium* frequently causes their ineffectiveness. On the other hand, Atkins and Hayes (1972) tested four phage resistant mutants of *R. trifolii* and concluded that these mutations were concerned with modification of constitution or organization of the polymers of the cell wall which caused degradation of lipopolysaccharide, known as phage receptors in gram-negative bacteria. Abdel-Wahab (1977) reported that single auxotrophic mutants induced from the effective *Rhizobium trifolii* WTG, requiring *His*, *Lys*, *Leu*, *Ade* or *Ura* were ineffective. However, these requiring *Arg*, *Meth* or *Trp* more effective in nitrogen fixation as the original WTG. Also, double auxotrophic mutants, *Arg*, *His*, *Leu*, *Meth*, *Lys*, *Tryp*, *Leu Ade* and *Lys Ura* were ineffective in nitrogen fixation. In conclusion the results show that, using the "treated-and-plate" test in mutant experiments, one has to control the physiological state of the cells very carefully. This is especially important when the bacteria are treated in non-growth medium with simple acridine and ascorbic acid which do not damage DNA, but acridine induce frame shift mutations only when present during replication. On the other hand, ascorbic acid is mutagenic and cytotoxic probably through the generation of hydrogen peroxide if added to bacteria in a medium prepared with sterile tap water.

References

- Abdel-Wahab, S.M., 1977. Genetic control of nitrogen fixation in *Rhizobium trifolii* strains. Ph.D. Thesis, Fac. Agric., Cairo Univ., Egypt.
- Adams, M.H., 1959. Bacteriophages. Interscience Publishers Inc. New York.
- Allen, O.O., 1959. Experiments in Soil Bacteriology. 3rd edition, Burges Publishing Co., Minneapolis, USA.
- Amabile-Cuevas, C.F., R.M. Pina-Zentella and M.E. Wah-Laborde, 1991. Decreased resistance to antibiotics and plasmid loss in plasmid-carrying strains of *Staphylococcus aureus* treated with ascorbic acid. *Mutation Res.*, 264: 119-125.
- Arias, A., C. Corvenansky, A. Gardiol, G. Martinez-Prets, 1979. Phosphoglucose isomerase mutant of *Rhizobium meliloti*. *J. Bacteriol.*, 137: 409-414.
- Atkins, G.J. and A.H. Hayes, 1972. Surface changes in a strain of *Rhizobium trifolii* on mutation to bacteriophage resistance. *J. Gen. Microbiol.*, 73: 273
- Balasa, G., 1963. Genetic transformation in *Rhizobium*. A review of the work of R. Blassa. *Bacteriol. Rev.*, 27: 228-241.
- Barnet, Y.M. and P.C. Catt, 1991. Distribution and characterization of root-nodule bacteria isolated from Australian *Acacia* sp. *Plant and Soil*, 135: 109-120.
- Batzli, J.M., W.R. Graves and P. Berkum, 1992. Diversity among rhizobia effective with *Robinia pseudoacacia* L. *Appl. Env. Microbiol.*, 58: 2137-2143.
- Beale, G.H., 1948. A method for measurement of mutation rate from phage sensitivity to phage resistance in *Escherichia coli*. *J. Gen. Microbiol.*, 2: 131-197.
- Beringer, J.E., 1974. R factor transfer in *Rhizobium leguminosarum*. *J. Gen. Microbiol.*, 84: 188-198.
- Bode, V.C., 1967. *J. Mol. Biol.*, 26: 125-129. (Cited from Stich, H.F.; J. Karim; J. Koropatnick and L. Lo. 1976. *Nature*, 260: 722-724.
- Brenner, S., L. Barnett, F.H. Crick and A. Orgel, 1961. The theory of mutagenesis. *J. Mol. Biol.*, Vol., 121.
- Bromfield, E.S.P. and P.G. Jones, 1979. The competitive ability and symbiotic effectiveness of doubly labeled antibiotic resistant mutants of *Rhizobium trifolii*. *J. Ann. Appl. Biol.*, 91: 211-219.
- Chan, C.L., T.A. Lumpkin and C.S. Root, 1988. Characterization of *Bardyrhizobium* sp. (*Astragalus sinicus* L.) using serological agglutination, intrinsic resistance, plasmid visualization, and field performance. *Plant and Soil*, 109: 85-91.
- Chao, C.C., 1987. Antibiotic and pesticide resistant characteristics of indigenous rhizobia in Taiwan. *J. Chin. Agric. Chem. Soc.*, 25: 308-317.
- Cole, M.A. and G.H. Elkan, 1973. Transmissible resistance to penicillin G, neomycin and chloramphenicol in *Rhizobium japonicum*. *Antimicrob. Agent Chemother.*, 4: 248-253.
- Cole, M.A. and G.H. Elkan, 1979. Multiple antibiotic resistance in *Rhizobium japonicum*. *Appl. Env. Microbiol.*, 37: 867-870.
- Dowling, D.N. and W.J. Broughton, 1986. Competition for nodulation of legumes. *Ann. Rev. Microbiol.*, 40: 131-157.
- Date, R.A. and L.S. Hurs, 1992. Growth, competitiveness and effectiveness of spontaneous antibiotic resistant strains of *Bradyrhizobium* for *Desmodium intortum* cv. green leaf. *Soil Biol. Biochem.*, 24: 33-39
- Eckardt, F. and R.H. Haynes, 1977. Induction of pure and sectoral mutant colonies in excision. proficient and deficient strains of yeast. *Mutation Res.*, 43: 327-338.
- Fitzsimon, A., F. Duffner, D. Curtin, G. Brophy, P. O'Kiely and M. O'Connell, 1992. Assessment of *Pedococcus acidilactici* as potential silage inoculant. *J. Appl. Microbiol.*, 58: 3047-3052.
- Galloway, S.M. and R.B. Painter, 1979. Vitamin C is positive in the DNA-synthesis inhibition and sister-chromatid exchange tests. *Mutation Res.*, 60: 321-327.
- Graham, P.H., 1963. Antibiotic sensitivities of the root-nodule bacteria. *Australian J. Biol. Sci.*, 16: 557-559.
- Graham, P.H. and C.A. Parker, 1964. Diagnostic features in the characterization of root-nodule bacteria of legumes. *Plant and Soil*, 20: 383-396.
- Hashem, F.M., D.M. Swelim, L.D. Kuykendall, A.I. Mohamed, S.M. Abdel-Wahab and M.N. Hegazi, 1998. Identification and characterization of salt- and thermo-tolerant *Leucaena* nodulating *Rhizobium* strains. *Bio. Fertil. Soils*, 27: 474.
- Holliday, R., 1956. A new method for the identification of biochemical mutants of micro-organisms. *Nature*, 177: 987.
- Jordan, D.C., 1984. Rhizobiaceae. In *Bergey's Manual Systematic Bacteriology*. Vol. 1. Eds. N.R. Krieg and J.G. Holt. pp: 235-256.
- Kerppola, T.M. and M.L. Kahn, 1988. Symbiotic phenotypes of auxotrophic mutants of *Rhizobium meliloti* 104A14. *J. General Microbiol.*, 134: 913-919.
- Kingsley, M.T. and B.B. Bohlool, 1983. Characterization of *Rhizobium* sp. (*Cicer arietinum* L.) by immunofluorescence, immunodiffusion, and intrinsic antibiotic resistance. *Can. J. Microbiol.*, 29: 518-526.
- Kleczkowska, J., 1965. Mutation in symbiotic effectiveness in *Rhizobium trifolii* caused by transforming DNA and other agents. *J. Gen. Microbiol.*, 40: 377.
- Kremer, R.J. and H.L. Peterson, 1982. Nodulation efficiency of legume inoculation as determined by intrinsic antibiotic resistance. *Appl. Env. Microbiol.*, 43: 636-642.
- Lafontaine, P.J., C. Lafreniere and H. Antoun, 1989. Some properties of carbohydrate and C₄-dicarboxylic acid utilization negative mutants of *Rhizobium leguminosarum* biovar *phaseoli* strain P 121. *Plant and Soil*, 120: 195-201.
- Lafreniere, C., P. Lafontaine, C. Marion and H. Antaun, 1987. Oxidation of substrates in organic acids utilization negative mutants and the wild type *Rhizobium meliloti* strain S⁺. *Plant and Soil*, 101: 73-78
- Lenhard, G., 1956. Die Dehydrogenaseaktivität des Bodens als Maß für die Mikroorganismen-tätigkeit im Boden. *Z. Pflanzenernähr. Düng. Bodenkd.*, 73: 1-11.
- Levin, D.E., M. Hollstein, M.F. Christman and B.N. Ames, 1984. Detection of oxidative mutagens with a new *Salmonella* tester strain (TA 102). *Methods in Enzymology*, 105: 249-254.
- Levin, R.A. and M.P. Montgomery, 1974. Symbiotic effectiveness of antibiotic-resistant mutants of *Rhizobium japonicum*. *Plant and Soil*, 41: 669-676.
- Malek, W., 1974. Relation between phage sensitivity and symbiotic properties of auxotrophic mutant of *Rhizobium meliloti*. *Acta Microbiol. Polon. Ser. A*, 6: 225-229.
- McCarty, M., 1945. Reversible inactivation substance inducing transformation of pneumococcal types. *J. Exp. Med.*, 81: 507-514.

Nassef *et al.*: Auxotrophic mutants induced in rhizobia

- Milnitsky, F., L. Frioni and F. Agius, 1997. Characterization of Rhizobia that nodulate native legume trees from Uruguay. *Soil Biol. Biochem.*, 29: 989-992.
- Mohn, G.R. and J. Ellenberger, 1977. The use of *E. coli* K 12/343/113 (λ) as a multipurpose indicator strain in various mutagenicity testing procedures, in: B. Kilbey, M.S. Legator, W. Nichols and C. Ramel (Eds.), *Handbook of Mutagenicity Test Procedures*, Elsevier / North Holland, Amsterdam, pp: 95-118.
- Murata, A., H. Suenaga, S. Hideshima, Y. Tanaka and F. Kato, 1986. Hydroxyl radical as the reactive species in the inactivation of phages by ascorbic acid. *Agric. Biol. Chem.*, 50: 1481-1487.
- Murphy, S.C. and G.H. Elkan, 1963. Growth inhibition by biotin in strain of *R. japonicum*. *J. Bacteriol.*, 86: 848-885.
- Norris, D.O., 1965. Acid production by *Rhizobium*, a unifying concept. *Plant and Soil*, 22:143-166.
- Pain, A.N., 1979. Symbiotic properties of antibiotic resistant and auxotrophic mutants of *Rhizobium leguminosarum*. *J. Appl. Bacteriol.*, 47: 53-64.
- Pandey, K.D. and A.K. Kashyap, 1992. Induction of mutation in the cyanobacterium *Anabaena doliolum*: A strain-specific property. *J. Folia Microbiol.*, 37: 377-380
- Pankhurst, C.E., 1977. Symbiotic effectiveness of antibiotic resistant mutants of fast and slow growing strains of *Rhizobium* nodulating lotus species. *Can. J. Microbiol.*, 23: 1026-1033.
- Parker, C.A., M.R. Trinick and D.L. Chatel, 1977. Rhizobia as soil and rhizosphere inhabitants. P. 311-352. In: R.W.F. Hardy and A.H. Gibson (ed.), *Atreatise on dinitrogen fixation*, Vol. 4. John Wiley & Sons, Inc., New York.
- Pillai, R.N. and A. Sen, 1973. Salt tolerance of *Rhizobium* from *Polichos lablab*. *Zentralbl. Bakteriell. Abt. Parasitenkd, Infektionsk. Hyg.*, 2: 538-542.
- Pons, F.W. and P. Müller, 1989. On the glucose effect in acridine-induced frame shift mutagenesis in *E. coli*. *Mutation Res.*, 210: 71-77.
- Postgate, J.R., 1982. *The fundamentals of Nitrogen Fixation*. Cambridge University Press, Cambridge.
- Saxena, D., M. Amin and S. Khanna, 1996. Modulation of protein profiles in *Rhizobium sp.* under salt stress. *Can. J. Microbiol.*, 42: 617-620.
- Schlegel, H.G., 1985. *General Microbiology*. 6th ed. Cambridge University Press, Cambridge. (Cited from Luchansky *et al.*, 1992. *Appl. Environ. Microbiol.*, 58: 3053-3059).
- Schwinghamer, E.A. and W.F. Dudman, 1973. Evaluation of spectomycin resistance as a marker for ecological studies with *Rhizobium* spp. *J. Appl. Bacteriol.*, 36: 263-272.
- Steel, R.G.D. and J.H. Torrie, 1980. *Principles and Procedures of Statistics: A Biometric Approach*. McGraw-Hill, New York, NY, USA. pp: 633.
- Stein, M., E.S.P. Bromfield and M. Dye, 1982. An assessment of a method based on intrinsic antibiotic resistance for identifying *Rhizobium* strains. *Ann. Appl. Biol.*, 101: 261-267.
- Steinborne, J. and R.J. Roughley, 1975. Toxicity of sodium chloride ions to *Rhizobium spp.* in broth and peat culture. *J. Appl. Bacteriol.*, 34: 133-138.
- Stewers, M.D. and A.R.J. Eaglesman, 1984. Physiological and symbiotic characteristic of fast-growing *Rhizobium japonicum*. *Plant and Soil.*, 77: 3-14.
- Stich, H.F., L. Wei and R.F. Whiting, 1979. Enhancement of the chromosome-damaging action of ascorbic by transition metal. *Cancer Res.*, 39: 4145-4152.
- Sutherland, J.M. and J.I. Sprent, 1993. Nitrogen fixation by legumes trees. In *Symbiosis in Nitrogen-Fixing Trees*. (N.S. Subba Rao and C. Rodriguez-Barrueco, Eds.), pp: 33-63. Oxford and IBH Publishing Co. New Delhi.
- Swelim, D.M., F.M. Hashem, L.D. Kuykendall, N.I. Hegazi and S.M. Abdel-Wahab, 1997. Host specificity and phenotypic diversity of *Rhizobium* strains nodulating *Leucaena*, *Acacia* and *Sesbani* in Egypt. *Biol. Fertil. Soils*, 25: 224-232.
- Thorne, S.H. and H.D. Williams, 1997. Adaptation to nutrient starvation in *Rhizobium leguminosarum* bv. *phaseoli*: analysis of survival, stress resistance, and change in macro-molecular synthesis during entry to and exit from stationary phase. *J. Bacteriol.*, 179: 6894-6901.
- Thomas, S.M. and D.G. MacPhee, 1987. Rec-A independent mutagenesis in *Escherichia coli* may be subjected to glucose repression. *Mutation Res.*, 180: 67-73.
- Trinick, M.J., 1982. Competition between rhizobial strains for nodulation. In *Nitrogen Fixation Legumes* ed. Vincent, J.M. pp: 229-238. Sydney: Academic Press.
- Turco, R.F., T.B. Moorman and D.F. Bezdicek, 1986. Effectiveness and competitiveness of spontaneous antibiotic resistant mutants of *Rhizobium leguminosarum* and *Rhizobium japonicum*. *Soil Biol. Biochem.*, 18: 259-262.
- Van Sluys, M.A., R. Alcantara-Gomes and C.F.M. Menck, 1986. *Escherichia coli* 10th A. Mutant is not hypersensitive to ascorbic acid copper treatment - an H₂O₂ generating reaction. *Mutation Res.*, 174: 265-269.
- Vincent, J.M., 1970. *A Manual for the Practical Study of the Root Nodule Bacteria*. 13 P 15. Blackwell Scientific Publications. Oxford.
- Woodworth, D.L. and K.N. Kreuzer, 1996. Bacteriophage T₄ mutants hypersensitive to an antitumor agent that induces top isomerase DNA cleavage complexes. *Genetics*, 143: 1081-1090.