

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Biological Control Potential of Two *Streptomyces* Isolates on *Rhizoctonia solani*, the Causal Agent of Damping-off of Sugar Beet

¹Sadeghi, A., ²A.R. Hessian, ³H. Askari, ⁴S. Aghighi and ⁴G.H. Shahidi Bonjar

¹Department of Genomics, Agricultural Biotechnology Research Institute of Iran, Mahdasht Road, Karaj, Iran

²Plant Diseases Lab., Plant Pests and Diseases Research Institute, Mahdasht Road, Karaj, Iran

³Department of Plant Physiology, Biochemistry and Proteomics, Agricultural Biotechnology Research Institute of Iran

⁴Departments of Plant Protection and Biotechnology, College of Agriculture, Shahid Bahonar University, 22 Bahman Blvd, Kerman, Iran

Abstract: Biological control of sugar beet damping-off of *Rhizoctonia solani* by two *Streptomyces* isolates (S2 and C) was evaluated in this study. The *in vitro* antagonism assays showed that active isolates had inhibitory effects on mycelium growth of the three *R. solani* AG-4 isolates (Rs1, Rs2 and Rs3). Soil treatment either with isolate S2 or C formulation inhibited the disease completely and increased seedling stand in infected and uninfected treatments significantly ($p < 0.05$). Compared to controls, all treatments containing bacteria had enhanced shoot and root dry weight and root density. Both bacterial isolates maintained normal growth in pH ranges of 5.6, 7.2 and 8.0 at 29°C. Isolate C grew in pH ranges 5.6, 7.2 and 8.0 at 23-37°C, isolate S2 grew at 18-37°C in pH 5.6-8.0 but did not grow in acidic medium at 37°C. When pH decreased to 5.0, growth of C decreased and S2 inhibited. To elucidate the mode of antagonism, chitinase activity and siderophore production were evaluated. Both isolates showed chitinase activity on medium containing colloidal chitin. Biosynthesis of siderophore was detectable in isolate C but not in S2. The results of this study showed that these isolates had antifungal activities by production of siderophore and chitinase.

Key words: Antagonism, biological control, damping-off, *Rhizoctonia solani*, *Streptomyces*, Sugar beet

INTRODUCTION

Damping-off is a common problem in almost all field and greenhouse crops (Georgakopoulos *et al.*, 2002). *Rhizoctonia solani* Kuhn is one of soilborne pathogenic fungi have been associated with damping off (Moussa, 2002). Protection of sugar beet against this pathogen is important to maximize the crop yield. The use of fungicide-treated seeds is the main commercially option for control of damping off. Development of biocontrol systems has provided an effective approach for reducing environmental pollution associated with fungicides and the risk of fungicide resistance development. Biological control of soil borne diseases by microorganisms especially *Streptomyces* species have already been published (Crawford *et al.*, 1993; EL-Abyad *et al.*, 1993; Asaka and Shoda, 1996; Jones and Samac, 1996; Chamberlain and Crawford, 1999; Whipps, 2001; Xiao *et al.*, 2002).

Several species of *Streptomyces* have inhibitory effects on the most common soil borne fungi like

Fusarium oxysporum (Smith *et al.*, 1990; Abd-Allah, 2001; Getha and Vikineswary, 2002), *Pythium ultimum* (Crawford *et al.*, 1993; Yuan and Crawford, 1995; Paulitz and Belanger, 2001), *Verticillium* spp. (EL-Abyad *et al.*, 1993; Aghighi *et al.*, 2004), *Rhizoctonia solani* (Asaka and Shoda, 1996; Sabaratnam and Traquair, 2002), *Sclerotinia homeocarpa*, *Gaeumannomyces graminis* and *Microdochium nivale* (Chamberlain and Crawford, 1999). According to recent publications, application of *Streptomyces*, (Sabaratnam and Traquair, 2002; Chung *et al.*, 2005) or their active ingredient from the culture filtrate (Shih *et al.*, 2003) are effective for control of *Rhizoctonia* damping off. Beside, some species are plant growth promoting agents that it may be due to the ability of these bacteria to produce hydroxamate-type siderophores and/or other plant growth-promoting metabolites (Nassar *et al.*, 2003).

The mode of *Streptomyces* action includes inhibition of the pathogen by antifungal antibiotics (Smith *et al.*, 1990; Gupte and Naik, 1999) competition for iron through production of siderophore; production of

degradative enzymes such as chitinase and glucanase (El-Tarabily *et al.*, 2000) and chitin binding proteins (Bormann *et al.*, 1999). Trejo-Estrada (1998) reported antagonistic effects of a *Streptomyces violaceusniger* isolate on plant pathogens. They showed that biocontrol activity of this bacterium was resulted from three antifungal compounds; also the isolate was able to produce chitinase and glucanase. Results of De Boer (1998) revealed that antibiotics were involved in antifungal activities of chitinolytic bacteria. El-Tarabily *et al.* (2000) determined that production of chitinase and glucanase was the main mechanisms related to biocontrol activity of *Streptomyces viridodiasticus*. Besides, other compounds such as phenylacetic and sodium phenylacetate from *Streptomyces humidus* showed antifungal activity on plant pathogens (Hwang *et al.*, 2001). Practical application of some beneficial biocontrol agents such as Pseudomonads has been limited due to decreased bacterial survival during commercial processing and packaging of formulation or after application in the field (Walker, 2004). In contrast, *Streptomyces* are able to produce spores to maintain their viability at non-specific conditions (Ensign, 1978). However, Yuan and Crawford (1995) showed that the density of the viable *Streptomyces* spores in formulation was only 10 fold less than origin over a 6.5-month period of storage at room temperature. Other significant parameter influencing the survival of bacteria in soil is temperature. Survival and antifungal activity (antibiotic and siderophore production) of Pseudomonads strains decrease at temperatures above 24°C (Schmidt *et al.*, 2004) but *Streptomyces* are mesophilic bacteria and grow at a wide range of temperature between 15 and 37°C (Locci, 1994). These beneficial characteristics of *Streptomyces* have attracted many investigators for isolating and improvement of these bacteria in biocontrol programs.

In our preliminary study, *in vitro* inhibitory effects of two isolates of *Streptomyces* (S2 and C) were assessed against some important pathogenic fungi including *Fusarium graminearum*, *F. oxysporum* f. sp. *melonis*, *Pyricularia grisea* and *Rhizoctonia solani* which cause rice root rot, wheat head blight, melon damping-off, rice blast and rice sheath blight, respectively. Our main objectives in the present study were introduction of potentially active biocontrol *Streptomyces* isolates against *R. solani*, the causal agent of damping-off in sugar beets and partial characterization of their effectiveness on sugar beet growth. We also surveyed growth sensitivity of the *Streptomyces* isolates to temperature and pH ranges. To assess the mode of action of antagonism, chitinase activity and siderophore production of active isolates were ruled out.

MATERIALS AND METHODS

Bacterial and fungal strains: Isolates of *Streptomyces* designated as S2 and C were used in this study. The strains were isolated from a soil sample collected from a wheat field at Kerman, Iran as described by Bonjar *et al.* (2005) and maintained lyophilized before use. Bacteria were revived and cultured on MYA medium containing 10 g Malt extract, 4 g Yeast extract, 4 g Glucose and 15 g agar, adjusted to pH 7.2 and incubated at 29°C for 7-10 days. These cultures were stored at 4°C until used. *R. solani* AG-4 (Rs1, Rs2 and Rs3 isolates) were provided by Plant Diseases Laboratory of Plant Pests and Diseases Research Institute, Karaj, Iran. Fungal stock cultures were maintained on PDA (Potato Dextrose Agar) at 23°C.

Effect of different temperatures and pH on bacterial growth: The growth behavior of *Streptomyces* isolates (S2 and C) was evaluated on MYA medium at temperatures of 18, 23, 29 and 37°C and pH ranges of 5.0, 5.6, 7.2 and 8.0.

In vitro antifungal bioassay: Procedure of Yuan and Crawford, (1995) was used to examine the biocontrol activity of *Streptomyces* isolates S2 and C against *R. solani* isolates Rs1, Rs2 and Rs3. A loopful of each *Streptomyces* isolate spores was streaked onto one side of each PDA plate. The plates were incubated at 29°C for 72 h. A 0.45 cm diameter agar plug from the growing margin of fungal colonies was transferred onto the center of the each plate. Fungal plugs were also placed on uninoculated PDA plates separately as uninhibited controls. Cultures were incubated at 23°C. Inhibitory effect of the bacterial isolates on fungal mycelium growth was estimated by subtraction of the fungal growth radius of a control culture (X₀ in centimeters) from the distance of the growth in the direction of *Streptomyces* isolates (X in centimeters) after 5 days in dual cultures.

Viability test of treated fungal hyphae: Fungicidal/fungistatic effect of the antagonists, at the end of 5th day of paired incubation, mycelial plugs from the edge of fungal colony in the vicinity of inhibitory zones were transferred to fresh medium for evaluation of the viability of the mycelia in that area.

Preparation of inoculum for in vivo studies: Dehulled wheat seeds were soaked overnight in tap water, excess of water was removed and the wet seeds were autoclaved twice at 121°C for 20 min, cooled and used as basal medium. For inoculation, pieces of agar with actively growing hyphae of *R. solani* were placed 2-3 cm below the surface of the medium in 1 L Erlenmeyer flasks. The flasks were shaken every 2 days. After four weeks of incubation

at 23°C, the medium was completely covered with the fungal mat. A mixture of the colonized medium, sterilized corn flour and sterilized sand (5, 5 and 90% w/w, respectively) was used as fungal inoculum for *in vivo* investigations.

Formulation of streptomyces isolates (S2 and C): A spore suspension of each *Streptomyces* isolates S2 and C was used for *in vivo* antagonism test. Spores and mycelia from 3 to 6 well grown MYA plates were scraped off of the surfaces with a sterile spatula and added to 20 mL of sterile physiological serum (0.9% NaCl). Spore suspension of each bacterial isolate was added to sterilized sand and mixed carefully, then the formulation was stored at 42°C for 2 h. One g sample from each isolate formulation was used to make serial dilutions in sterile physiological serum and was cultured on MYA plates for measurement of CFU g⁻¹ sand.

Soil treatment and bioassay study: Sterilized soil was infected with inoculum at a rate of 1:10 w/w. In uninoculated pots mixture of 5% w/w sterilized corn flour and 95% w/w sterilized sand was used instead of inoculum. The infected and uninfected soils were placed in 20 cm diameter plastic pots. For plants treated with both bacteria and fungus or plants treated with only bacteria, 80 g (10⁶ CFU g⁻¹ sand) from each formulation was added to inoculated and uninoculated pots. Eighty gram from sand treated with 0.9% sterile physiological serum was added to uninfected soil as control. Pots were prepared in replicates of three, in all treatments and were arranged in a complete randomized design in the greenhouse at 23°C±1 and cycles of 12 h light and 12 h darkness. After 2 days, each pot was planted with 8 monogerm sugar beet seeds and watered every 2 days. After 14 days the numbers of healthy seedlings were recorded. Two months after planting, the plants were harvested and root volume, dry weight of leave, stem and root were measured.

Chitinase activity: Three 0.45 cm diameter agar plug containing mycelium of each *Streptomyces* isolates (S2 and C) were placed on the surface of 9-cm petri plates containing 0.4% colloidal chitin (Hsu and Lockwood, 1975) and 1.5% agar at pH 7.2 and incubated at 29°C. After 3, 5, 7 and 10 days incubation, clear zones (>5 mm in diameter) around each colony was recorded and used as indicator of chitinolytic activity. There were three replicates and the experiments were repeated.

Siderophore detection: A procedure of Schupp *et al.* (1988) was used to siderophore detection. In brief, Whatman No. 1 filter paper soaked in indicator solution containing 1% ammonium ferric sulfate

(FeNH₄(SO₄)₂.12H₂O) in sulfuric acid was put on 4-days single colonies of S2 and C grown up on Des4 medium containing 2% Dextrin, 2% Manitol, 1.2% L-asparagine, 0.025% L-lysine, 0.01% L-methionine, 0.01% L-threonine, 0.5% CaCO₃, 0.025% MgSO₄, 0.05% K₂HPO₄, 0.005% ZnSO₄ and 2% agar. After 15 min brown or reddish brown halo around colonies were recorded and used to detect siderophore producing *Streptomyces* isolate.

Statistical analysis: Analysis of variance was performed based on used experimental design. Mean comparison was carried out using Duncan's multiple range test. MSTATC statistical package was used for the analyses.

RESULTS

Bacterial growth at different pHs and temperatures: Both bacterial isolates grew well at 29°C in pHs of 5.6, 7.2 and 8.0. The isolate C grew in acidic (pH 5.6), neutral (pH 7.2) and alkaline (pH 8.0) media at 23 to 37°C. The isolate S2 grew at 18 to 37°C in pH 5.6 to 8.0 but did not grow in acidic medium at 37°C. When pH decreased to 5.0, the growth of C and S2 isolates was decreased and inhibited, respectively.

In vitro antagonism test: The *in vitro* antagonism assays showed that both isolates, S2 and C had inhibitory effect on growth of fungal mycelium. The antagonistic effects were appeared after two days of paired bacteria and fungi incubation and advanced during the days after incubation. Comparison of inhibition zones of S2 and C illustrated that isolate C with an inhibition zone more than 2 cm had nearly two fold more inhibitory effect than S2 with an inhibition zone of less than 1 cm. Since *Streptomyces* have a relatively lower growth rate on agar plates than most of the fungi, preinoculation (72 h) was done to establish bacterial colonies on the agar surface first. Although, inhibition was appeared after two days of simultaneous culture of bacteria and fungi. The viability tests showed that after 5 days of incubation, mycelium of *Rhizoctonia* isolates (RS1, RS2 and RS3) taken from peripheral areas of the inhibited colonies facing S2 or C, were culturable on fresh agar medium.

Chitinase activity: Clear zone around S2 and C appeared 3 and 5 days after incubation on chitin agar, respectively indicated that both isolates were able to use chitin as a carbon source. Isolate C grew as fast as isolate S2 on chitin agar but did not appear clear zone before 5 days.

Siderophore production: Reaction between siderophore and iron existed in indicator solution ((FeNH₄(SO₄)₂.12H₂O)) creates a colored halo around




Fig. 1: Effect of *Streptomyces* spp. isolates S2 and C on plant stand of sugar beet seedlings grown in soil infected with *R. solani* spp. isolate (RS1, RS2 and RS3). Means in each column followed by the same letter are not significantly different from each other according to Duncan's Multiple Range Test ($p < 0.05$)




Fig. 2: Effect of *Streptomyces* spp. isolate S2 and C on sugar beet dry matter and root density in soil infected with *R. solani* spp. isolate RS1, RS2 and RS3. Numbers in each column followed by the same letter are not significantly different from each other according to Duncan's Multiple Range Test ($p < 0.05$)

siderophore producing bacteria. According to that, after covering Des4 medium with Whatman paper soaked in indicator solution the halo appeared around isolate C colonies, but not for isolate S2.

Shelf life of the S2 and C formulation and recovery of bacteria from soil: Density of spores in both formulations of S2 and C was 10^6 CFU g^{-1} sand before adding to pots. After 1 months of storage at room temperature ($25^{\circ}C \pm 2$), the density of spores in both formulations have declined to 10^5 CFU g^{-1} sand. The level of viability was increased up to 10^6 CFU/g for S2 and decreased to 10^4 CFU g^{-1} for C in samples collected from 3 cm from the soil surface 1 month after treatment with the formulations.

Bioassay study: Soil treatment with S2 and C formulation increased seed germination (16.6 and 25% respectively) from uninfected soil significantly ($p < 0.05$). *Streptomyces* isolates S2 and C inhibit Rhizoctonia damping-off completely (Fig. 1). All treatments containing bacteria increased shoot and root dry weight and root density compared to control (Fig. 2).

DISCUSSION

Enhancement of fungal biological control agents may improve alternative measures for chemical control management. In this way, several attempts have been made to use the biocontrol approach against major diseases in crop plants.

Several studies have reported the antagonistic effects of *Trichoderma* spp. (Abada, 1994), *Pseudomonas* spp. (Shah-Smith and Burns, 1996; Georgakopoulos *et al.*, 2002) and *Streptomyces* spp. (Crawford *et al.*, 1993; Yuan and Crawford, 1995; Paulitz and Belanger, 2001) against *Pythium* spp. causal agents of sugar beet damping-off. Biocontrol of Rhizoctonia damping off with *Streptomyces* strains were published for crops, Chinese cabbage by Chung *et al.* (2005) and tomato by Sabaratnam and Traquair (2002). Also there is a commercial product, Mycostop, driven from *Streptomyces griseoviridis* has been registered to control of crop damping-off (Lahdenpera *et al.*, 1991). The data presented in this study demonstrated that two *Streptomyces* isolates

Table 1: Effect of pH (5.0-8.0) and temperature (18-37°C) on growth of *Streptomyces* spp. isolates S2 and C

Temp (°C)	pH							
	5.0		5.6		7.2		8.0	
	S2	C	S2	C	S2	C	S2	C
18	-	-	M*	-	M	-	M	-
23	-	M	M	M	M	M	M	M
29	-	M	W**	W	W	W	W	W
37	-	M	-	W	W	W	W	W

M*: Moderate growth; W**: Well growth

(S2 and C) had both *in vitro* and *in vivo* antagonistic effects against *R. solani*, the causal agent of sugar beet damping-off.

Our result showed there is relationship between *in vitro* antagonism and *in vivo* disease suppression for potential biocontrol *Streptomyces* as reported by the others (Yuan and Crawford, 1995; El-Tarabily, 2000).

The *in vitro* antagonism assays revealed that three pathogenic *R. solani* isolates were sensitive to the antifungal metabolites released by S2 and C in solid media. Isolate S2 with a smaller zone of inhibition than C was completely able to suppress the disease similar to C based on results from *in vivo* experiments. It seems that results from *in vitro* study cannot simply be up scaled to *in vivo* tests. On the other words, *in vitro* screening of biocontrol agents should be confirmed using *in vivo* assays. It may help us to save some beneficial isolates are less effective in *in vitro* assays.

Soil treatment with both two biocontrol agents increased plant dry matter and root density compared with those were cultured in untreated (with biocontrol agent formulation), infected and uninfected (control) soils (Fig. 2).

Isolates C and S2 showed that differentially promote sugar beet growth. Isolate C increased seed germination (25%) rather than S2 (16%) and total dry matters were more for plants grown in soils treated with C than S2 in infected soil.

Since isolate C, is a siderophore producing and S2 isn't, it may be concluded that a part of efficiency of C is related to siderophore production.

Dicotyledonous plants have been shown to acquire Fe⁺ from microbial chelates, siderophores (Crowley *et al.*, 1991). Direct utilization of microbial siderophores as iron sources examined and its effect in preventing iron stress and increasing plant biomass reported previously (Wang *et al.*, 1993). On the other hand, siderophore is one of the compounds has been indicated to cause Induced Systemic Resistance (ISR) in plants providing protection against a broad spectrum of phytopathogenic organisms (Bloembergen and Lugtenberg, 2001). Tokala *et al.* (2002) demonstrated that *Streptomyces lydicus* WYEC108, an

antifungal biocontrol agent, is also a plant growth-promoting bacterium producing hydroxamate-type siderophores. With basis of the results it may point out that siderophore production has a positive correlation with plant growth promotion.

Chitin degradation by isolates S2 and C have been demonstrated on colloidal chitin but its involvement in lyses of living mycelia was not studied. Although, according to De Boer *et al.* (1998) and Gupta *et al.* 1995, it seems that the basic mechanism for chitinolytic bacteria against fungi is antibiosis. On the other hand, because mycelial plugs from the fungal colony edge facing S2 and C colonies were recovered on fresh medium, it may be suggested that fungistatic metabolites secretion by antagonist isolates which diffused in solid agar had the most inhibitory effects on *R. solani* isolates.

Isolates C and S2 grew well at 29°C in pH 7.2, that have been cited as optimal conditions for *Streptomyces* (Locci, 1994). At 37°C colonies appeared faster and at lower temperature, 23°C both isolates grew moderately. When temperature was decreased to 18°C, the growth of S2 did not change but the growth of C was inhibited (Table 1). Hence both isolates grew moderately to well at temperatures between 25-33°C which *R. solani* is active (Gallian, 2001) and is suitable for sugar beet. Isolate C at 18°C and isolate S2 in pH 5.0 were inhibited, consequently integration of these two isolates may be more effective to biocontrol of *R. solani* at some environmental conditions that is not suitable for S2 or C.

Based on these results a biocontrol composite prepared from these two bacterial isolates probably should be viable and active in wide range of temperature (18-37°C) and acidity (pH 5.0-8.0) of planting soils.

Since isolates C and S2 showed inhibitory effects on different genera of fungi (*Rhizoctonia*, *Fusarium* and *Pyricularia*) (unpublished data) seems that these two isolates may be able to suppress all anastomosis groups of *Rhizoctonia solani* included of the major AG worldwide, *R. solani* AG-2-2, causes root and crown rot (Scholten *et al.*, 2001) too. Therefore these two isolates have the potential to be developed into commercial biofungicides in control of sugar beet *Rhizoctonia* diseases.

We hope that further study lead us to identify the mode of action and genes involved in antifungal activity of these isolates.

ACKNOWLEDGMENTS

We thank Reza Ostadsaraie for technical assistance. This work was supported by the Agricultural Biotechnology Research Institute of Iran (ABRII).

REFERENCES

- Abada, K.A., 1994. Fungi causing damping-off and root-rot on sugar-beet and their biological control with *Trichoderma harzianum*. Agric. Ecosys. Environ., 51: 333-337.
- Abd-Allah, E.F., 2001. *Streptomyces plicatus* as a model biocontrol agent. Folia Microbiol., 46: 309-314.
- Aghighi, S., G.H. Shahidi Bonjar and I. Saadoun, 2004. First report of antifungal Properties of a new strain of *Streptomyces plicatus* (Strain101) against four Iranian phytopathogenic isolates of *Verticillium dahliae*, a new horizon in biocontrol agents. Biotechnology, 3: 90-97.
- Asaka, O. and M. Shoda, 1996. Biocontrol of *Rhizoctonia solani* damping-off of tomato with *Bacillus subtilis* RB14. Applied Environ. Microbiol., 62: 4081-4085.
- Bloemberg, G.V. and B.J.J. Lugtenberg, 2001. Molecular basis of plant growth promotion and biocontrol by rhizobacteria. Curr. Opin. Plant Biol., 4: 343-350.
- Bonjar, G.H.S., P.R. Farrokhi, S. Aghighi, L.S. Bonjar and A. Aghelizadeh, 2005. Antifungal characterization of actinomycetes isolated from Kerman, Iran and their future prospects in biological control strategies in greenhouse and field conditions. Plant Pathol. J., 4: 78-84.
- Bormann, C., D. Baier, I. Horr, C. Raps, J. Berger, G. Jung and H. Schwarz, 1999. Characterization of a novel, antifungal, chitin-binding protein from *Streptomyces tendae* Tu901 that interferes with growth polarity. J. Bacteriol., 181: 7421-7429.
- Chamberlain, K. and D.L. Crawford, 1999. *In vitro* and *in vivo* antagonism of pathogenic turfgrass fungi by *Streptomyces hygroscopicus* strains YCED9 and WYE53. J. Ind. Microbiol. Biotechnol., 23: 641-646.
- Chung, W.C., J.W. Huang and H.C. Huang, 2005. Formulation of a soil biofungicide for control of damping-off of Chinese cabbage (*Brassica chinensis*) caused by *Rhizoctonia solani*. Biol. Control, 32: 287-294.
- Crawford, D.L., J.M. Lynch, J.M. Whipps and M.A. Ousley, 1993. Isolation and characterization of actinomycete antagonists of a fungal root pathogen. Applied Environ. Microbiol., 59: 3899-3905.
- Crowley, D.E., Y.C. Wang, C.P.P. Reid and P.J. Szanislo, 1991. Mechanisms of iron acquisition from siderophores by microorganisms and plants. Plant Soil, 130: 179-198.
- De Boer, W., P.J.A. Klein Gunnewiek, P. Lafeber, J.D. Janse, B.E. Spit and J.W. Woldendorp, 1998. Antifungal properties of chitinolytic dune soil bacteria. Soil Biol. Biochem., 30: 193-203.
- EL-Abyad, M.S., M.A. EL-Sayed, A.R. EL-Shanshoury and S.M. EL-Sabbagh, 1993. Toward the biological control of fungal and bacterial diseases of tomato using antagonistic *Streptomyces* spp. Plant Soil, 149: 185-195.
- El-Tarabily, K.A., M.H. Soliman, A.H. Nassar, H.A. Al-Hassani, K. Sivasithamparam, F. McKenna and G.E.St.J. Hardy, 2000. Biological control of *Sclerotinia minor* using chitinolytic bacterium and actinomycetes. Plant Pathol., 49: 573-583.
- Ensign, J.C., 1978. Formation, properties and germination of actinomycete spores. Ann. Rev. Microbiol., 32: 185-219.
- Gallian, J.J., 2001. Management of sugar beet root rots. Pacific Northwest Publication. University of Idaho. Washington State University.
- Georgakopoulos, D.G., P. Fiddaman, C. Leifert and N.E. Malathrakis, 2002. Biological control of cucumber and sugar beet damping-off caused by *Pythium ultimum* with bacterial and fungal antagonists. J. Applied Microbiol., 92: 1078-1086.
- Getha, K. and S. Vikineswary, 2002. Antagonistic effects of *Streptomyces violaceusniger* strain G10 on *Fusarium oxysporum* f. sp. cubense race 4: Indirect evidence for the role of antibiosis in the antagonistic process. J. Ind. Microbiol. Biotechnol., 28: 303-310.
- Gupta, R., R.K. Saxena, P. Chaturvedi and J.S. Virdi, 1995. Chitinase production by *Streptomyces viridificans*: Its potential in fungal cell wall lysis. J. Applied Bacteriol., 78: 378-383.
- Gupte, T.E. and S.R. Naik, 1999. Isolation, taxonomic and fermentation studies on a new strain of *Streptomyces arenae* var *ukrainiana* producing a tetraene antibiotic. World J. Microbiol. Biotechnol., 15: 545-552.
- Hsu, S.C. and J.L. Lockwood, 1975. Powered chitin agar as a selective medium for enumeration of actinomycetes in water and soil. Applied Microbiol., 29: 422-426.
- Hwang, B.K., S.W. Lim, B.S. Kim, J.Y. Lee and S.S. Moon, 2001. Isolation and *in vivo* and *in vitro* antifungal activity of phenylacetic acid and sodium phenylacetate from *Streptomyces humidus*. Applied Environ. Microbiol., 67: 3739-3745.
- Jones, C.R. and D.A. Samac, 1996. Biological control of fungi causing alfalfa seedling damping-off with a disease-suppressive strain of *Streptomyces*. Biol. Control, 7: 196-204.

- Lahdenpera, M.L., E. Simon and J. Uoti, 1991. Mycostop-A Novel Biofungicide Based on Streptomyces Bacteria. In: Biotic Interactions and Soil-Borne Disease, Volume 23, Beemster, A.B.R., G.J. Bollen, M. Gerlagh, M.A. Ruissen, B. Schippers and A. Tempel (Eds.), Elsevier, Amsterdam, The Netherlands, pp: 258-263.
- Locci, R., 1994. *Streptomyces* and related genera. In Bergeys manual of systematic bacteriology (Eds., Williams, S.T., M.E. Sharpe and J.G. Holt), Williams and Wilkins, Baltimore, 4: 2451-2508.
- Moussa, T.A.A., 2002. Studies on biological control of sugar beet pathogen *Rhizoctonia solani* Kühn. Online J. Biol. Sci., 2: 800-804.
- Paulitz, T.C. and R.R. Belanger, 2001. Biological control in greenhouse systems. Ann. Rev. Phytopathol., 39: 103-133.
- Nassar, A.H., A. Khaled, A. El-Tarabily and K. Sivasithamparam, 2003. Growth promotion of bean (*Phaseolus vulgaris* L.) by a polyamine-producing isolate of *Streptomyces griseoluteus*. Plant Growth Regul., 40: 97-106.
- Sabaratham, S. and J.A. Traquair, 2002. Formulation of a Streptomyces biocontrol agent for the suppression of *Rhizoctonia* damping-off in tomato transplants. Biol. Control, 23: 245-253.
- Schmidt, C.S., F. Agostini, C. Leifert, K. Killham and C.E. Mullins, 2004. Influence of soil temperature and matric potential on sugar beet seedling colonization and suppression of *Pythium* damping-off by the antagonistic bacteria *Pseudomonas fluorescens* and *Bacillus subtilis*. Phytopathology, 94: 351-363.
- Scholten, O.E., L.W. Panella, T.S.M. De Bock and W. Lange, 2001. A greenhouse test for screening sugar beet (*Beta vulgaris*) for resistance to *Rhizoctonia solani*. Eur. J. Plant Pathol., 107: 161-166.
- Schupp, C., C. Toupet and M. Divers, 1988. Cloning and Expression of two genes of *Streptomyces pilosus* involved in the biosynthesis of the siderophore desferrioxamine B. Gene, 64: 179-188.
- Shah-Smith, D.A. and R.G. Burns, 1996. Biological control of damping-off of sugar beet by *Pseudomonas putida* applied to seed pellets. Plant Pathol., 45: 572-582.
- Shih, H.D., Y.C. Liu, F.L. Hsu, V., Mulabagal, R. Dodda and J.W. Huang, 2003. Fungichromin: a substance from *Streptomyces padanus* with inhibitory effects on *Rhizoctonia solani*. J. Agric. Food Chem., 51: 95-99.
- Smith, J., A. Putnam and M. Nair, 1990. *In vitro* control of *fusarium* diseases of *Asparagus officinalis* L. with a Streptomyces or its polyene antibiotic, faeriefungin. J. Agric. Food Chem., 38: 1729-1733.
- Tokala, R.K., J.L. Strap, C.M. Jung, D.L. Crawford, M.H. Salove, L.A. Deobald, J.F. Bailey and M.J. Morra, 2002. Novel plant-microbe rhizosphere interaction involving *Streptomyces lydicus* WYEC108 and the pea plant (*Pisum sativum*). Applied Environ. Microbiol., 68: 2161-2171.
- Trejo-Estrada, S.R., I.R. Sepulveda and D.L. Crawford, 1998. *In vitro* and *in vivo* antagonism of *Streptomyces violaceusniger* YCED9 against fungal pathogens of turfgrass. World J. Microbiol. Biotechnol., 14: 865-872.
- Walker, R., 2004. The delivery of biological control agents to the sugar beet crop. British Sugar Beet Rev., 72: 20-24.
- Wang, Y., H.N. Brown, D.E. Crowley and P.J. Szanislo, 1993. Evidence for direct utilization of a siderophore, ferroxamine B, in axenically grown cucumber. Plant Cell Environ., 16: 579-585.
- Whipps, J.M., 2001. Microbial interactions and biocontrol in the rhizosphere. J. Exp. Bot., 52: 487-511.
- Xiao, K., L.L. Kinkel and D.A. Samac, 2002. Biological control of *Phytophthora* root rots on alfalfa and soybean with *Streptomyces*. Biol. Control, 23: 285-295.
- Yuan, W.M. and D.L. Crawford, 1995. Characterization of *Streptomyces lydicus* WYEC108 as a potential biocontrol agent against fungal root and seed rots. Applied Environ. Microbiol., 61: 3119-3123.