

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan

Screening Plant Growth Promoting Rhizobacteria for Improving Seed Germination, Seedling Growth and Yield of Maize

S. Nezarat and A. Gholami
Department of Agronomy, College of Agriculture,
Shahrood University of Technology, Shahrood, Iran

Abstract: The effect of Plant Growth-Promoting Rhizobacteria (PGPR) on seed germination, seedling growth and yield of field grown maize were evaluated in three experiments. In these experiments six bacterial strains include *P. putida* strain R-168, *P. fluorescens* strain R-93, *P. fluorescens* DSM 50090, *P. putida* DSM291, *A. lipoferum* DSM 1691 and *A. brasilense* DSM 1690 were used. Results of first study showed seed inoculation significantly enhanced seed germination and seedling vigour of maize. In second experiment, leaf and shoot dry weight and also leaf surface area significantly were increased by bacterial inoculation in both sterile and non-sterile soil. The results showed that inoculation with bacterial treatments had a more stimulating effect on growth and development of plants in nonsterile than sterile soil. In the third experiment, Inoculation of maize seeds with all bacterial strains significantly increased plant height, 100 seed weight, number of seed per ear and leaf area. The results also showed significant increase in ear and shoot dry weight of maize.

Key words: Maize, PGPR, *Azospirillum*, *Pseudomonas*, biofertilizer

INTRODUCTION

During the last couple of decades given the negative environmental impact of chemical fertilizers and their increasing costs, the use of soil microorganisms for sustainable agriculture has increased in various parts of the world. Plant Growth Promoting Rhizobacteria (PGPR) are a group of bacteria that actively colonize plant roots and increase plant growth and yield (Wu *et al.*, 2005). The main active strains of PGPR belong to a range of genera, including *Azospirillum*, *Azotobacter*, *Bacillus*, *Pseudomonas* and *Serratia* (Bashan *et al.*, 2004; Ramamoorthy *et al.*, 2002; Zhang *et al.*, 1996). The mechanisms by which PGPRs promote plant growth are not fully understood, but are thought to include: the ability to produce phytohormones (Egamberdiyeva, 2007; Shaharoon, 2006), symbiotic N₂ fixation (Mrkovački and Milic, 2001; Salantur *et al.*, 2006), against phytopathogenic microorganisms by production of siderophores, the synthesis of antibiotics, enzymes and/or fungicidal compounds (Ahmad *et al.*, 2006; Bharathi *et al.*, 2004; Jeun *et al.*, 2004) and also solubilisation of mineral phosphates and other nutrients (Cattelan *et al.*, 1999).

Significant increases in growth and yield of agronomical important crops in response to inoculation with PGPR have been reported by Asghar *et al.* (2002),

Bashan *et al.* (2004) and Biswas *et al.* (2000). *Azospirillum*, *Pseudomonas* and *Azotobacter* strains could affect seed germination and seedling growth (Shaukat *et al.*, 2006). Kloepper *et al.* (1992) has been shown that wheat yield increased up to 30% with *Azotobacter* inoculation and up to 43% with *Bacillus* inoculation. Strains of *Pseudomonas putida* and *Pseudomonas fluorescens* could increase root and shoot elongation in canola, (Glick *et al.*, 1997) as well as wheat and potato (De Freitas and Germida, 1992; Frommel *et al.*, 1993). Inoculation of plants with *Azospirillum* could result in significant changes in various growth parameters, such as increase in plant biomass, nutrient uptake, tissue N content, plant height, leaf size and root length of cereals (Bashan *et al.*, 2004). Thus it has been shown that *Azospirillum* and *Pseudomonas* had the potential for agricultural exploitation and could use as natural fertilizers (Bashan *et al.*, 1989; Cakmakc *et al.*, 2006).

However, the effects of these bacteria on growth and yield of some crop plants studied in earlier studies. But the effects of PGPR on growth parameter from germination to yield were not evaluated simultaneously. The main objective of this research was to determine if PGPR strains could affect on seed germination, growth parameters of maize seedling in greenhouse and also grain yield of field grown maize.

MATERIALS AND METHODS

PGPR strains and inoculum's preparation: Six bacterial strains include *P. putida* strain R-168, *P. fluorescens* strain R-93, *P. fluorescens* DSM 50090, *P. putida* DSM291, *A. lipoferum* DSM 1691 and *A. brasilense* DSM 1690 were obtained from the Soil Microbiology Department, Soil and Water Research, Tehran and were used in this study that conducted at 2005-2006.

Mode of application: Bacterial strains were used as maize seed treatments. Seeds of maize (SC 647) were surface-sterilized with 0.02% sodium hypochlorite for 2 min and rinsed thoroughly in sterile distilled water. For inoculation seeds were coated with 20% gum arabic as an adhesive and rolled into the suspension of bacteria (10^8 cfu mL⁻¹) with perlite until uniformly coated. (Salanture *et al.*, 2006). Seeds treated with sterile distilled water amended with gum arabic served as the nontreated control.

Seed germination bioassay: Germination tests were carried out by the paper towel method (Niranjan *et al.*, 2004). Twenty five seeds for each treatment with three replications in completely randomized design and incubated in growth chamber at 28°C. After 7 days the number of germinated seeds was counted. Root and shoot length of individual seedling was measured to determine the vigor index with following formula (Abdul Baki and Anderson, 1973):

$$\text{Vigor index} = (\text{Mean root length} + \text{Mean shoot length}) \times \text{Germination (\%)}$$

Pot culture experiment: For the evaluation of maize seedling growth promotion with PGPRs, above bacterial strains were tested in both non-sterile and sterile soils at 2005. The plastic pots had 15 cm diameter and capacity to hold 2 kg of soil. For preparation of sterile soil, field soil was autoclaved twice for 20 min at 120°C with a 24 h interval. All treatments (bacterial inoculation × soil condition) arranged in 48 pots i.e., 3 replicates with 14 pots per replication and a double seed per pot. Treatments were arranged in a factorial experiment based on completely randomized design. Seedlings were watered daily and no artificial fertilization was used. After 30 days, fresh weight was determined and dry weight calculated by drying plants in an oven at 75°C until the weight remained constant. For leaf area determination, the area of each expanded leaf was calculated as:

$$K \times \text{length} \times \text{width}$$

Where:

$$k = 0.75 \text{ (Ruget } et al., 1996).$$

Field experiment: Field experiment was conducted at research farm of Shahrood University of technology (latitude of 36° 25 'N and longitude of 54° 57' E with an elevation of 1345 m) in the period of May-September, 2006. The field soil was silty clay loam in texture, having pH, 7.8; EC, 3.9 dS m⁻¹; 0.75% of organic carbon; 0.04% N, 6.4 and 320 ppm of available P and K, respectively. Seeds of maize were washed with distilled water then inoculation was performed by a suspension of any bacteria (10^8 cfu mL⁻¹) with perlite mixture. Treatments were arranged as randomized complete block design with three replications. There were four rows in each plot. Which the row width and length was 0.7 and 9 m, respectively. Before sowing, the soil was fertilized with N, P and K at rate of 300, 150 and 50 kg ha⁻¹ as urea, single super phosphate and potassium sulphate, respectively. Half of nitrogen was applied at sowing time and residue at the start of reproductive stage. Seeds were placed at 5 cm depth. At the third leaf stage, plants were thinned to 1 plant hill⁻¹ for the appropriate final stand of 75000 plants ha⁻¹. The soil was subsequently irrigated 10 days interval at the first and shortened to 7 days interval to summer. Ten mature maize plants were sampled from each treatment for final measurements in October (120 days after sowing). Leaf surface area was measured as describe above. In laboratory, samples were separated into different components and oven-dried at 75°C until reached to constant weight.

Data analysis: Data were statistically treated by ANOVA, Least Significant Difference (LSD) test at probability level 0.05 was used to separate the means when the ANOVA F-test indicated a significant effect of the treatments.

RESULTS

Seed germination: Seed inoculation significantly enhanced seed germination and seedling vigour of maize. However, the rate of enhancement varied with bacterial strains. All bacteria except *A. lipoferum* DSM1691, increased seed germination up to 18.5% over nontreated control (Fig. 1). The highest enhancement of vigor indexes were obtained from *A. brasilense* DSM 1690 and *P. putida* strain R-168, which recorded 975 and 873 vigor index, respectively.

Effects of inoculation with bacteria on maize plant growth and development: The results of pot study showed that inoculation of maize seeds with bacterial strains did not affect leaf fresh weight and stem dry weight (Table 1). In

Table 1: Effect of bacterial inoculation on growth characteristics of maize seedlings at 30 days after sowing in different soil condition

| Treatments | Fresh weight (g) | | | Dry weight (g) | | | Leaf area (cm ²) |
|--|------------------|---------|---------|----------------|------|----------|------------------------------|
| | Leaf | Stem | Total | Leaf | Stem | Shoot | |
| <i>P. putida</i> strain R-168 | | | | | | | |
| Non sterile | 5.28* | 4.55abc | 9.84abc | 0.29ab | 0.62 | 0.900abc | 250.60ab |
| Sterile | 3.57 | 3.20abc | 6.78abc | 0.27ab | 0.57 | 0.840abc | 173.10bc |
| <i>P. fluorescens</i> strain R-93 | | | | | | | |
| Non sterile | 5.04 | 4.49abc | 9.53abc | 0.33ab | 0.61 | 0.940abc | 274.60ab |
| Sterile | 1.42 | 1.15c | 2.56c | 0.11b | 0.38 | 0.490abc | 109.30 |
| <i>P. fluorescens</i> DSM50090 | | | | | | | |
| Non sterile | 5.09 | 4.31abc | 9.31abc | 0.27ab | 0.59 | 0.860abc | 244.10ab |
| Sterile | 4.32 | 3.80abc | 8.12abc | 0.26ab | 0.55 | 0.810abc | 169.20bc |
| <i>P. putida</i> DSM291 | | | | | | | |
| Non sterile | 5.28 | 4.55abc | 9.83abc | 0.34ab | 0.65 | 0.990abc | 252.40ab |
| Sterile | 3.56 | 2.89abc | 6.45abc | 0.17ab | 0.43 | 0.600abc | 166.70bc |
| <i>A. lipoferum</i> DSM1691 | | | | | | | |
| Non sterile | 6.13 | 6.07a | 12.20a | 0.52a | 0.61 | 1.130ab | 350.00a |
| Sterile | 1.50 | 1.57 bc | 3.07c | 0.08b | 0.20 | 0.280bc | 87.50c |
| <i>A. brasilense</i> DSM1690 | | | | | | | |
| Non sterile | 6.32 | 5.39ab | 11.72ab | 0.36ab | 0.72 | 1.080ab | 267.30ab |
| Sterile | 6.65 | 5.62a | 12.27a | 0.37ab | 0.93 | 1.300a | 325.40a |
| Control | | | | | | | |
| Non sterile | 2.47 | 1.08c | 3.88c | 0.05b | 0.12 | 0.170c | 196.30bc |
| Sterile | 2.45 | 2.13abc | 4.58bc | 0.18ab | 0.36 | 0.540abc | 170.00bc |
| LSD (0.05) | - | 4.002 | 7.607 | 0.385 | - | 0.857 | 112.20 |

*In a column, means followed by a common letter(s) are not significantly different at 5% level by LSD




Fig. 1: Effects of bacterial inoculations on seed germination and vigor index of maize 7 days after germination *in vitro* conditions

contrast, stem and total fresh weight significantly increased by inoculation in sterile and non-sterile soil. The highest stem and total fresh weight were recorded from *A. lipoferum* DSM 1691 in non sterile soil and from *A. brasilense* DSM 1690 in sterile soil. Leaf and shoot dry weight were significantly enhanced by bacterial inoculation and soil type ($p < 0.05$). Application of *A. lipoferum* DSM 1691 in non sterile soil and *A. brasilense* DSM 1690 in sterile soil, respectively had the best effect on leaf dry weight (0.52 g plant⁻¹) and shoot dry weight (1.30 g plant⁻¹). Furthermore, inoculation with bacterial strains had significant effect on leaf surface area under both soil conditions. The results

revealed that in non sterile soil, *A. lipoferum* DSM 1691 caused an increase of 78.3% in leaf area while in sterile soil, leaf area was increased up to 65% in response to inoculation with *A. brasilense* DSM 1690 compared to control.

The results showed that inoculation with bacterial treatments had a more stimulating effect on growth and development of plants in nonsterile soil than sterile condition. *P. putida* strain R-168 and *A. lipoferum* DSM 1691 performed better than other strains in stimulating plant growth in pot experiment especially at nonsterile soil.

Inoculation of maize seeds with all bacteria strains significantly increased the plant height (14.3-21.7%) and leaf area (Table 2). However, seed dry weight increased with bacterial inoculation, but there were no significant differences between all treatments and control. 100 seed weight was significantly affected due to treatments with *Azospirillum* and *Pseudomonas* strains over the control. The most effective strain was *P. fluorescens* DSM50090 which increased 100 seed weight up to 44% over control. Effect of inoculation on number of seed per ear was also significant. Application of *P. fluorescens* strain R-93 and *A. lipoferum* DSM1691 gave the maximum number of seed per ear.

The results showed a clear and significant increase in ear and shoot dry weight of maize ($p < 0.05$). The increase in these parameters, with different inoculation ranged from *P. putida* DSM291 (up to 70%) to *A. lipoferum* DSM1691 (up to 100 %) when compared with control (Table 2).

Table 2: Effect of seed treatment with PGPR on growth parameters of maize under field conditions

| Treatments | Plant height (cm) | Leaf area (cm ²) | Seed dry weight (g plant ⁻¹) | 100-seed weight (g) | No. of seed ear ⁻¹ | Ear dry weight (g plant ⁻¹) | Soot dry weight (g plant ⁻¹) |
|-----------------------------------|-------------------|------------------------------|--|---------------------|-------------------------------|---|--|
| <i>P. putida</i> strain R-168 | 183.00a* | 5838.0a | 195.21 | 25.84bcd | 761.36ab | 241.37a | 340.66a |
| <i>P. fluorescens</i> strain R-93 | 178.33a | 5762.5a | 226.41 | 24.35cd | 867.36a | 267.16a | 351.90a |
| <i>P. fluorescens</i> DSM50090 | 185.00a | 6137.7a | 203.65 | 31.00a | 736.23ab | 258.26a | 355.45a |
| <i>P. putida</i> DSM291 | 176.58a | 6322.7a | 190.63 | 30.31ab | 616.88bc | 235.16a | 331.20a |
| <i>A. lipoferrum</i> DSM1691 | 187.75a | 5968.0a | 231.47 | 28.41abc | 814.79a | 280.95a | 388.98a |
| <i>A. brasilense</i> DSM1690 | 176.33a | 5628.0a | 205.02 | 29.98ab | 685.83abc | 259.63a | 355.97a |
| Control | 154.25b | 2176.0b | 109.00 | 21.41d | 509.56c | 133.72b | 192.97b |
| LSD (0.05) | 19.614 | 1743.4 | 78.40 | 4.509 | 193.48 | 84.306 | 102.69 |

*In a column, means followed by a common letter(s) are not significantly different at 5% level by LSD

DISCUSSION

Plant growth promoting effects of PGPR strains in different crops were clearly demonstrated (El-Hawary *et al.*, 2002; Wu *et al.*, 2005). Bacterial inoculants are able to increase plant growth and germination rate, improve seedling emergence, responses to external stress factors and protect plants from disease (Lugtenberg *et al.*, 2002).

This present investigation confirms the earlier studies. It revealed that under *in vitro* conditions, seed treatment with PGPR strains improved seed germination, seedling vigor, seedling emergence and seedling stand over the control. Similar improvement of seed germination parameters by rhizobacteria has been reported in other cereals such as sorghum (Raju *et al.*, 1999) and pearl millet (Niranjan *et al.*, 2003, 2004). The improvement in seed germination by PGPR was also found in work with wheat and sunflower (Shaukat *et al.*, 2006a, b), where it was found that some PGPR induced increases in seed emergence, in some cases achieving increases up to 100% greater than controls. These findings may be due to the increased synthesis of hormones like gibberellins, which would have triggered the activity of specific enzymes that promoted early germination, such as α -amylase, which have brought an increase in availability of starch assimilation. Beside, significant increase in seedling vigor would have occurred by better synthesis of auxins (Bharathi *et al.*, 2004).

In pot experiment, it was observed that inoculation with PGPR strains significantly promoted growth of seedling maize under different soil conditions. In general, inoculation resulted in early seedling growth and development. These results are similar with the findings of Dobbelaere *et al.* (2001, 2002), who assessed the inoculation effect of PGPR *Azospirillum brasilense* on growth of spring wheat. They observed that inoculated plants resulted in better germination, early development and flowering and also increase in dry weight of both the root system and the upper plant parts. Similarly, promotion in growth parameters and yields of various

crop plants in response to inoculation with PGPR were reported by Gravel *et al.* (2007), Kozdroja *et al.* (2004) and Shaharoon *et al.* (2006). Inoculation of maize seeds with *Azospirillum* strains compared with *Pseudomonas* strains under experiment conditions resulted in a more visible increase in shoot development, especially during the establishment of the plant. Khalid *et al.* (2004) showed that responses of wheat growth to inoculation with rhizobacteria depend on plant genotype and PGPR strains as well as environmental conditions.

Soil condition influenced growth promotion by bacterial strains. *Azospirillum brasilense* DSM 1690 had more effect on growth parameters in sterile soil compared with non sterile soil. Martinez-Toledo *et al.* (1988) showed that the numbers of *Azotobacter* decreased as plant growth continued in non-sterile agricultural soils, while the numbers of *Azotobacter* associated with maize roots grown in sterile agricultural soils remained similar to those of the original inoculums. In contrast, inoculation with other bacterial treatments had a more stimulating effect on growth of plants in non-sterile soil than sterile condition. Abbass and Okon (1993) hypothesized that IAA and other plant hormones were responsible for increased growth of canola, tomato (*Lycopersicon esculentum* Mill.) and wheat (*Triticum turgidum* L.) in non-sterile soil inoculated with *Azotobacter paspali*. Auxins produced by rhizobacteria can influence plants growth, including root development which improve uptake of essential nutrients thus increasing plant growth (Vikram *et al.*, 2007). This may imply rhizobacteria had more competitive ability to survive and affect the growth of inoculated plants in the presence of indigenous micro flora (Khalid *et al.*, 2004).

In the other hand, as suggested by Roesti *et al.* (2006) this result mean that inoculums of the PGPR strains on the seeds may have shifted the bacterial community equilibrium at early stages of plant growth and favoured for growth of beneficial populations.

In this study, inoculation of PGPR strains increased all parameters determined in field experiment. The positive

effects of PGPRs on the yield and growth of crops such as wheat (Ozturk *et al.*, 2003; Salanture *et al.*, 2006) maize (Egamberdiyeva, 2007) soybean (Cattelan *et al.*, 1999) and sugar beet (Cakmakc *et al.*, 2006) were explained by N₂ fixation ability, phosphate solubilizing capacity and phytohormones production.

The present experiment revealed that seed inoculation with all bacteria resulted in an increased plant height and leaf area (Table 2). Similar increases in plant height and leaf area were observed in different crops inoculated with *Pseudomonas*, *Azospirillum* and *Azotobacter* strains (Martinez-Toledo *et al.*, 1988; Shaukat *et al.*, 2006a, b; Siddiqui and Shaukat., 2002). Burd *et al.* (2000) reported that plant growth promoting rhizobacteria might enhance plant height and productivity by synthesizing phytohormones, increasing the local availability of nutrients, facilitating the uptake of nutrients by the plants decreasing heavy metal toxicity in the plants antagonizing plant pathogens.

Results have also showed that plants inoculated with PGPRs generally have higher seed dry weight than un-inoculated plants. The increases in seed dry weight were derived mainly from increase in 100 seed weight and number of seed per ear. This finding was supported by Yasari and Patwardhan (2007) reported that application of *Azotobacter* and *Azospirillum* strains increased canola yield (21.17%), pod per plant (16.05%), number of branches (11.78%) and weight of 1000 grain (2.92%).

The higher ear and shoot dry weight response to all inoculants compared to control clearly showed the beneficial role of these rhizobacteria. The enhancing effect of seed inoculation with rhizobacteria on shoot dry weight and yield of maize were reported by Pandey *et al.* (1998) and Shaharoon *et al.* (2006). Such an improvement might be attributed to N₂-fixing and phosphate solubilizing capacity of bacteria as well as the ability of these microorganisms to produce growth promoting substances (Bashan *et al.*, 2004; Vessey, 2003). In conclusion the results of this study suggest that simultaneous screening of rhizobacteria for growth and yield promotion under pot and field experiment is a good tool to select effective PGPR for biofertilizer development biotechnology.

ACKNOWLEDGMENTS

The researchers are thankful to Shahrood University of Technology and the Soil Microbiology Department, Soil and Water Research, Tehran for helping to carry out this study.

REFERENCES

- Abbass, Z. and Y. Okon, 1993. Plant growth promotion by *Azotobacter paspali* in the rhizosphere. *Soil Biol. Biochem.*, 25: 1075-1083.
- Abdul Baki, A.A. and J.D. Anderson, 1973. Vigor determination in soybean seed by multiple criteria. *Crop Sci.*, 13: 630-633.
- Ahmad F., I. Ahmad and M.S. Khan, 2008. Screening of free-living rhizospheric bacteria for their multiple plant growth promoting activities. *Microbiol. Res.*, 168: 173-181.
- Asghar, H.N., Z.A. Zahir, M. Arshad and A. Khaliq, 2002. Relationship between *in vitro* production of auxins by rhizobacteria and their growth promoting activities in *Brassica juncea* L. *Bio. Fertility Soil*, 35: 231-237.
- Bashan, Y., Y. Ream, H.L. Levanony and A. Sade, 1989. Non-specific response in plant growth yields and root colonization of non-cereal crop plants to inoculation with *Azospirillum brasilense* cd. *Can. J. Bot.*, 67: 1317-1324.
- Bashan, Y., G. Holguin and L.E. De-Bashan, 2004. *Azospirillum*-plant relationships: physiological, molecular, agricultural and environmental advances (1997-2003). *Can. J. Microbiol.*, 50: 521-577.
- Bharathi, R., R. Vivekananthan, S. Harish, A. Ramanathan and R. Samiyappan, 2004. Rhizobacteria-based bio-formulations for the management of fruit rot infection in chillies. *Crop Prot.*, 23: 835-843.
- Biswas, J.C., L.K. Ladha and F.B. Dazzo. 2000. Rhizobia inoculation improves nutrient uptake and growth of lowland rice. *Soil Sci. Soc. Am. J.*, 64: 1644-1650.
- Burd, G.I., D.G. Dixon and B.R. Glick, 2000. Plant growth promoting rhizobacteria that decrease heavy metal toxicity in plants. *Can. J. Microbiol.*, 33: 237-245.
- Cakmakc, R.I., D.F. Ayдын and A.F. Sahin, 2006. Growth promotion of plants by plant growth-promoting rhizobacteria under greenhouse and two different field soil conditions. *Soil Biol. Biochem.*, 38: 1482-1487.
- Cattelan, A.J., P.G. Hartel and J.J. Fuhrmann, 1999. Screening for plant growth-promoting rhizobacteria to promote early soybean growth. *Soil Sci. Soc. Am. J.*, 63: 1670-1680.
- De Freitas, J.R. and J.J. Gemida, 1992. Growth promotion of winter wheat by fluorescent pseudomonads under growth chamber conditions. *Soil Biol. Biochem.*, 24: 1127-1135.
- Dobbelaere, S., A. Croonenborghs, A. Thys, D. Ptacek and J. Vanderleyden *et al.*, 2001. Response of agronomically important crops to inoculation with *Azospirillum*. *Aust. J. Plant Physiol.*, 28: 871-879.

- Dobbelaere, S., A. Croonenborghs, A. Thys, D. Ptacek, Y. Okon and J. Vanderleyden, 2002. Effect of inoculation with wild type *Azospirillum brasilense* and A. irakense strains on development and nitrogen uptake of spring wheat and grain maize. Biol. Fertil. Soils, 36: 284-297.
- Egamberdiyeva, D., 2007. The effect of plant growth promoting bacteria on growth and nutrient uptake of maize in two different soils. Applied Soil Ecol., 36: 184-189.
- El-Hawary, M.I., I. El-Hawary, Fatma, A.M. El-Ghamry and E. El-Naggar 2002. Effect of Application of Biofertilizer on the Yield and NPK Uptake of Some Wheat Genotypes as Affected by the Biological Properties of Soil Pak. J. Biol. Sci., 5: 1181-1185.
- Frommel, M.I., J. Nowak and G. Lazarovits, 1993. Treatment of potato tubers with a growth promoting *Pseudomonas* sp. Plant growth responses and bacterium distribution in the rhizosphere. Plant Soil, 150: 51-60.
- Glick, B.R., L. Changping, G. Sibdas and E.B. Dumbroff, 1997. Early development of canola seedlings in the presence of the plant growth-promoting rhizobacterium *Pseudomonas putida* GR12-2. Soil Biol. Biochem., 29: 1233-1239.
- Gravel, V., H. Antoun and R.J. Tweddell, 2007. Growth stimulation and fruit yield improvement of greenhouse tomato plants by inoculation with *Pseudomonas putida* or *Trichoderma atroviride*: Possible role of Indole Acetic Acid (IAA). Soil Biol. Biochem., 39: 1968-1977.
- Jeun, Y.C., K.S. Park, C.H. Kim, W.D. Fowler and J.W. Kloepper, 2004. Cytological observations of cucumber plants during induced resistance elicited by rhizobacteria. Biol. Control, 29: 34-42.
- Khalid, A., M. Arshad and Z.A. Zahir, 2004. Screening plant growth-promoting rhizobacteria for improving growth and yield of wheat. J. Applied Microbiol., 29: 473-480.
- Kloepper, J.W. and C.J. Beauchamp, 1992. A review of issues related to measuring of plant roots by bacteria. Can. J. Microbiol., 38: 1219-1232.
- Kozdroja, J., J.T. Trevors and J.D. Van Elsas, 2004. Influence of introduced potential biocontrol agents on maize seedling growth and bacterial community structure in the rhizosphere. Soil Biol. Biochem., 36: 1775-1784.
- Lugtenberg, B., T. Chin-A-Woeng and G. Bloemberg, 2002. Microbe-plant interactions: Principles and mechanisms. Antonie van Leeuwenhoek, 81: 373-383.
- Martinez-Toledo, M.V., J. Gonzalez-Lopez, T. De la Rubia, J. Moreno and A. Ramos-ormenzana, 1988. Effect of inoculation with *Azotobacter chroococcum* on nitrogenase activity of *Zea mays* roots grown in agricultural soils under aseptic and non-sterile conditions. Biol. Fertility Soils, 6: 170-173.
- Mrkovacki, N. and V. Milic, 2001. Use of *Azotobacter chroococcum* as potentially useful in agricultural application. Ann. Microbiol., 51: 145-158.
- Niranjan, S.R., S.A. Deepak, P. Basavaraju, H.S. Shetty, M.S. Reddy and J.W. Kloepper, 2003. Comparative performance of formulations of plant growth promoting rhizobacteria in growth promotion and suppression of downy mildew in pearl millet. Crop Protect., 22: 579-588.
- Niranjan, S.R., N.P. Shetty and H.S. Shetty, 2004. Seed bio priming with *Pseudomonas fluorescens* isolates enhances growth of pearl millet plants and induces resistance against downy mildew. J. Pest Manage., 50: 41-48.
- Ozturk, A., O. Caglar and F. Sahin, 2003. Yield response of wheat and barley to inoculation of plant growth promoting rhizobacteria at various levels of nitrogen fertilization. J. Plant Nutr. Soil Sci., 166: 262-266.
- Pandy, A., E. Sharma and L.K.S. Plani, 1998. Influence of bacterial inoculation on maize in upland farming systems of the sikkim himalaya. Soil Biol. Biochem., 30: 379-384.
- Raju, N.S., S. R. Niranjana, G.R. Janardhana, H.S. Prakash, H.S. Shetty and S.B. Mathur, 1999. Improvement of seed quality and field emergence of *Fusarium moniliforme* infected sorghum seeds using biological agents. J. Sci. Food Agric., 79: 206-212.
- Ramamoorthy, V., T. Raguchander and R. Samiyappan, 2002. Enhancing resistance of tomato and hot pepper to Pythium diseases by seed treatment with fluorescent pseudomonads. Eur. J. Plant Pathol., 108: 429-441.
- Roesti, D., R. Gaur, B.N. Johri, G. Imfeld, S. Sharma, K. Kawaljeet and M. Aragno, 2006. Plant growth stage, fertilizer management and bio-inoculation of Arbuscular mycorrhizal fungi and plant growth promoting rhizobacteria affect the rhizobacterial community structure in rain-fed wheat fields. Soil Biol. Biochem., 38: 1111-1120.
- Ruget, F., R. Bonhomme and M. Chartier, 1996. Estimation simple de la surface foliaire de plantes de maïs en croissance. Agronomie, 16: 553-562.
- Salantur, A., A. Ozturk and S. Akten, 2006. Growth and yield response of spring wheat (*Triticum aestivum* L.) to inoculation with rhizobacteria. Plant Soil Environ., 52: 111-118.

- Shaharoon, B., M. Arshad, Z.A. Zahir and A. Khalid, 2006. Performance of *Pseudomonas* sp. containing ACC-deaminase for improving growth and yield of maize (*Zea mays* L.) in the presence of nitrogenous fertilizer. *Soil Biol. Biochem.*, 38: 2971-2975.
- Shaukat, K., S. Affrasayab and S. Hasnain, 2006a. Growth responses of *Helianthus annuus* to plant growth promoting rhizobacteria used as a biofertilizer. *J. Agric. Res.*, 1: 573-581.
- Shaukat, K., S. Affrasayab and S. Hasnain, 2006b. Growth responses of *Triticum aestivum* to plant growth promoting rhizobacteria used as a biofertilizer. *Res. J. Microbiol.*, 1: 330-338.
- Siddiqui, I.A. and S.S. Shaukat, 2002. Mixtures of plant disease suppressive bacteria enhance biological control of multiple tomato pathogens. *Biol. Fertility Soil*, 36: 260-268.
- Vessey, J.K., 2003. Plant growth promoting rhizobacteria as biofertilizers. *Plant Soil*, 255: 571-586.
- Vikram, A., 2007. Efficacy of phosphate solubilizing bacteria isolated from vertisols on growth and yield parameters of sorghum. *Res. J. Microbiol.*, 2: 550-559.
- Wu, S.C., Z.H. Cao, Z.G. Li, K.C. Cheung and M.H. Wong, 2005. Effects of biofertilizer containing N-fixer, P and K solubilizers and AM fungi on maize growth: A greenhouse trial. *Geoderma*, 125: 155-166.
- Yasari, E. and A.M. Patwardhan, 2007. Effects of *Azotobacter* and *Azospirillum* inoculations and chemical fertilizers on growth and productivity of Canola. *Asian J. Plant Sci.*, 6: 77-82.
- Zhang, F., N. Dashti, H. Hynes and D.L. Smith, 1996. Plant growth promoting rhizobacteria and soybean [*Glycine max* (L.) Merr.] nodulation and nitrogen fixation at suboptimal root zone temperatures. *Ann. Bot.*, 77: 453-459.